



ELSEVIER

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

Spectrochimica Acta Part A 70 (2008) 674–681

SPECTROCHIMICA  
ACTA  
PART A[www.elsevier.com/locate/saa](http://www.elsevier.com/locate/saa)

## Model-based analysis for kinetic complexation study of Pizda and Cu(II)

M. Vosough<sup>a</sup>, M. Maeder<sup>b</sup>, M. Jalali-Heravi<sup>c,\*</sup>, S.E. Norman<sup>b</sup><sup>a</sup> Chemistry and Chemical Engineering Research Center of Iran, Tehran, Iran<sup>b</sup> Department of Chemistry, University of Newcastle, Callaghan, NSW, Australia<sup>c</sup> Department of Chemistry, Sharif University of Technology, P.O. Box 11365-9516, Tehran, Iran


Received 19 March 2007; received in revised form 25 August 2007; accepted 27 August 2007

## Abstract

In the present work, the multivariate kinetic complexation of a new synthesized ligand, 1-(2'-hydroxyl cyclohexyl)-3'-[aminopropyl]-4-[3'-aminopropyl]piperazine (Pizda) and Cu<sup>2+</sup> in 50% ethanol–water solution is investigated using the UV–vis stopped-flow technique and state-of-the-art multi-wavelength numerical analysis. Model-based least squares fitting analysis or hard modeling is a specific part of chemometrics which is based on mathematical relationships for describing the measurements. Some recent developments include the incorporation of the effects of non-ideal experimental conditions into the fitting algorithm so it can substantially simplify experimental procedures. In this study no buffers are required because pH changes are taken into computations. Some 21 multi-wavelength kinetic measurements, taken at various initial concentrations of [H<sup>+</sup>] were analyzed globally, i.e. simultaneously applying an all inclusive reaction mechanism and a common set of species spectra. Using numerical analysis, the pH of the experimental solutions was allowed to vary as a consequence of the proceeding reactions. This enabled the complete kinetic analysis of the formation and dissociation of Cu(Pizda)<sup>n+</sup>. Here protonation equilibria have been directly incorporated into the rate law, so thus variable pH values have been allowed during each measurement. Using the independently estimated stability constants (from spectrophotometric and potentiometric measurements) for the Cu(Pizda)<sup>n+</sup> complexes, a total of six rate constants and one protonation constant could be elucidated. The results of the analysis include the concentration distribution and spectra of all chemical species involved in the reaction. A low standard deviation and residual profiles obtained validate the results.


© 2007 Elsevier B.V. All rights reserved.

**Keywords:** Hard modeling; Stopped-flow measurements; Global analysis; Pizda





Самая маленькая разновеска для аналитических весов.



Рекомендации для пользователей Windows.



(фотография ulibnis1710.h18.ru)