


DOI: 10.1002/anie.200501276

Approach to the Blues: A Highly Flexible Route to the Azulenes**

Sébastien Carret, Aurélien Blanc, Yoann Coquerel,
Mikaël Berthod, Andrew E. Greene, and
Jean-Pierre Deprés*

*Dedicated to Dr. Jean-Louis Luche
on the occasion of his retirement*

Nearly a century and a half ago, Septimus Piesse applied the descriptive name azulene to azure-blue distillates from various sources, such as chamomile, yarrow, and wormwood.^[1] However, it was only in 1937 that Placidus Plattner and Alexander Pfau successfully carried out the first synthesis of azulene (**1**),^[2] the appellation given to the parent compound




of the family of bicyclo[5.3.0]decapentaenes, which are now collectively referred to as the azulenes. Since this historical event, additional syntheses of the parent azulene and preparations of various other azulenes, such as guaiazulene (**2**), which is the archetype of the major azulene class, have been reported.^[3] Although the approaches to these fascinating and electronically unique compounds are often ingenious, they suffer for the most part from being excessively long, low yielding, or lacking in generality. Many substitution patterns of the azulenes are still difficult to access, if they can be accessed at all.^[3]

We have found that chlorotrienones **5a** and **5b** are readily prepared from cycloheptatriene and 7-methylcycloheptatriene, respectively, through a doubly regioselective cycloaddition of dichloroketene, a regioselective ring expansion with ethereal diazomethane, and dehydrochlorination in dimethylformamide (DMF; Scheme 1).^[4]

[*] S. Carret, A. Blanc, Y. Coquerel, M. Berthod, Dr. A. E. Greene, Prof. J.-P. Deprés
Chimie Recherche (LEDSS)
Université Joseph Fourier
B.P. 53X, 38041 Grenoble, Cedex 9 (France)
Fax: (+33) 4-7651-4494
E-mail: jean-pierre.depres@ujf-grenoble.fr

[**] We thank Prof. P. Dumy for his interest in our research, the Research Ministry for fellowship awards to S.C. and Y.C., and the Université Joseph Fourier and the CNRS (UMR 5616, FR 2607) for financial support.




Scheme 1. Preparation of chlorotrienones **5a** and **5b**: a) Zn/Cu, POCl₃, CCl₃COCl, Et₂O, 20 °C, 20 h; b) CH₂N₂, MeOH/Et₂O (5:95), 0→20 °C, 35 min; c) DMF, 20 °C, 12 h (44–45 % yield, 3 steps).

As other C7-substituted cycloheptatrienes and diazoalkane reagents are also readily available^[4a,5] for use in this sequence, it appeared to us that a broad approach to the azulenes might now be possible if 1) the oxygen atom at C3 could be excised, 2) the fused rings could be dehydrogenated to aromaticity, 3) the chlorine atom at C4 could be replaced, and 4) additional substituents could be regioselectively introduced into the trienones, particularly at C7. Herein, we disclose an efficient and exceptionally versatile preparation of azulenes, which are compounds of considerable interest for use in cosmetics,^[6a] pharmaceutical preparations,^[6b] and new molecular materials,^[6c] through effective resolution of these concerns.

Azulene itself was targeted initially so that the first three critical points could be addressed at the outset. It was pleasing to find, after a number of failed attempts, that sequential treatment of **5a** with NaBH₄/CeCl₃, the Burgess reagent,^[7] and *p*-chloranil could transform **5a** into chloroazulene **6a** in 45 % overall yield (Scheme 2).^[8] Azulene could then be obtained from **6a** in 92 % yield through modification of the reduction recently described by Rahaim and Maleczka.^[9] Analogously, 4-methylazulene (**7**)^[2,10] was readily prepared from **5b** in a slightly higher overall yield.


As most naturally occurring azulenes have methyl groups at C1 and C4, the discovery that **6b** could also be methylated by reaction with methylboronic acid, dpdb, and K₃PO₄^[11] to give 1,4-dimethylazulene (**8**)^[12a] in 98 % yield was welcomed



Scheme 2. Preparation of azulene (**1**), 4-methylazulene (**7**), and 1,4-dimethylazulene (**8**): a) NaBH₄, MeOH/CeCl₃, 0 °C, 1.5 h (from **5a**: 96 %; from **5b**: 98 %); b) the Burgess reagent, THF, 0→20 °C, 1 h; *p*-chloranil, 20 °C, 24 h (from **6a**: 47 %; from **6b**: 52 %); c) PMHS, Pd(OAc)₂, dpdb, K₃PO₄, THF, 80 °C, 12 h (**1**: 92 %; **7**: 91 %); d) MeB(OH)₂, Pd(OAc)₂, dpdb, K₃PO₄, PhMe, 100 °C, 24 h (98 %). Burgess reagent = (methoxycarbonylsulfonyl)triethylammonium hydroxide inner salt, *p*-chloranil = tetrachloro-1,4-benzoquinone, PMHS = poly(methylhydrosiloxane), dpdb = dicyclohexylphosphano-2',6'-dimethoxybiphenyl.

and augured well for the application of this approach to many other naturally occurring azulenes.^[12b]

Considerable study was required to resolve point (4) satisfactorily. Although several different types of organometallic reagents did indeed add regio- (and stereo-)selectively to **5b** at C7, the yields were modest and, furthermore, aromatization of the resultant dienones proved difficult to achieve efficiently. We were, therefore, pleased to find that organozinc reagents, in the presence of CuOTf (cat.) and TMSCl,^[13] were highly effective in transforming **5b** into the corresponding 1,6-conjugate addition products: Me, 91 %; Et, 98 %; *i*Pr, 71 %. Even better, it was discovered that simply by adding PhSeCl to the reaction mixture and then effecting oxidative (H₂O₂) elimination, the chlorotrienone system could be smoothly regenerated (Scheme 3).^[14] Thus, chloro-




Scheme 3. Preparation of chamazulene (**10**), 7-isopropyl-4-methylazulene (**12**), and guaiazulene (**2**): a) 1. Et₂Zn, CuOTf, HMPA, TMSCl, −50 °C, 6 h; 2. PhSeCl, −50→0 °C, 2 h; b) H₂O₂, pyridine, 0 °C, 1 h (**9**: 71 %, 2 steps; **11**: 62 %, 2 steps); c) NaBH₄, MeOH/CeCl₃, 0 °C, 1.5 h (58–59 %); e) MeB(OH)₂, Pd(OAc)₂, dpdb, K₃PO₄, PhMe, 100 °C, 24 h (**10**: 89 %; **2**: 94 %); f) 1. *i*Pr₂Zn, CuOTf, HMPA, TMSCl, −100 °C, 6 h; 2. PhSeCl, −50→0 °C, 2 h; g) PMHS, Pd(OAc)₂, dpdb, K₃PO₄, THF, 80 °C, 12 h (95 %). Tf = trifluoromethanesulfonyl, HMPA = hexamethylphosphoramide, TMSCl = trimethylsilyl chloride.

trienones **9** and **11** were prepared in 71 and 62 % yields, respectively. The former, through application of the above chemistry, provided chamazulene (**10**), and the latter, 7-isopropyl-4-methylazulene (**12**) and guaiazulene (**2**).^[15]


Ketene acetals can also be employed for 1,6-conjugate addition, thus adding to the synthetic possibilities: the treatment of **5b** with the *tert*-butyldimethylsilyl (TBDMS) ketene acetal, derived from methyl propionate, in the presence of lithium perchlorate^[16] readily generated the 1,6-conjugate addition product, which on successive treatment with PhSeCl and H₂O₂ afforded chlorotrienone **13** (Scheme 4). Aromatization, methylation, and saponification then provided chamazulenecarboxylic acid (**14**).^[17]

The attractiveness of the present approach resides, in part, in the diversity of structures that can readily be accessed with total regiocontrol, a feature that few other approaches enjoy. To further illustrate this point, regiochemically pure 7-isopropyl-2,4-dimethylazulene (**15**)^[18a] (a naturally occurring isomer of guaiazulene (**2**)),^[15b] 1-chloro-3-methylazulene (**16a**),^[18b] and the phenylazulenes **16b**^[18c] and **16c**^[18d] have



Scheme 4. Preparation of (±)-chamazulenecarboxylic acid (**14**):

a) 1. MeCH=C(OMe)OTBDMS, LiClO₄, CH₂Cl₂, 20 °C, 12 h; 2. PhSeCl, –50 → 0 °C, 2 h; b) H₂O₂, pyridine, 0 °C, 1 h (46%, 2 steps); c) NaBH₄, MeOH/CeCl₃, 0 °C, 1.5 h (92%); d) 1. Burgess reagent, THF, 0 °C, 1 h; 2. *p*-chloranil, 20 °C, 24 h (55%); e) MeB(OH)₂, Pd(OAc)₂, dpdb, K₃PO₄, PhMe, 100 °C, 24 h (99%); f) LiOH, THF, H₂O, 20 °C, 12 h (94%).



also been efficiently prepared, thereby demonstrating four distinctly different substitution options as well. Finally, acetoxy and aza derivatives **17a**^[18e] and **17b**^[18f] are examples of novel azulenes that are readily prepared with this methodology.

In conclusion, modern chemistry has been applied to the long-standing problem of azulene synthesis and an efficient, highly flexible approach has resulted. Our approach allows controlled access to a wide variety of substituents and substitution arrays and should prove to be among the most broadly useful methods to prepare these important compounds.

Experimental Section

Experimental procedure for the preparation of **6b** from **5b**: NaBH₄ (43 mg, 1.13 mmol) was added to a stirred solution of **5b** (200 mg, 1.03 mmol) in MeOH/CeCl₃ (2.80 mL, 0.40 M, 1.12 mmol) at 0 °C. The reaction mixture was stirred for 1.5 h at 0 °C and then treated with a saturated solution of aqueous NaH₂PO₄. The crude product was isolated with EtOAc/pentane (1:1) in the usual manner and purified by column chromatography on dry silica gel with diethyl ether/pentane (40:60) to afford 198 mg (98%) of the *trans* alcohol as a white solid. M.p. 67–68 °C; ¹H NMR (300 MHz, CDCl₃): δ = 1.17 (d, *J* = 7.1 Hz, 3H), 1.38–1.48 (m, 1H), 2.16–2.24 (m, 2H), 2.33 (pseudo q, *J* = 7.3 Hz, 1H), 2.68–2.77 (m, 1H), 4.57 (pseudo t, *J* = 7.2 Hz, 1H), 5.66 (dd, *J* = 11.4, 3.3 Hz, 1H), 5.85–5.94 (m, 1H), 6.01 (dd, *J* = 11.5, 6.9 Hz, 1H), 6.51 ppm (d, *J* = 11.5 Hz, 1H); ¹³C NMR (75.5 MHz, CDCl₃): δ = 19.2, 38.9, 40.3, 46.6, 74.6, 123.7, 125.3, 127.7, 133.6, 139.6, 141.5 ppm; FTIR: $\tilde{\nu}$ = 3360, 3016, 1700 cm^{–1}. A solution of the Burgess reagent (347 mg, 1.46 mmol) in dry THF (5 mL) was added to a stirred solution of the above alcohol (198 mg, 1.01 mmol) in dry THF (6.0 mL) at 0 °C. The reaction mixture was allowed to warm to 20 °C and stirred for 1 h, whereupon *p*-chloranil (745 mg, 3.03 mmol) was added and the resulting mixture was stirred for a further 24 h. The crude product was isolated with pentane in the usual way and purified by column chromatography on dry silica gel with pentane to give 92 mg (52%) of chloroazulene **6b** as a blue solid. M.p. 36 °C; ¹H NMR (300 MHz, CDCl₃): δ = 2.86 (s, 3H), 7.13 (d, *J* = 10.3 Hz,

1H), 7.14 (pseudo t (dd), *J* = 9.5 Hz, 1H), 7.30 (d, *J* = 4.2 Hz, 1H), 7.53 (pseudo t (dd), *J* = 10.3 Hz, 1H), 7.67 (d, *J* = 4.2, 1H), 8.40 ppm (d, *J* = 9.5 Hz, 1H); ¹³C NMR (75.5 MHz, CDCl₃): δ = 24.1, 113.4, 117.1, 122.0, 127.2, 133.3, 133.5, 134.8, 136.7, 137.8, 148.0 ppm; FTIR: $\tilde{\nu}$ = 3085, 3021, 2957, 2923, 2853, 1591, 1560, 1487, 1419, 1389, 1358, 906, 772, 743 cm^{–1}; MS (DCI): *m/z*: 177 [M+H]⁺; HRMS calcd for [C₁₁H₉Cl+H]⁺: 177.0471; found: 177.0484; elemental analysis (%) calcd for C₁₁H₉Cl: C 74.79, H 5.14; found: C 74.84, H 5.18.

Data for selected compounds: **2**: M.p. 31–32 °C; ¹H NMR (300 MHz, CDCl₃): δ = 1.36 (d, *J* = 6.9 Hz, 6H), 2.66 (s, 3H), 2.83 (s, 3H), 3.08 (sept, *J* = 6.9 Hz, 1H), 7.01 (d, *J* = 10.7 Hz, 1H), 7.22 (d, *J* = 3.7 Hz, 1H), 7.42 (dd, *J* = 10.7, 1.8 Hz, 1H), 7.62 (d, *J* = 3.7 Hz, 1H), 8.19 ppm (d, *J* = 1.8 Hz, 1H); ¹³C NMR (75.5 MHz, CDCl₃): δ = 13.0, 24.2, 24.9, 38.4, 112.9, 125.2, 125.3, 133.5, 135.0, 136.2, 136.4, 137.4, 140.0, 144.4 ppm; FTIR: $\tilde{\nu}$ = 3095, 3064, 2958, 2924, 2854, 1554, 1527, 1462, 1420, 1387, 1367, 772 cm^{–1}; MS (DCI): *m/z*: 199 [M+H]⁺; HRMS calcd for [C₁₅H₁₉+H]⁺: 199.1487; found: 199.1482.

10: ¹H NMR (300 MHz, CDCl₃): δ = 1.34 (t, *J* = 7.6 Hz, 3H), 2.65 (s, 3H), 2.82 (s, 3H), 2.83 (q, *J* = 7.6 Hz, 2H), 6.98 (d, *J* = 10.5 Hz, 1H), 7.21 (br s, 1H), 7.38 (dd, *J* = 10.5, 1.9 Hz, 1H), 7.61 (br s, 1H), 8.15 ppm (d, *J* = 1.9 Hz, 1H); ¹³C NMR (75.5 MHz, CDCl₃): δ = 13.0, 17.5, 24.2, 34.0, 112.9, 125.1, 125.2, 134.8, 135.9, 136.3, 136.4, 136.5, 137.5, 144.4 ppm; FTIR: $\tilde{\nu}$ = 3098, 3063, 2960, 2926, 2866, 1555, 1561, 1526, 1452, 1422, 1364, 772 cm^{–1}; MS (DCI): *m/z*: 185 [M+H]⁺; HRMS calcd for [C₁₄H₁₆+H]⁺: 185.1330; found: 185.1334; elemental analysis (%) calcd for C₁₄H₁₆: C 91.25, H 8.75; found: C 91.44, H 8.86.

12: ¹H NMR (300 MHz, CDCl₃): δ = 1.35 (d, *J* = 6.8 Hz, 6H), 2.88 (s, 3H), 3.07 (sept, *J* = 6.8 Hz, 1H), 7.11 (d, *J* = 10.6 Hz, 1H), 7.29 (d, *J* = 3.8 Hz, 1H), 7.31 (d, *J* = 3.8 Hz, 1H), 7.46 (dd, *J* = 10.6, 2.0 Hz, 1H), 7.80 (pseudo t (dd), *J* = 3.8 Hz, 1H), 8.31 ppm (d, *J* = 2.0 Hz, 1H); ¹³C NMR (75.5 MHz, CDCl₃): δ = 24.3, 24.7, 38.2, 114.7, 118.0, 126.0, 135.3, 135.4, 136.6, 137.4, 140.2, 141.8, 145.2 ppm; FTIR: $\tilde{\nu}$ = 3093, 3064, 2958, 2924, 1556, 1529, 1461, 1422, 1389, 1362, 749 cm^{–1}; MS (DCI): *m/z*: 185 [M+H]⁺; HRMS calcd for [C₁₄H₁₆]⁺: 184.1252; found: 184.1259.

14: M.p. 86–87 °C; ¹H NMR (300 MHz, CDCl₃): δ = 1.61 (d, *J* = 7.2 Hz, 3H), 2.65 (s, 3H), 2.83 (s, 3H), 3.88 (q, *J* = 7.2 Hz, 1H), 7.00 (d, *J* = 10.7 Hz, 1H), 7.28 (d, *J* = 3.8 Hz, 1H), 7.44 (dd, *J* = 10.7, 1.9 Hz, 1H), 7.64 (d, *J* = 3.8 Hz, 1H), 8.22 ppm (d, *J* = 1.9 Hz, 1H); ¹³C NMR (75.5 MHz, CDCl₃): δ = 13.0, 19.2, 24.2, 48.8, 114.3, 125.1, 127.1, 130.9, 133.9, 135.7, 135.8, 136.9, 137.6, 145.6 ppm; FTIR: $\tilde{\nu}$ = 3585, 3103, 3065, 2961, 2923, 2853, 1704, 1556, 1454, 1261, 1023, 774 cm^{–1}; MS (DCI): *m/z*: 229 [M+H]⁺.

Received: April 12, 2005

Published online: July 13, 2005

Keywords: aromatic compounds · azulenes · cycloaddition · natural products · synthesis design

- [1] S. Piesse, C. R. *Hebd. Seances Acad. Sci.* **1863**, 57, 1016.
- [2] P. A. Plattner, A. S. Pfau, *Helv. Chim. Acta* **1937**, 20, 224–232; see, also: H.-J. Hansen, *Chimia* **1996**, 50, 489–496.
- [3] For reviews on the azulenes, see: a) M. Gordon, *Chem. Rev.* **1952**, 52, 127–200; b) V. B. Mochalin, Y. N. Porshnev, *Russ. Chem. Rev.* **1977**, 46, 530–547; c) K.-P. Zeller in *Methoden der Organischen Chemie (Houben-Weyl)*, Vol. V/2c (Ed.: H. Kropf), Georg Thieme, Stuttgart, **1985**, pp. 127–418; for a recent synthetic approach, see: d) A. L. Crombie, J. L. Kane, Jr., K. M. Shea, R. L. Danheiser, *J. Org. Chem.* **2004**, 69, 8652–8667.
- [4] a) Y. Coquerel, PhD Thesis, University of Grenoble, **2001**; b) Y. Coquerel, A. Blanc, J.-P. Depr s, A. E. Greene, M.-T. Averbuch-Pouchot, C. Philouze, A. Durif, *Acta Crystallogr. Sect. C* **2000**, 56, 1480–1481; c) Y. Coquerel, A. E. Greene, J.-P. Depr s, *Org. Lett.* **2003**, 5, 4453–4455; see, also: d) R. Yokoyama, S. Ito, M.

- Watanabe, N. Harada, C. Kabuto, N. Morita, *J. Chem. Soc. Perkin Trans. 1* **2001**, 2257–2261.
- [5] H. Günther, M. Görliitz, H. H. Hinrichs, *Tetrahedron* **1968**, *24*, 5665–5676; tropylium tetrafluoroborate is now commercially available.
- [6] a) C. S. Slavtcheff, S. R. Barrow, V. D. Kanga, M. C. Cheney, A. Znaiden (Unilever PLC, UK) WO9503779, **1995** [*Chem. Abstr.* **1995**, *123*, 17488w]; b) T. Yanagisawa, S. Wakabayashi, T. Tomiyama, M. Yasunami, K. Takase, *Chem. Pharm. Bull.* **1988**, *36*, 641–647; A. E. Asato, A. Peng, M. Z. Hossain, T. Mirzadegan, J. S. Bertram, *J. Med. Chem.* **1993**, *36*, 3137–3147; W.-G. Friebe, J. Dickhaut, R. Haag, H. Leinert, F. Grams (Roche Diagnostics GmbH, Germany), EP922702, **1999** [*Chem. Abstr.* **1999**, *131*, 58665k]; T. Toyama, I. Toyama, M. Yokota (Kotobuki Seiyaku Co. Ltd., Japan), JP2000007611, **2000** [*Chem. Abstr.* **2000**, *132*, 64075e]; B.-C. Hong, Y.-F. Jiang, E. S. Kumar, *Bioorg. Med. Chem. Lett.* **2001**, *11*, 1981–1984; D. A. Becker, J. J. Ley, L. Echegoyen, R. Alvarado, *J. Am. Chem. Soc.* **2002**, *124*, 4678–4684; H. Wakabayashi, K. Hashiba, K. Yokoyama, K. Hashimoto, H. Kikuchi, H. Nishikawa, T. Kurihara, K. Satoh, S. Shioda, S. Sato, S. Kusano, H. Nakashima, N. Motohashi, H. Sakagami, *Anticancer Res.* **2003**, *23*, 4747–4755; c) L. Cristian, I. Sasaki, P. G. Lacroix, B. Donnadieu, I. Asselberghs, K. Clays, A. C. Razus, *Chem. Mater.* **2004**, *16*, 3543–3551, and references therein.
- [7] E. M. Burgess, H. R. Penton, Jr., E. A. Taylor, *J. Org. Chem.* **1973**, *38*, 26–31.
- [8] Application of the Shapiro reaction, a) chlorotrienone, TsNHNH₂; b) MeLi, failed to produce the expected dihydroazulene; alternative allylic alcohol treatments, Al₂O₃, Pd/C; *p*-chloranil, Δ; HCl, Δ; Martin sulfurane reagent, were not fruitful.
- [9] R. J. Rahaim, Jr., R. E. Maleczka, Jr., *Tetrahedron Lett.* **2002**, *43*, 8823–8826; the reported conditions gave **1** in only 54% yield.
- [10] P. A. Plattner, E. Heilbronner, A. Fürst, *Helv. Chim. Acta* **1947**, *30*, 1100–1105.
- [11] S. D. Walker, T. E. Barder, J. R. Martinelli, S. L. Buchwald, *Angew. Chem.* **2004**, *116*, 1907–1912; *Angew. Chem. Int. Ed.* **2004**, *43*, 1871–1876.
- [12] a) P. A. Plattner, J. Wyss, *Helv. Chim. Acta* **1940**, *23*, 907–911; b) 1,4-dimethylazulene is naturally occurring, see: D. Meuche, S. Huneck, *Chem. Ber.* **1966**, *99*, 2669–2674; U. Siegel, R. Mues, R. Dönig, T. Eicher, M. Blechschmidt, H. Becker, *Phytochemistry* **1992**, *31*, 1671–1678.
- [13] M. Kitamura, T. Miki, K. Nakano, R. Noyori, *Bull. Chem. Soc. Jpn.* **2000**, *73*, 999–1014.
- [14] It should be noted that the aromatization of the azulenes, not unexpectedly, is directly dependent on the degree of unsaturation in the substrates; see reference [3b].
- [15] Chamazulene is an artifact from various sesquiterpene oils; 7-isopropyl-4-methylazulene and guaiazulene are found naturally in different sources, see: a) T. K. Devon, A. I. Scott in *Handbook of Naturally Occurring Compounds*, Vol. 2, Academic Press, New York, **1972**; b) *Dictionary of Terpenoids*, Vol. 2 (Eds.: J. D. Connolly, R. A. Hill), Chapman and Hall, New York, **1991**, and references therein.
- [16] C. S. Wilcox, R. E. Babston, *J. Org. Chem.* **1984**, *49*, 1451–1453; M. T. Reetz, D. D. A. Fox, *Tetrahedron Lett.* **1993**, *34*, 1119–1122.
- [17] E. Stahl, *Naturwissenschaften* **1954**, *41*, 257; E. Stahl, *Chem. Ber.* **1954**, *87*, 505–507; an earlier synthesis has been reported in a patent, see: Robugen GmbH Pharmazeutische Fabrik Esslingen A.N., Germany DE 10065683, **2001** [*Chem. Abstr.* **2001**, *135*, 76647t].
- [18] a) Prepared by sequential treatment of **11** with MeMgBr, the Burgess reagent, *p*-chloranil, and PMHS/Pd(OAc)₂; b) synthesized by treatment of **4a** with MeCHN₂ rather than CH₂N₂,

followed by DMF, NaBH₄/CeCl₃, the Burgess reagent, and *p*-chloranil; c) obtained from 7-phenylcycloheptatriene; see references [4a,5]; d) prepared from **6b** by using phenylboronic acid rather than methylboronic acid (98% yield); e) obtained by heating **5b** with Ac₂O and Pd/C in toluene (50% yield), for several related examples, see reference [4a]; f) synthesized from cyclobutanone **4b** by sequential treatment with *O*-mesitylene-sulfonylhydroxylamine, [Me₃O][BF₄], and DMF; for other examples of aza azulenes, see: M. Kitamura, S. Chiba, O. Saku, K. Narasaka, *Chem. Lett.* **2002**, 606–607.