

TOPICS IN  
ORGANOMETALLIC CHEMISTRY

34

# Iridium Catalysis

 Springer

**34**

**Topics in Organometallic Chemistry**

**Editorial Board:**

**M. Beller • J. M. Brown • P. H. Dixneuf**

**A. Fürstner • L. S. Hegedus • P. Hofmann**

**T. Ikariya • L. A. Oro • M. Reetz • Q.-L. Zhou**

# Topics in Organometallic Chemistry

## Recently Published Volumes

### **Iron Catalysis – Fundamentals and Applications**

Volume Editor: B. Plietker  
Vol. 33, 2010

### **Medicinal Organometallic Chemistry**

Volume Editors: G. Jaouen, N. Metzler-Nolte  
Vol. 32, 2010

### **C-X Bond Formation**

Volume Editor: A. Vigalok  
Vol. 31, 2010

### **Transition Metal Complexes of Neutral $\eta^1$ -Carbon Ligands**

Volume Editors: R. Chauvin, Y. Canac  
Vol. 30, 2010

### **Photophysics of Organometallics**

Volume Editor: A. J. Lees  
Vol. 29, 2010

### **Molecular Organometallic Materials for Optics**

Volume Editors: H. Le Bozec, V. Guerschais  
Vol. 28, 2010

### **Conducting and Magnetic Organometallic Molecular Materials**

Volume Editors: M. Fourmigué, L. Ouahab  
Vol. 27, 2009

### **Metal Catalysts in Olefin Polymerization**

Volume Editor: Z. Guan  
Vol. 26, 2009

### **Bio-inspired Catalysts**

Volume Editor: T. R. Ward  
Vol. 25, 2009

### **Directed Metallation**

Volume Editor: N. Chatani  
Vol. 24, 2007

### **Regulated Systems for Multiphase Catalysis**

Volume Editors: W. Leitner, M. Hölscher  
Vol. 23, 2008

### **Organometallic Oxidation Catalysis**

Volume Editors: F. Meyer, C. Limberg  
Vol. 22, 2007

### **N-Heterocyclic Carbenes in Transition Metal Catalysis**

Volume Editor: F. Glorius  
Vol. 21, 2006

### **Dendrimer Catalysis**

Volume Editor: L. H. Gade  
Vol. 20, 2006

### **Metal Catalyzed Cascade Reactions**

Volume Editor: T. J. J. Müller  
Vol. 19, 2006

### **Catalytic Carbonylation Reactions**

Volume Editor: M. Beller  
Vol. 18, 2006

### **Bioorganometallic Chemistry**

Volume Editor: G. Simonneaux  
Vol. 17, 2006

### **Surface and Interfacial Organometallic Chemistry and Catalysis**

Volume Editors: C. Copéret, B. Chaudret  
Vol. 16, 2005

### **Chiral Diazaligands for Asymmetric Synthesis**

Volume Editors: M. Lemaire, P. Mangeney  
Vol. 15, 2005

### **Palladium in Organic Synthesis**

Volume Editor: J. Tsuji  
Vol. 14, 2005

### **Metal Carbenes in Organic Synthesis**

Volume Editor: K. H. Dötz  
Vol. 13, 2004

# Iridium Catalysis

Volume Editor: Pher G. Andersson

With Contributions by

John F. Bower · Daniel Carmona · Jongwook Choi ·  
Carmen Claver · Robert H. Crabtree · Montserrat  
Diéguez · Alan S. Goldman · John F. Hartwig ·  
Michael J. Krische · Luis A. Oro · Oscar Pàmies ·  
Andreas Pfaltz · Mark J. Pouy · Ourida Saidi ·  
Jonathan M. J. Williams · David H. Woodmansee

 Springer

*Editor*

Prof. Pher G. Andersson  
Uppsala University  
Department of Biochemistry and Organic Chemistry  
Box 576, BMC  
S-751 23 Uppsala  
Sweden  
pher.andersson@biorg.uu.se

ISSN 1436-6002

e-ISSN 1616-8534

ISBN 978-3-642-15333-4

e-ISBN 978-3-642-15334-1

DOI 10.1007/978-3-642-15334-1

Springer Heidelberg Dordrecht London New York

© Springer-Verlag Berlin Heidelberg 2011

This work is subject to copyright. All rights are reserved, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilm or in any other way, and storage in data banks. Duplication of this publication or parts thereof is permitted only under the provisions of the German Copyright Law of September 9, 1965, in its current version, and permission for use must always be obtained from Springer. Violations are liable to prosecution under the German Copyright Law.

The use of general descriptive names, registered names, trademarks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

*Cover design:* KünkelLopka GmbH

Printed on acid-free paper

Springer is part of Springer Science+Business Media ([www.springer.com](http://www.springer.com))

---

## Volume Editor

Prof. Pher G. Andersson

Uppsala University  
Department of Biochemistry  
and Organic Chemistry  
Box 576, BMC  
S-751 23 Uppsala  
Sweden  
*pher.andersson@biorg.uu.se*

## Editorial Board

Prof. Matthias Beller

Leibniz-Institut für Katalyse e.V.  
an der Universität Rostock  
Albert-Einstein-Str. 29a  
18059 Rostock, Germany  
*matthias.beller@catalysis.de*

Prof. John M. Brown

Chemistry Research Laboratory  
Oxford University  
Mansfield Rd.,  
Oxford OX1 3TA, UK  
*john.brown@chem.ox.ac.uk*

Prof. Pierre H. Dixneuf

Campus de Beaulieu  
Université de Rennes 1  
Av. du Gl Leclerc  
35042 Rennes Cedex, France  
*pierre.dixneuf@univ-rennes1.fr*

Prof. Alois Fürstner

Max-Planck-Institut für Kohlenforschung  
Kaiser-Wilhelm-Platz 1  
45470 Mülheim an der Ruhr, Germany  
*fuerstner@mpi-muelheim.mpg.de*

Prof. Louis S. Hegedus

Department of Chemistry  
Colorado State University  
Fort Collins, Colorado 80523-1872, USA  
*hegedus@lamar.colostate.edu*

Prof. Peter Hofmann

Organisch-Chemisches Institut  
Universität Heidelberg  
Im Neuenheimer Feld 270  
69120 Heidelberg, Germany  
*ph@uni-hd.de*

Prof. Takao Ikariya

Department of Applied Chemistry  
Graduate School of Science and Engineering  
Tokyo Institute of Technology  
2-12-1 Ookayama, Meguro-ku,  
Tokyo 152-8550, Japan  
*tikariya@apc.titech.ac.jp*

Prof. Luis A. Oro

Instituto Universitario de Catálisis Homogénea  
Department of Inorganic Chemistry  
I.C.M.A. - Faculty of Science  
University of Zaragoza-CSIC  
Zaragoza-50009, Spain  
*oro@unizar.es*

Prof. Manfred Reetz

Max-Planck-Institut für Kohlenforschung  
Kaiser-Wilhelm-Platz 1  
45470 Mülheim an der Ruhr, Germany  
*reetz@mpi-muelheim.mpg.de*

Prof. Qi-Lin Zhou

State Key Laboratory of Elemento-organic  
Chemistry  
Nankai University  
Weijin Rd. 94, Tianjin 300071 PR China  
*qlzhou@nankai.edu.cn*



# Topics in Organometallic Chemistry

## Also Available Electronically

*Topics in Organometallic Chemistry* is included in Springer's eBook package *Chemistry and Materials Science*. If a library does not opt for the whole package the book series may be bought on a subscription basis. Also, all back volumes are available electronically.

For all customers who have a standing order to the print version of *Topics in Organometallic Chemistry*, we offer free access to the electronic volumes of the Series published in the current year via SpringerLink.

If you do not have access, you can still view the table of contents of each volume and the abstract of each article by going to the SpringerLink homepage, clicking on "Chemistry and Materials Science," under Subject Collection, then "Book Series," under Content Type and finally by selecting *Topics in Organometallic Chemistry*.

You will find information about the

- Editorial Board
- Aims and Scope
- Instructions for Authors
- Sample Contribution

at [springer.com](http://springer.com) using the search function by typing in *Topics in Organometallic Chemistry*.

Color figures are published in full color in the electronic version on SpringerLink.

## Aims and Scope

The series *Topics in Organometallic Chemistry* presents critical overviews of research results in organometallic chemistry. As our understanding of organometallic structures, properties and mechanisms grows, new paths are opened for the design of organometallic compounds and reactions tailored to the needs of such diverse areas as organic synthesis, medical research, biology and materials science. Thus the scope of coverage includes a broad range of topics of pure and applied organometallic chemistry, where new breakthroughs are being made that are of significance to a larger scientific audience.

The individual volumes of *Topics in Organometallic Chemistry* are thematic. Review articles are generally invited by the volume editors.

In references *Topics in Organometallic Chemistry* is abbreviated Top Organomet Chem and is cited as a journal.





# Preface

The use of iridium in catalysts for organic reactions is a rapidly evolving field that covers a wide range of chemical reactivity. With this in mind, it has been a pleasure to edit this volume on Iridium Catalysis, which aims to present the reader with a comprehensive resource that covers the field and focuses on recent advances in the area. All chapters have been written by leading chemists in the corresponding area and I am extremely thankful to them for their contributions and efficient collaboration in this project: Robert H Crabtree, Carmen Claver, Montserrat Dieguez, Oscar Pamies, Andreas Pfaltz, David H. Woodmansee, Jonathan M.J. Williams, Ourida Saidi, Michael J. Krische, John F. Bower, Alan S. Goldman, Jongwook Choi, John F. Hartwig, Mark J. Pouy, Luis A. Oro, and Daniel Carmona. I am also grateful for all help received from SPRINGER, especially from Jutta Lindenborn.

I hope that this book will be useful to researchers, teachers, and students involved in organic synthesis and that it might also attract new people to the interesting field of catalysis.

Uppsala, Sweden  
2010

Pher G. Andersson



# Contents

<b>Introduction and History</b> .....	1
Robert H. Crabtree	
<b>Iridium-Catalyzed Hydrogenation Using Phosphorus Ligands</b> .....	11
Montserrat Diéguez, Oscar Pàmies, and Carmen Claver	
<b>Iridium-Catalyzed Asymmetric Hydrogenation of Olefins with Chiral N,P and C,N Ligands</b> .....	31
David H. Woodmansee and Andreas Pfaltz	
<b>Iridium-Catalyzed Hydrogen Transfer Reactions</b> .....	77
Ourida Saidi and Jonathan M. J. Williams	
<b>Formation of C–C Bonds via Iridium-Catalyzed Hydrogenation and Transfer Hydrogenation</b> .....	107
John F. Bower and Michael J. Krische	
<b>Ir-Catalyzed Functionalization of C–H Bonds</b> .....	139
Jongwook Choi and Alan S. Goldman	
<b>Iridium-Catalyzed Allylic Substitution</b> .....	169
John F. Hartwig and Mark J. Pouy	
<b>Iridium-Catalyzed 1,3-Dipolar Cycloadditions</b> .....	209
Daniel Carmona and Luis A. Oro	
<b>Index</b> .....	231



# Introduction and History

Robert H. Crabtree

**Abstract** Iridium had shown no special advantage in heterogeneous catalysis and this situation seemed to carry over into the early work in homogeneous catalysis. In particular, the failure of  $\text{IrCl}(\text{PPh}_3)_3$ , the Ir analog of Wilkinson's catalyst, to give catalytic hydrogenation strongly influenced thinking at the time. The developments that changed the situation were Shrock and Osborn's work on  $[(\text{cod})\text{RhL}_2]^+$  complexes that showed how lowering the L:M ratio from 3 to 2 gave improved catalysts, followed by our own finding that  $[(\text{cod})\text{IrL}_2]^+$  and especially  $[(\text{cod})\text{IrLL}']^+$  complexes are even more highly active, particularly for highly substituted alkenes. This performance is absent in the classical coordinating solvents then standard in the field but only appears in weakly coordinating solvents like dichloromethane. A special property of the Ir complexes is a pronounced directing effect caused by catalyst binding to substrate functionality, leading to very high diastereoselectivity of the reduction. Asymmetric hydrogenations consistently gave lower *ee* values for iridium over rhodium, however, so academic work still strongly emphasized Rh. The exceedingly rapid rates of hydrogenation with Ir attracted industrial attention, because rate is such an important factor in that case, and Novartis' catalytic asymmetric route to the pesticide (S)-Metolachlor employs iridium, even though the *ee* value is only ca 80%. Leading applications of these and other Ir catalysts up to about 1,990 are also discussed. Work by Pfaltz, Burgess, and Andersson showed how efficient asymmetric Ir catalysts could be made by judicious choice of ligand. Alkane dehydrogenation by reverse hydrogenation also proved possible with Ir catalysts.

**Keywords** CH activation · Hydrogenation · Isotope exchange

## Contents

1	Introduction .....	2
2	Vaska's Complex .....	2
3	Schrock and Osborn Catalysts .....	3
4	Distinctive Reactivity for Ir in Hydrogenation Catalysis .....	4
5	Further Developments in Hydrogenation .....	6
6	Mechanism .....	7
7	Alkane Activation .....	7
8	Conclusion .....	8
	References .....	9

## 1 Introduction

The catalysis field began in the nineteenth century with work on heterogeneous platinum group metals from such contributors as Berzelius, Faraday, Döbereiner, Mitscherlich, Davy, Sabatier, and Ostwald [1, 2]. Iridium never held a very high place in the field, however, because it showed no special advantages. Rylander expresses this point well in his classic 1967 work on heterogeneous catalysis [3].

Iridium... makes a fair catalyst and its lack of use stems partly from neglect and partly from the fact that some platinum metal has usually proved more suitable whenever a comparison was made.

The position for heterogeneous Ir is little changed today, as shown by the entries in the Encyclopedia of Reagents for Organic Synthesis [4]. Heterogeneous Pd, Pt, Ru, and Rh catalysts are all discussed in detail but Ir and Os are omitted.

This background influenced the first work in homogeneous catalysis, where Pd, Pt, Ru, and Rh all gained early favor over Ir and Os. Neglect again played a role – investigators tend to follow up lines previously shown to be successful – but an important factor was the failure of  $\text{IrCl}(\text{PPh}_3)_3$ , the Ir analog of Wilkinson's catalyst, to give catalytic hydrogenation.

## 2 Vaska's Complex

The work of Lauri Vaska [5] at Clarkson College in the 1960s deserves special attention. His investigations on  $\text{IrCl}(\text{CO})(\text{PPh}_3)_2$ , now known as Vaska's complex, were the first to give iridium a major and distinctive role for organometallic chemists in contributing to the understanding of oxidative addition, a fundamental reaction step in homogeneous catalysis. This work was particularly impressive since it was carried out at a small college with undergraduate coworkers. He showed that a variety of molecules A–B were able to react with  $\text{IrCl}(\text{CO})(\text{PPh}_3)_2$  to give  $\text{Ir}(\text{A})(\text{B})\text{Cl}(\text{CO})(\text{PPh}_3)_2$ , the oxidative addition product. By monitoring the CO stretching frequencies of the adducts, he was able to establish the oxidative

character of the reaction and rank different A–B molecules in terms of their relative oxidizing tendency. The stability of the adducts did not bode well for their intermediacy in catalysis, however, and indeed Vaska's complex never showed any more than a modest degree of catalytic activity. Iridium was therefore considered to be interesting only in giving isolable analogs of species thought to be important in catalysis by rhodium complexes.

Just as the H<sub>2</sub> oxidative addition product of Vaska's complex is relatively stable, the same holds for the iridium analog of Wilkinson's catalyst. The reason for the lack of high catalytic activity of IrCl(PPh<sub>3</sub>)<sub>3</sub> is that once the required oxidative addition of H<sub>2</sub> has taken place, the product, IrH<sub>2</sub>Cl(PPh<sub>3</sub>)<sub>3</sub>, is stable and does not dissociate a phosphine to provide a site for binding of substrate alkene. The essence of catalysis is the involvement of unstable catalytic intermediates. For example, RhH<sub>2</sub>Cl(PPh<sub>3</sub>)<sub>3</sub>, the analogous intermediate from Wilkinson's catalyst, is unstable in that it does easily lose a PPh<sub>3</sub> ligand – the one trans to the high trans effect hydride – and thus liberates a coordination site at which the substrate alkene can bind. The resulting alkene dihydride complex, RhH<sub>2</sub>Cl(alkene)(PPh<sub>3</sub>)<sub>2</sub>, now has both substrates bound, both alkene and hydrogen, and can thus proceed directly to product by alkene insertion into the Rh–H bond to give an alkyl hydride that can reductively eliminate the alkene product. The reason for the difference between Rh and Ir is probably the systematic increase in M–L bond strengths on going from the second row of the transition metals to the third. In line with this idea, moving to a Wilkinson type complex of a smaller and more tightly bound phosphine, PEt<sub>3</sub>, is enough to halt the catalysis even for Rh because RhH<sub>2</sub>Cl(PEt<sub>3</sub>)<sub>3</sub> is a stable species.

### 3 Schrock and Osborn Catalysts

The situation changed when Schrock and Osborn [6, 7] at Harvard developed catalysts in which the ligand to metal ratio was 2:1 and not 3:1, thus avoiding the need for phosphine dissociation. At the same time, they also removed the chloride ligand from the inner sphere and replaced it by an outer sphere “noncoordinating” anion, often BF<sub>4</sub>. Building on Chatt's work [8] showing that 1,5-cyclooctadiene (cod) is a reliable ligand for Rh(I), they synthesized [(cod)Rh(PPh<sub>3</sub>)<sub>2</sub>]BF<sub>4</sub>. Under hydrogen, the cod is removed by hydrogenation to cyclooctane and the solvated [(sol)<sub>2</sub>Rh(PPh<sub>3</sub>)<sub>2</sub>]<sup>+</sup> cation is formed. This is an active hydrogenation catalyst, but the key point is that the system is no longer limited to PPh<sub>3</sub> as in the Wilkinson case. Instead, a wide variety of phosphines, L, or chelating phosphines, L–L, can give catalytically active species [(cod)Rh(L)<sub>2</sub>]BF<sub>4</sub> or [(cod)Rh(L–L)]BF<sub>4</sub>. This led to the possibility of using asymmetric ligands like Kagan's chelating *diop* that allow asymmetric hydrogenation of prochiral alkenes to give homochiral products with high ee.

While the Schrock and Osborn catalysts worked well in the Rh series, the analogous iridium compounds, also studied by them, did not show any special



advantages. The reason proved to be that instead of forming analogs of the  $[(\text{solv})_2\text{Rh}(\text{PPh}_3)_2]^+$  cation, the iridium species  $[(\text{cod})\text{Ir}(\text{L})_2]\text{BF}_4$  reacted differently with hydrogen. The product was the relatively stable  $[(\text{solv})_2\text{IrH}_2(\text{PPh}_3)_2]^+$  cation. This gave a modest catalytic rate but, as in the case of heterogeneous Ir catalysts, no special properties were apparent that would make them the focus of an investigation. The reason the Ir compounds behave differently is again the rise in M–L bond strengths on going from Rh to Ir. This allows the  $\text{H}_2$  oxidative addition product to form for Ir but not for Rh. The two M–H bonds formed in the oxidative addition have to “pay for” breaking the H–H bond of  $\text{H}_2$  if the oxidative addition is to be thermodynamically allowed.

## 4 Distinctive Reactivity for Ir in Hydrogenation Catalysis

The next step came in the laboratory of Hugh Felkin, a very distinguished physical organic chemist, working at the CNRS Natural Products Institute at Gif-sur-Yvette near Paris. Felkin is perhaps best known for his formulation of the Felkin–Anh rule for predicting the diastereoselectivity of nucleophilic attack in organic carbonyl compounds [9]. He had become interested in the organic applications of transition metals and had hired a series of postdoctoral coworkers from the UK to look into the area. On this basis, I was working as an *Attaché de Recherche* with a 3<sup>ème</sup> cycle doctoral thesis student from Oxford, George Morris. The plan was to look at the  $[(\text{cod})\text{Ir}(\text{L})_2]\text{BF}_4$  series with the idea of isolating stable intermediates analogous to the ones that were presumed to be important in the Schrock–Osborn hydrogenation catalysts.

One of the fixed assumptions in homogeneous hydrogenation catalysis at that time was that reducing, coordinating solvents like MeOH or EtOH were preferred and oxidizing solvents like dichloromethane were best avoided. This came from the work on the Wilkinson catalyst, showing that the alcohols acted well but the catalyst deactivated in chlorinated solvents such as dichloromethane. The Schrock–Osborn hydrogenation work confirmed the role of coordinating solvents with the finding that the catalyst precursor forms solvated  $[(\text{solv})_2\text{Rh}(\text{PPh}_3)_2]^+$  cations under hydrogen. For Rh, catalytic activity was maintained, however, because the solvent ligands being labile, alkene substrates are able to gain access to the metal, a step that is followed by oxidative addition of  $\text{H}_2$ . The resulting alkene dihydride intermediate undergoes insertion and reductive elimination just as in the Wilkinson cycle.

Working from the principle that a catalyst should be maximally unstable without actually decomposing, I had tried “noncoordinating” solvents like dichloromethane for the  $[(\text{cod})\text{Rh}(\text{L})_2]\text{BF}_4$  case and seen that a high catalytic rate was achievable in certain cases. When George Morris began work on the Ir series, he included noncoordinating solvents and found high activity for dichloromethane. The solvent effect in this case was extremely high: dichloromethane not only gave much faster catalysts than the traditional solvents but now even highly substituted alkenes – poor

ligands and therefore poor substrates for classical catalysts – were reactive. None of these effects were seen with the Rh series, so iridium was special. A factor that may well explain the inertness of dichloromethane in these cases is the cationic charge on the metal, making them more resistant to deactivation by oxidizing solvents.

During a discussion with G Pannetier (Université Paris VI) in Paris, he mentioned that he thought he had seen evidence for mixed pyridine–phosphine derivatives of the type  $[(\text{cod})\text{Rh}(\text{PPh}_3)(\text{C}_5\text{H}_5\text{N})]\text{BF}_4$  in the Rh series. On returning to Gif, I confirmed this result by isolating the compound and mentioned it to Morris, who tried to obtain the iridium complex. Our initial idea was that a 1:1 phosphine to metal catalyst might be even more active than the 2:1 species.

The  $\text{PPh}_3$  derivative showed very promising activity, but Morris tried a few other phosphines as well and identified tricyclohexylphosphine as the best of these. In fact, the new  $\text{PCy}_3$  catalyst was spectacularly active, particularly for highly substituted alkenes for which classical catalysts gave zero rate. Indeed, the lower the catalyst concentration, the higher the apparent rate, presumably due to mass transfer limitations for dissolution of the hydrogen. Our first attempt to publish these results in *J. Am. Chem. Soc.* failed because the referees were not impressed. We therefore published then in *J. Organometal Chem.*; but at the time, the results attracted little attention [10]. This was a period when organic chemists were already familiar with organolithium reagents, but transition metals had not yet been fully integrated into their thinking. In addition, we had used substrates like hexene, cyclohexene, 2-methyl-2-butene, and 2,3-dimethyl-2-butene that were regarded as unrepresentative of complex organic molecules.

By 1980, I was already at Yale and met Bill Suggs at Bell Labs, where he was working on catalysis from the organic perspective. With characteristic generosity, he suggested we collaborate on trying to use  $[(\text{cod})\text{Ir}(\text{PCy}_3)(\text{C}_5\text{H}_5\text{N})]\text{BF}_4$  for hydrogenation of substrates that were more complex than simple hydrocarbons. This proved successful, was published in *Tet Lett*, and did begin to interest organic chemists [11]. This experience made me think that it would be good to try the catalyst for more complex substrates.

I consulted Sam Danishevsky's wife, Sarah, who was then working in the Yale Department on synthetic problems. She suggested that a good problem would be to look for directing effects of preexisting functionality on the diastereoselectivity of hydrogen addition. My graduate student, Mark Davis, took up the problem and found evidence for very high selectivity in unsaturated cyclic alcohols. An OH group on one face of the substrate molecule was able to bind to the metal in such a way that the two hydrogen atoms could only add to that same face of the ring that contained the OH group. These papers [12, 13], published in *Organometallics* and *J. Org. Chem.*, together with parallel work from Stork and Kahne [14] at Columbia, were the trigger that attracted the interest of a much broader organic community. No doubt Stork's eminence in the field played a major role in acceptance of the work. From that point onward, development of the iridium catalyst and its many analogs passed into the hands of organic chemists. In the first such example, Arthur Schultz at RPI published a synthesis of (+)-Pumiliotoxin C in 1985–1987 that

employed  $[(\text{cod})\text{Ir}(\text{PCy}_3)(\text{C}_5\text{H}_5\text{N})]\text{BF}_4$  for carbonyl-directed hydrogenation of a trisubstituted  $\text{C}=\text{C}$  bond [15].

## 5 Further Developments in Hydrogenation

The Ir catalysts are also active for a number of other reactions. In a purely mechanistic study from 1985, we looked at H/D exchange between deuterium and a number of organic substrates [16]. We found that initial binding of the catalyst to a ligating group such as a quinoline nitrogen allowed the catalyst to cyclometalate at a specific adjacent site, followed by rapid deuteration at that site. This was much later adapted for selective tritiation by pharmaceutical chemists, who needed to be able to follow the metabolic fate of drugs by radioisotope labeling. Traditional syntheses had incorporated tritium at an early stage, resulting in tedious syntheses and production of radioactive waste on a significant scale. In the catalytic variant, the tritium is introduced in one step, making the iridium catalyst a standard in the field. In mechanistic work with Odile Eisenstein, the noted computational organometallic chemist from Montpellier (France), the reason for the rapid isotope exchange was traced to the presence of an unusually flat potential energy surface that in turn results from compensatory changes in M–L bond energies through the cycle [17].

Asymmetric hydrogenation with analogous catalysts proved possible. Pfaltz introduced cationic iridium complexes with chiral mixed P,N-donor ligands as catalysts for asymmetric hydrogenation. A phosphino-oxazoline (PHOX) was found to be best when combined with the large, weakly coordinating anion tetrakis[3,5-bis(trifluoromethyl)-phenyl]borate (BARF); the anion helps avoid deactivation and reduces moisture sensitivity. Hydrogenation of trisubstituted stilbenes were carried out with high levels of enantioselectivity. Under optimized conditions, turnover numbers were  $>5,000$  [18–20]. Burgess and coworkers have had success with asymmetric carbene oxazoline ligands [21]. An unusual sulfinylimine phosphine ligand, applied to challenging trisubstituted stilbene substrates by Ellman, gives ee's of up to 94% [22]. Andersson has shown advantages from chiral bidentate phosphine thiazoles in asymmetric hydrogenation of aryl alkenes and aryl alkene esters, substrates that can be difficult to hydrogenate with classical catalysts [23].

In a commercially important example, a related iridium catalyst equipped with a ferrocenyl bis phosphine ligand of Togni's proved active for the key asymmetric hydrogenation in the production of an important agrochemical, the herbicide (S)-metolachlor, having the trade name Dual Magnum<sup>®</sup>. The key step in the process is the asymmetric hydrogenation of an imine intermediate with an iridium ferrocenyl diphosphine catalyst. The isolated imine can be hydrogenated at a hydrogen pressure of 80 bar and 50°C with a substrate-to-catalyst ratio of  $>10^6$ . Complete conversion is reached within 4 h with an enantioselectivity of 79% with an initial turnover frequency (TOF) exceeding  $1.8 \times 10^6 \text{ h}^{-1}$  [24].

## 6 Mechanism

Mechanistic aspects of the iridium catalysts have attracted attention. For catalytic reactions where Ir does show special properties, we need to ask what mechanistic factors are at work that differentiate the two metals. An important difference between Rh and Ir is the much easier accessibility of the M(V) state for Ir. For example,  $\text{IrH}_5(\text{PPh}_3)_2$  is a stable species while no Rh analog has ever been reported. The mechanism invoked for Rh(I) hydrogenation catalysts involves cycling between Rh(I) and Rh(III) oxidation states. An attractive possibility for the Ir catalyst is an Ir(III)–Ir(V) cycle, although this is still subject to debate. Hall, Burgess, and their coworkers report computational data that favors this III–V cycle for diene hydrogenation by a closely related Ir catalyst, for example [25].

In all the applications described above, the iridium catalyst showed special properties not so far obtainable with any other metal. We now move on to reactions in which iridium is active but where other metals have also been shown to be suitable.

## 7 Alkane Activation

A point of particular interest to me even at Gif but particularly on my move to Yale in 1977 was whether such an active hydrogenation catalyst as we had in hand could be induced to reverse direction to provide catalytic alkane dehydrogenation to alkene or arene. This would constitute a CH activation. Granted, the reaction would be endoergic, but I felt it might be possible to capture or remove the hydrogen in some way to drive the reaction. The importance of CH activation was impressed on me right from my doctoral work with Joseph Chatt, who held that CH activation would be the next big field in transition metal chemistry after nitrogen fixation.

This approach seemed feasible because Alexander Shilov, working at the Chernogolovka Institute of Chemical Physics, had already shown catalytic deuteration and chlorination of alkanes by Pt complexes. My discussions with colleagues in the late 1970s suggested that this result was discredited by the majority of workers in the West – as I felt, unjustly so. It was thought that colloidal Pt metal, a well-known heterogeneous catalyst for alkane conversions, might be the true active species. Personally, I felt that the selectivity for reaction at the terminal methyl of the alkane substrates seen by Shilov was much more consistent with homogeneous than heterogeneous catalysis. Pt-catalyzed reactions of alkanes are now fully confirmed and widely studied and the field is called “Shilov chemistry” [26].

My initial plan involved dehydrogenation of cyclic alkanes to give stable arenes, particularly of cyclopentane to give cyclopentadienyl derivatives. My own experimental efforts with Pt were unavailing, but Jennifer Quirk, a graduate student of mine working on Ir, took an interest in the project and found a small yield of Cp

complex from  $[\text{IrH}_2(\text{acetone})_2(\text{PPh}_3)_2]\text{BF}_4$  in dichloromethane. This was too small a yield to publish and so she looked for hydrogen acceptors that could drive the reaction. After trying more than 70 alkenes, she obtained success with  $\text{tBuCH}=\text{CH}_2$ . The yield was now high enough to publish. This time, *J. Am. Chem. Soc.* accepted the manuscript, which was published in 1979 [27]. George Whitesides was clearly a referee on the manuscript because he contacted me before proceeding with his report to warn me, in an entirely kindly way, that such results were often misconceived and later might have to be retracted with consequent damage to my career. I admitted that there was a risk but wanted to go ahead anyway. At a Gordon Conference several years later, he mentioned in his talk that he had been doubtful about our results, but with the accumulation of later results, most notably from Bob Bergman at UC Berkeley [28], he was happy to state that he now believed the work was correct. Bergman, being a highly respected figure in the field, did the most to convince colleagues of the reality of homogeneous CH activation. His work, being stoichiometric, does not directly enter into this discussion of catalysis, but it was undoubtedly very influential. The groups of Bill Jones at Rochester and Bill Graham at Edmonton also published very similar findings during the same period [29, 30]. These key contributions demonstrated stoichiometric oxidative addition of alkane CH bonds to highly reactive Ir or Rh intermediates generated by photolysis of a stable precursor.

Hugh Felkin was also working in the area and, by 1982, developed the first truly catalytic systems for alkane conversion to alkenes, using  $\text{tBuCH}=\text{CH}_2$  to drive the process [31–33]. A variety of catalysts proved useful including  $[\text{IrH}_5(\text{PPh}_3)_2]$ . An important point that emerged was the strong kinetic selectivity for the formation of the 1-alkene from n-alkane, implying that the same selectivity for initial attack at the terminal methyl group applied here as Shilov had seen earlier. This outcome is entirely opposed to the selectivity seen for electrophilic and radical attack, which occurs preferentially at the 2° positions or 3° for branched alkanes. Once again, this provided evidence for the absence of any classical organic reaction pathways in this chemistry. Our own later catalyst  $[\text{Ir}(\text{O}_2\text{CCF}_3)\text{H}_2(\text{PPh}_3)_2]$  shows similar behavior and dates from 1987 [34].

Alan Goldman developed much improved catalysts and also obtained detailed mechanistic information in the 1990s [35, 36]. An unexpected variant developed by Saito, by our own group and most extensively by Goldman, was acceptorless dehydrogenation. It turned out that the reaction could be driven by reflux because the hydrogen was continuously pumped out of the solvent by the reflux action [37–39].

## 8 Conclusion

Iridium is now a well-accepted choice for homogeneous catalysis, thanks to the historical developments discussed. Shrock and Osborn's work on  $[(\text{cod})\text{RhL}_2]^+$  complexes that showed how lowering the L:M ratio from 3 to 2 gave excellent

catalysts. We found that [(cod)IrL<sub>2</sub>]<sup>+</sup> and especially [(cod)IrLL']<sup>+</sup> complexes become very highly active, even for highly substituted alkenes, but only in weakly coordinating solvents such as dichloromethane. The Ir complexes show a pronounced directing effect of existing functionality on the diastereoselectivity of the reduction. The exceedingly rapid rates of hydrogenation with Ir attracted industrial attention, and Novartis' catalytic asymmetric route to the pesticide (S)-Metolachlor employs iridium, even though the ee value is only ca 80%. Work by Pfaltz, Burgess, and Andersson showed how efficient asymmetric Ir catalysts could be made by a judicious choice of ligand. Alkane activation by reverse hydrogenation also proved possible with Ir catalysts.

## References

1. Lindström B, Pettersson LJ (2004) *Cattech* 130:7
2. Nye MJ (1977) *Isis* 68:375
3. Rylander PN (1967) *Catalytic hydrogenation over platinum metals*. Academic, New York
4. Paquette LA (ed) (2006) *Encyclopedia of reagents for organic synthesis*. Wiley, New York
5. Vaska L (1968) *Accounts Chem Res* 1:335
6. Schrock RR, Osborn JA (1971) *J Am Chem Soc* 93:3089
7. Schrock RR, Osborn JA (1976) *J Am Chem Soc* 98:2134, 2143, 4450
8. Chatt J, Venanzi LM (1957) *J Chem Soc* 4735
9. Cherest M, Felkin H, Prudent N (1968) *Tetrahedron Lett* 18:2199
10. Crabtree RH, Felkin H, Morris GE (1977) *J Organomet Chem* 141:205
11. Suggs JW, Cox SD, Crabtree RH, Quirk JM (1981) *Tetrahedron Lett* 22:303
12. Crabtree RH, Davis MW (1983) *Organometallics* 2:681
13. Crabtree RH, Davis MW (1986) *J Org Chem* 51:2655
14. Stork G, Kahne DE (1983) *J Am Chem Soc* 105:1072
15. Schultz AG, McCloskey PJ, Court JJ (1987) *J Am Chem Soc* 109:6493
16. Crabtree RH, Holt EM, Lavin M, Morehouse SM (1986) *Inorg Chem* 1985:24
17. Clot E, Chen JY, Lee DH, Sung SY, Appelhans LN, Faller JW, Crabtree RH, Eisenstein O (2004) *J Am Chem Soc* 126:8795
18. Roseblade SJ, Pfaltz ACR (2007) *Chimie* 10:178
19. Smidt SP, Zimmermann N, Studer M, Pfaltz A (2004) *Chem Eur J* 10:4685
20. Helmchen G, Pfaltz A (2000) *Acc Chem Res* 33:336
21. Hou D-R, Reibenspies JH, Colacot TJ, Burgess K (2001) *Chem Eur J* 7:5391
22. Schenkel LB, Ellman JA (2004) *J Org Chem* 69:1800
23. Källström K, Hedberg C, Brandt P, Hansen LK, Andersson PG (2004) *J Am Chem Soc* 126:14308
24. Blaser HU, Buser HP, Coers K, Hanreich R, Jalett HP, Jelsch E, Pugin B, Schneider HD, Spindler F, Wegmann A (1999) *Chimie* 53:275
25. Cui X, Fan Y, Hall MB, Burgess K (2005) *Chem Eur J* 11:6859
26. Shilov AE, Shul'pin GB (2000) *Activation and catalytic reactions of saturated hydrocarbons*. Kluwer, Dordrecht
27. Crabtree RH, Mihelcic JM, Quirk JM (1979) *Iridium complexes in alkane dehydrogenation*. *J Am Chem Soc* 101:7738
28. Janowicz AH, Periana RA, Buchanan JM, Kovac CA, Stryker JM, Wax MJ, Bergman RG (1984) *Pure Appl Chem* 56:13
29. Jones WD, Feher FJ (1984) *J Am Chem Soc* 106:1650
30. Hoyano JK, Graham WAG (1982) *J Am Chem Soc* 104:3723

31. Baudry D, Ephritikhine M, Felkin H (1980) Chem Commun 1243
32. Baudry D, Ephritikhine M, Felkin H, Zakrzewski J, (1982) Chem Commun 1235
33. Felkin H, Fillebeen-Khan T, Holmes-Smith R, Yingrui L (1985) Tetrahedron Lett 26:1999
34. Burk MJ, Crabtree RH (1987) J Am Chem Soc 109:8025
35. Maguire JA, Boese WT, Goldman AS (1990) Abstracts Paper ACS 199: 459-INOR
36. Liu FC, Pak EB, Singh B, Jensen CM, Goldman AS (1999) J Am Chem Soc 121:4086
37. Yukawa K, Fujii T, Saito Y (1991) Chem Commun 1548
38. Aoki T, Crabtree RH (1993) Organometallics 12:294
39. Xu WW, Rosini GP, Gupta M, Jensen CM, Kaska WC, Krogh-Jespersen K, Goldman AS (1997) Chem Commun 2273

# Iridium-Catalyzed Hydrogenation Using Phosphorus Ligands

Montserrat Diéguez, Oscar Pàmies, and Carmen Claver

**Abstract** Due to the requirement of obtaining enantiomerically pure products through environmentally friendly technologies, the development of new methods in asymmetric catalysis is becoming an important approach to asymmetric synthesis. Asymmetric hydrogenation is one of the most applied catalytic reactions for the preparation of enantiomerically pure products. Iridium complexes containing phosphorus ligands have been less applied as catalysts in hydrogenation reaction than their corresponding rhodium complexes. On the other hand, most of the iridium complexes used as catalyst in hydrogenation reaction contain phosphorus–nitrogen ligands. However, iridium complexes containing a great variety of mono- and bidentate phosphorus ligands (phosphines, phosphinites, phosphites, phosphoramidites, and the combination of these functionalities) have been applied in hydrogenation reaction with different grades of success. In the early days, diphosphines were the most widely used phosphorus ligands. However, they are generally difficult to synthesize and prone to oxidation. In recent years, chiral phosphites, phosphinites, phosphonites, and phosphoroamidites have emerged as new types of ligands, which present several advantages and high catalytic activity and selectivity. Homogeneous and heterogeneous systems have been studied, allowing the recycling of catalysts. Slight modifications in the electronic and steric properties of the ligands influence the catalytic results, which indicates the potentiality of modular ligands. Imines, including unfunctionalized N–H imines and cyclic imines as quinolines and other substrates, have been transformed in the corresponding chiral amines with ee higher than 95% in many cases. Binol-derived phosphoroamidite constitute one of the most successful ligands in these asymmetric reaction achieving, in some cases, practically total enantioselectivity.

**Keywords** Asymmetric catalysis · Asymmetric hydrogenation · Imine · Iridium catalysts · Phosphorus ligands

---

M. Diéguez, O. Pàmies, and C. Claver (✉)

Departament de Química Física i Inorgànica, Universitat Rovira i Virgili, C/Marcel·lí Domingo s/n, Tarragona 43007, Spain

e-mail: montserrat.dieguez@urv.cat; oscar.pamies@urv.cat; carmen.claver@urv.cat



## Contents

1	Introduction .....	12
2	Application of Bidentate Phosphorus Ligands .....	13
2.1	Bidentate Diphosphines .....	13
2.2	Bidentate Diphosphinite, Phosphite–Phosphinite, Diphosphite and Phosphine–Phosphoroamidite Ligands .....	19
3	Application of Monodentate Phosphorus Ligands .....	23
3.1	Monodentate Phosphite, Phosphoroamidite, and Phosphorothioite Ligands .....	23
3.2	Monodentate Phosphines .....	27
4	Conclusions .....	28
	References .....	28

## 1 Introduction

Hydrogenation catalyzed by transition metals is one of the most efficient methods for reduction of C=C, C=O, and C=N double bonds [1, 2]. Among the different transition metals used, iridium plays an important role in the organic transformations through hydrogenation [3]. To achieve the highest levels of reactivity and selectivity in catalytic hydrogenation, a careful selection and design of the ligand is perhaps the most crucial step. Particular effort has been developed in the last years in the synthesis of phosphorus ligands due to the effectiveness of these ligands in homogenous catalysis. Their diversity (phosphines, phospholes, phosphetanes, phosphinites, phosphonites, phosphites, phosphoramidites, etc.) together with the possibilities of variation of electronic and steric properties make these ligands extremely attractive for catalysis, providing, in many cases, successful results. Chelate phosphorus chiral ligands are considered to be the best ligands for transition metals. In the early days, diphosphines were the most widely used phosphorus ligands. However, they are generally difficult to synthesize and prone to oxidation. In recent years, chiral diphosphites, diphosphinites, diphosphonites, and phosphoroamidites have emerged as new types of ligands, which present several advantages and high catalytic activity and selectivity in metal catalyzed reactions. These ligands, however, are more prone to hydrolysis than phosphines. Their modular nature enables their steric and electronic properties to be finely tuned. For example, binaphthol and its derivatives are a class of chiral diols that are useful for synthesizing these ligands and varying their properties. Interestingly, each catalytic process requires a particular type of ligand. There is no universal type of ligand that can be considered to be more efficient than the others because efficiency also depends on the nature of the substrate [4].

Due to the requirement of obtaining enantiomerically pure products through environmentally friendly technologies, the development of new methods in asymmetric catalysis is becoming an important approach to asymmetric synthesis. Asymmetric hydrogenation is one of the most applied catalytic reactions for the

preparation of enantiomerically pure products. Many chiral phosphorus ligands have been developed to be applied in this important reaction [5–7].

Iridium complexes containing phosphorus ligands have been less applied as catalysts in hydrogenation reaction than the corresponding rhodium complexes containing phosphorus ligands. On the other hand, most of the iridium complexes used as catalyst in hydrogenation reaction contain phosphorus–nitrogen ligands. However, iridium complexes containing a great variety of phosphorus ligands (phosphines, phosphinites, phosphites, phosphoramidites) have been applied in hydrogenation reaction with different grades of success. Due to the effectiveness of the iridium catalysts in C=N hydrogenation, the iridium complexes are mainly used as catalysts in imines and related substrates. Although the asymmetric hydrogenation of imines presents important drawbacks (the coordination of substrate, which can take place through both the nitrogen donor atom and the double bond, the *E/Z* isomeric mixture of imine substrates, and the poisoning effect of the resultant amines on the catalysts), recent progress in the field of asymmetric hydrogenation of imines allow high activities and enantioselectivities [8].







In general, two different kinds of iridium precursors are used. Isolated cationic  $[\text{Ir}(\text{COD})_2]\text{A}$  complexes and the “in situ” precursors formed from the neutral dimer  $[\text{Ir}(\mu\text{-Cl})(\text{COD})]_2$  by addition of the corresponding phosphorus ligand.

This chapter collects the most relevant cases of iridium complexes applied in hydrogenation reaction classified according to the nature of the phosphorus ligand.

## 2 Application of Bidentate Phosphorus Ligands

### 2.1 Bidentate Diphosphines

Iridium complexes containing diphosphines have been mainly used as catalysts in the asymmetric hydrogenation of imines for the preparation of chiral amines. Optically active amines are useful intermediates in the synthesis of biologically active compounds. Therefore, the development of efficient methods for their preparation is synthetically important. The first reports concern the use of iridium with the well-known diphosphines DIOP [9, 10], BINAP [10], and BDPP [10, 11] (Fig. 1) as ligands in the hydrogenation of acyclic and cyclic imines. Excellent enantioselectivities (ee's up to 90% at  $T = 0^\circ\text{C}$ ,  $P_{\text{H}_2} = 40$  bar) were obtained in the hydrogenation of *N*-[(*Z*)-1-methoxypropan-2-ylidene]-2,6-dimethylaniline, which is used as a model herbicide precursor [12, 13], with the  $[\text{Ir}(\text{BDPP})(\text{OCOCF}_3)_3]$  catalyst system. The use of a mixture of THF/ $\text{CH}_2\text{Cl}_2$  (3/1) was found to be crucial, although in some cases, the use of pure THF provided faster hydrogenation rates. The authors also observed that increasing the hydrogen pressure increases the catalytic activity without changing the enantiomeric excess. For substrates 2,3,3-trimethyl-3H-indole and *N*-[(*E*)-1-phenylethylidene]aniline, moderate enantioselectivities were obtained. Interestingly, for the latter substrate Ir-BINAP catalytic


	 <b>DIOP</b>	 <b>BINAP</b>	 <b>BDPP</b>
	29 ( <i>S</i> )	67 ( <i>S</i> )	90 ( <i>R</i> )
	35 (-)	-	-
	-	60 ( <i>R</i> )	33 ( <i>S</i> )

**Fig. 1** DIOP, BINAP, and BDPP diphosphine ligands. Summary of the best results obtained in the Ir-catalyzed hydrogenation of several imines


system provides higher enantioselectivities than Ir-BDPP. Later, Claver and coworkers found that using  $[\text{Ir}(\text{COD})((S,S)\text{-BDPP})\text{PF}_6$  (COD = 1,5-cyclooctadiene), at 5 bar of hydrogen and 40°C lead to complete conversion of *N*-( $\alpha$ -methyl-*p*-methoxybenzylidene)-benzylamine. The best enantioselectivity was obtained in the case of the corresponding immobilized catalyst in montmorillonite K-10 (MK-10) providing ee's up to 59% [11].

Diphosphine ferrocene derivatives as the axially chiral (*R,R*)-*f*-binaphane or (*S,S*)-*f*-binaphane (**1**) ligands (Fig. 2) were successfully used in the iridium-catalyzed hydrogenation of different imines using the dinuclear  $[\text{Ir}(\mu\text{-Cl})(\text{COD})_2]$  as iridium source. In the case of acyclic arylimines, the Ir/(*R,R*)-*f*-binaphane (**1a**) catalyst achieve high enantioselectivities (up to 99%). The addition of iodine as additive was required in the hydrogenation of these substrates [14]. The corresponding (*S,S*)-*f*-binaphane was the only ligand, among a number of electron-rich chiral phosphines evaluated, providing high enantioselectivity in the Ir-catalyzed hydrogenation of unfunctionalized N–H imines. The use of a solvent mixture of MeOH/CH<sub>2</sub>Cl<sub>2</sub> (2:1) leads to complete conversion and enantioselectivity up to 95% [15]. The authors proposed a mechanism were the Ir<sup>III</sup>-catalyzed hydrogenation of imines in the presence of I<sub>2</sub> as the additive is possible (Scheme 1). The oxidative addition of I<sub>2</sub> to the Ir<sup>I</sup> precursor **A** generates the Ir<sup>III</sup> complex **B**. The heterolytic cleavage of H<sub>2</sub> can occur in the presence of an amine to form the Ir<sup>III</sup>–H species **C**. An imine substrate can coordinate with **C** in a  $\eta^1$  or  $\eta^2$  fashion. Migratory insertion of the  $\eta^2$ -imine into the Ir<sup>III</sup>–H bond forms an Ir<sup>III</sup>–amide complex. The heterolytic cleavage of H<sub>2</sub> by the Ir<sup>III</sup>–amide intermediate gives an amine and regenerates the Ir<sup>III</sup>–H species **C** [16].

**Fig. 2** f-Binaphane ligands  
**1.** Summary of the best results obtained in the Ir-catalyzed hydrogenation of *N*-arylimines using **1a**




R <sup>1</sup>	R <sup>2</sup>	%ee
Ph	Ph	84
Ph	2,6-Me <sub>2</sub> -C <sub>6</sub> H <sub>3</sub>	>99
4-OMe-C <sub>6</sub> H <sub>4</sub>	2,6-Me <sub>2</sub> -C <sub>6</sub> H <sub>3</sub>	98
4-CF <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	2,6-Me <sub>2</sub> -C <sub>6</sub> H <sub>3</sub>	99



**Scheme 1** Proposed mechanism for the Ir-f-binaphane-catalyzed hydrogenation of imines. S = substrate or solvent

Other 2, 2'-disubstitued biferrocene derivatives (ligands **2**, Fig. 3) also tested using the dinuclear  $[\text{Ir}(\mu\text{-Cl})(\text{COD})]_2$  as the iridium source provided total conversion in arylimine hydrogenation, although with poor-to-moderate enantioselectivity (20–33% ee) [17]. The low enantioselectivities are assumed to be related to the conformational flexibility of this type of bifep ligands.

Cationic iridium complexes with 3,6-bis(diphenylphosphino)hexahydrofuro [3,2-b]furan (**3**) as ligand (Fig. 4) [18], catalyzed asymmetric hydrogenation of several acyclic imines under atmospheric hydrogen pressure, providing chiral amines in quantitative yields and enantioselectivities from 80 to 94%. Contrary to the above-mentioned Ir-diphosphine catalysts, the Ir-**3** complexes formed efficient catalysts under atmospheric hydrogen pressure, whereas at higher pressures, the catalytic activity of the system was drastically reduced. Neutral  $[\text{Ir}(\mu\text{-Cl})(\text{COD})]_2$ -**3** systems were inactive, suggesting that cationic, unsaturated intermediates are preferred with the Ir-**3** catalyst. This was further supported from the pronounced effect that different solvents had on the catalyst's activity. Coordinating solvents such as THF and methanol effectively deactivated the catalyst, whereas in 1,2-dichloroethane, quantitative yields of the secondary amine were obtained.




**2a** R= Ph

**2b** R= 3,5-Me<sub>2</sub>-C<sub>6</sub>H<sub>3</sub>

**2c** R= 3,5-Me<sub>2</sub>-4-OMe-C<sub>6</sub>H<sub>2</sub>

**Fig. 3** Biferrocene diphosphine ligands **2**



**Fig. 4** Diphosphine ligand **3**. Summary of the best results obtained in the Ir-catalyzed hydrogenation of *N*-arylimines

	R <sup>1</sup>	R <sup>2</sup>	%ee
	H	H	84
	OMe	H	80
	Cl	H	81
	H	OMe	94



complete recovery of the starting materials. In an effort to rationalize the stereochemical outcome of the reaction, the authors assumed that the reaction proceeds through a four-membered transition state; namely, the C=N function interacts with the Ir–H bond trans to the Ir–P bond. Two transition states, **A** and **B** (Fig. 7), are possible, in which **A** is more preferable than **B** because there is a large steric repulsion between the N-aryl group (Ar') and the *tert*-butyl group on the phosphorus atom in **B**. It is generally known that unsymmetric imines exist mostly as *anti* isomers, and hence, the reaction via transition state **A** eventually leads to secondary amines with the **R** configuration [19].

Recently, several diphosphines (ligands **5–8**, Fig. 8) were used for the metal-catalyzed asymmetric hydrogenation of exocyclic enamines [20]. The best results were obtained for the MeO-BiPhep (**5**) ligand using the catalytic system [Ir( $\mu$ -Cl)(COD)]<sub>2</sub>/**5**/I<sub>2</sub> with up to 96% enantioselectivity at 50°C and 20 bar of pressure (Fig. 9). This system had previously been applied with excellent results, full conversion, and ees up to 96% at 50 bars, for the hydrogenation of quinolines [21]. Later on, the scope of this Ir/**5** system has been extended to the hydrogenation of 2-benzylquinolines and 2-functionalized and 2,3-disubstituted quinolines, achieving ees up to 96% [22]. Substituted 1,2,3,4-tetrahydroquinolines are an




Fig. 7 Possible transition states leading to secondary amines







Fig. 8 Diphosphine ligands **5–8** applied in the Ir-catalyzed asymmetric hydrogenation of exocyclic enamines




**5**



R <sup>1</sup> /R <sup>2</sup>	%ee
H/Ph	96
H/Me	94
H/n-C <sub>3</sub> H <sub>7</sub>	85
Me/Ph	90
F/Ph	96



R/Ar	%ee
H/C <sub>6</sub> H <sub>5</sub>	94
H/2-MeC <sub>6</sub> H <sub>4</sub>	95
H/1-C <sub>10</sub> H <sub>7</sub>	95
Me/C <sub>6</sub> H <sub>5</sub>	95
F/C <sub>6</sub> H <sub>5</sub>	96



R <sup>1</sup> /R <sup>2</sup>	%ee
H/COPh	96
H/COMe	90
H/(n-Pr)	95
Me/COPh	94
F/COPh	96

**Fig. 9** Summary of the best results obtained in the Ir-catalyzed hydrogenation of quinolines using ligand **5**


important class of biologically active compounds [23]. Mechanistic studies revealed that the hydrogenation mechanism of quinoline could involve a 1,4-hydride addition, isomerization, and 1,2-hydride addition, and the catalytic active species may be a Ir(III) complex with chloride and iodide [24].

Chan and coworkers developed a new diphenylphosphine **9**, related to MeO-BiPhep **5** (Fig. 10) [21]. [Ir( $\mu$ -Cl)(COD)]<sub>2</sub>/**9**/I<sub>2</sub> catalytic system provided similar enantioselectivities than [Ir( $\mu$ -Cl)(COD)]<sub>2</sub>/**5**/I<sub>2</sub> in the Ir-catalyzed hydrogenation of quinolines but higher enantioselectivities in the reduction of 2-methyl-quinoxaline and 2,3,3-trimethylindolenine (Fig. 10).

## 2.2 Bidentate Diposphinite, Phosphite–Phosphinite, Diposphite and Phosphine–Phosphoroamidite Ligands


Furanoside diposphinite ligands **10** and **11** (Fig. 11) were applied in the Ir-catalyzed asymmetric hydrogenation of several dehydroaminoacid derivatives [25]. The best enantioselectivities (ee's up to 78%) were obtained in the reduction of methyl *N*-acetamidoacrylate with ligand **10**. These results using the Ir/**10**–**11** catalysts precursor show that enantiomeric excesses are strongly dependent on the absolute configuration of the C-3 stereocenter of the carbohydrate backbone. The best enantioselectivity were therefore obtained with ligand **10** with an





**Fig. 10** Diphosphine ligand **9**. Comparison of the best results obtained in the Ir-catalyzed hydrogenation of several substrate types using ligands **5** and **9**


**Fig. 11** Summary of the best results obtained in the Ir-catalyzed hydrogenation using furanoside diphosphinite ligands **10** and **11**




*S* configuration of C-3 of the carbohydrate backbone. Interestingly, in the case of using Rh/**10–11** catalysts systems, the best enantioselectivity were obtained with ligand **11**.

Later, furanoside diphosphinite ligands **12**, but with  $C_2$ -symmetry, provided enantioselectivities up to 70% in the asymmetric hydrogenation of several *N*-arylimines (Fig. 12) [26]. The electronic effect of the ligand on enantioselectivity is considerable. Results were best when electron-donating groups are present on the phenyl rings (ligand **12b**). On the other hand, the use of additives was detrimental to both conversion and enantioselectivity.

The furanoside phosphite–phosphinite ligand **13**, related to the previously diphosphinite ligands **12**, was also applied to the asymmetric hydrogenation of *N*-arylimines, increasing the enantioselectivities up to 76% (Fig. 13) [26]. Precursors based on cationic  $[\text{Ir}(\text{cod})_2]\text{BF}_4$  provided better yields and enantioselectivities than




**Fig. 12** Summary of the best results obtained in the Ir-catalyzed hydrogenation using furanoside C<sub>2</sub>-diphosphinite ligands **12**




**Fig. 13** Summary of the best results obtained in the Ir-catalyzed hydrogenation using furanoside phosphinite–phosphinite ligand **13**

the neutral ones formed from the dimer  $[\text{Ir}(\mu\text{-Cl})(\text{COD})]_2$ . The use of benzylamine and phthalimide was beneficial in terms of enantioselectivity.

One of the challenges in hydrogenation is the reduction of substituted aromatic substrates. This transformation cannot be performed using molecular catalysts. Ru-nanoparticles have shown to be catalytically active [27]. In this aspect, iridium nanoparticles ( $[\text{Ir}(\text{THF})_x(\text{L})_y(\text{BF}_4)_z]$ , **L** = diphosphite ligands) stabilized with furanoside diphosphite ligands **14** and **15** (Fig. 14) have recently been applied in the asymmetric hydrogenation of *ortho*-methylanisole and *meta*-methylanisole with high *cis*-selectivity (up to 100%) but low enantioselectivity [28]. Although both diphosphite ligands stabilized iridium nanoparticles with spherical shape and a narrow size distribution, nanoparticles stabilized with ligand **14** were found to be slightly smaller than those bearing ligand **15**. This fact correlates with the higher



**Fig. 14** Summary of the best results obtained in the Ir-catalyzed hydrogenation using nanoparticles stabilized with furanoside diphosphite ligands **14** and **15**



**Fig. 15** Phosphine–phosphoramidite ligands **16–18**. Summary of the best results obtained in the Ir-catalyzed hydrogenation quinolines using ligand **18**

activity obtained with Ir-nanoparticles stabilized with ligand **14** than the ones with ligand **15** and is due to its concomitant higher surface area. It should be pointed out that rhodium and ruthenium nanoparticles provide higher reaction rates than the iridium ones.

A set of novel phosphine–phosphoramidite ligands **16–18** (Fig. 15) possessing two elements of chirality have been prepared through a modular synthetic approach and successfully applied in the Ir-catalyzed asymmetric hydrogenation of

2-substituted quinolines. Particularly remarkable are the enantiomeric excesses up to 97% achieved in the iridium-catalyzed hydrogenation of 2-substituted quinolines, where (1*bS*)-*N*-[2-(diphenylphosphino)phenyl]-*N*-[(*S*)-1-(naphthalen-1-yl)ethyl]-8,9,10,11,12,13,14,15-octahydrodinaphtho[2,1-*d*:1',2'-*f*] [1,3,2]dioxaphosphepin-4-amine **18** proved to be the most selective ligand. Substantially lower enantioselectivities were obtained with the mismatched diastereomer **17b** and with the *N*-phenyl-substituted ligand **16b**, missing a second element of chirality [29].

### 3 Application of Monodentate Phosphorus Ligands


#### 3.1 Monodentate Phosphite, Phosphoramidite, and Phosphorothioite Ligands

Bidentate chiral ligands were considered superior over monodentate ones in metal-catalyzed asymmetric hydrogenation for more than 30 years [1, 2, 5–7] as chelation was believed to be necessary to impart the rigidity to the metal complex for an efficient transfer of chirality. Recently, chiral monodentate phosphines, phosphonites, phosphoramidites, and phosphites were reported to lead to excellent results in the asymmetric hydrogenation of  $\alpha$ - and  $\beta$ -dehydroamino acids, itaconic acid derivatives, and enamides [30–33]. Monodentate phosphite and phosphoramidite ligands have the advantage of being readily accessible, highly diverse, air stable, and inexpensive compared to most bidentate ligands [31, 34, 35]. In addition, they are amenable to parallel synthesis [36–38].


Binol-derived phosphoramidite PipPhos (**19**) has been successfully used as a ligand for the Ir-catalyzed asymmetric hydrogenation of 2- and 2,6-substituted quinolines [39], 2- and 2,6-substituted quinoxalines [40], and *N*-aryl imines [41] (Fig. 16).

In the hydrogenation of quinolines, the presence of tri-*ortho*-tolylphosphine and/or chloride salts as additives was highly adventitious, increasing the enantioselectivities up to 89%. NMR studies indicated that no mixed complexes are formed upon addition of tri-*ortho*-tolylphosphine [39]. Moreover, in order to examine the stability of the catalyst during the reaction, the enantioselectivity was monitored over time. An increase of the enantioselectivity over 24 h was observed. The low ee at the beginning of the reaction may be explained by the slow formation of the catalytically active species during the first hour (10% of conversion). This is also confirmed by the induction time that is observed in the hydrogenation. So, pre-stirring the iridium complex in the presence of the ligand under the reaction conditions during 1 h is necessary to achieve the highest levels of enantioselectivity.

In the hydrogenation of 2- and 2,6-substituted quinoxalines, the presence of piperidine hydrochloride as additive full conversions and enantioselectivities of up to 96% were obtained [40]. These results represent the highest selectivity reached for this class of heterocyclic compounds reported as yet [22, 42, 43].




**19 (S)-PipPhos**




R <sup>1</sup>	R <sup>2</sup>	%ee
Me	H	89
Et	H	88
i-Pr	H	89
i-Bu	H	86
Me	Me	85
Me	OMe	82
n-pentyl	H	83



R <sup>1</sup>	R <sup>2</sup>	%ee
Me	H	96
Et	H	80
i-Bu	H	80
nBu	H	82
n-pentyl	H	85
Ph	H	86
Me	Cl	88



R <sup>1</sup>	R <sup>2</sup>	Ar	%ee
Ph	Me	Ph	87
Ph	Me	2-OMe-Ph	97
Ph	Me	3,5-Me <sub>2</sub> -Ph	>99
2-Naphthyl	Me	2-OMe-Ph	99
4-Me-Ph	Me	2-OMe-Ph	98
4-CF <sub>3</sub> -Ph	Me	2-OMe-Ph	97
Ph	Pr	2-OMe-Ph	97


**Fig. 16** Summary of the best results obtained in the Ir-catalyzed hydrogenation of quinolines, quinoxalines, and *N*-aryl imines using binol-derived phosphoramidite PipPhos **19** ligand

Ir-PipPhos catalysts lead also to very high enantioselectivity in the asymmetric reduction of several *N*-aryl acetophenone imines [41]. The reaction is strongly solvent dependent: in protic solvents such as methanol, no reaction was observed. Excellent conversions and high enantioselectivities were obtained in toluene and dichloromethane. It was also observed that pressures above 5 bar caused a slight decrease in enantioselectivity and shorter reaction times. An interesting anion effect was observed using different iridium precursors, with cationic  $[\text{Ir}(\text{COD})_2]\text{BARF}$  giving the best results. High reaction rates were observed with  $[\text{Ir}(\text{COD})_2]\text{PF}_6$  (full conversion in 30 min); however, the enantioselectivity remained at 65%. No conversion was observed using neutral  $[\text{Ir}(\mu\text{-Cl})(\text{COD})]_2$  as catalyst precursor at room temperature and 5 bar; however, at 50 bar and 60°C, the reaction goes to completion, yielding the product amine with 61% ee. In addition, it was also observed that the presence of electrodonating substituents in the 2-, 3- and 5-position of the aryl ring enhances enantioselectivities up to >99%. If the substituent is 2-methoxy, the resultant secondary amines are easily oxidatively deprotected using trichloroisocyanuric acid to give the primary amines in good yield with full retention of enantioselectivity.

In situ-generated iridium complexes containing bulky 3,3'-substituted H8-phosphoramidites **20** as ligands catalyzed the asymmetric hydrogenation of


nonfunctionalized enamines in high yield and good-to-excellent selectivities (Fig. 17). Enantioselectivities are positively affected by addition of noncoordinating salts (i.e., NaClO<sub>4</sub>, NaBF<sub>4</sub>, etc.). The authors also found an important effect of the substituents of the biaryl group on the catalytic performance. Therefore, activities and enantioselectivities were best using ligand **20c** [44]. These results represent the highest selectivity reached in the Ir-catalyzed hydrogenation of enamides reported as yet. So far, only two studies on this topic have been reported [45, 46], and the best enantioselectivity obtained was up to 60% [46].

The spiro-phosphoroamidite ligand **21** (Fig. 18) was successfully applied in the enantioselective hydrogenation of cyclic *N,N*-dialkyl enamines for the synthesis of optically active cyclic tertiary amines, including those having a six-membered ring or a fused ring (ee's up to 94%) [47]. The author used this asymmetric hydrogenation reaction to synthesize the isoquinoline alkaloid crispine A [48], which was isolated from *Carduus crispus* (welded thistle) and has a significant cytotoxic activity [49, 50]. The asymmetric hydrogenation of enamine **22** performed under optimized conditions using the catalyst Ir/**21** produced crispine A in 97% yield with 90% ee (Scheme 2).




R <sup>1</sup>	R <sup>2</sup>	%ee
Ph	H	93
4-OMe-Ph	H	88
3-OMe-Ph	H	82
2-OMe-Ph	H	64
4-CF <sub>3</sub> -Ph	H	70
t-Bu	H	73
Ph	Me	29

**Fig. 17** Summary of the best results obtained in the Ir-catalyzed hydrogenation of enamides using H8-binol-monophosphoroamidite ligands **20c**




R <sup>1</sup>	R <sup>2</sup>	%ee
Ph	Me	94
Ph	i-Pr	96
4-OMe-C <sub>6</sub> H <sub>4</sub>	Me	94
4-F-C <sub>6</sub> H <sub>4</sub>	Me	92

**Fig. 18** Summary of the best results obtained in the Ir-catalyzed hydrogenation of cyclic *N,N*-dialkyl enamines using spiro-phosphoroamidite ligand **21**



**Scheme 2** Synthesis of Crispine A through asymmetric hydrogenation





**Fig. 19** Combination of ligand **23** with several achiral monodentated P-ligands (phosphites, phosphonites, and phosphines)

Reetz and coworkers reported the use of ligand **23** in combination with several achiral monodentated P-ligands (phosphites, phosphonites, and phosphines) for the asymmetric Ir-catalyzed hydrogenation of imines (Fig. 19). The combination of ligand **23** with the achiral triphenylphosphine leads to remarkable efficient catalyst system reducing imines with enantioselectivities up to 92% (Scheme 3) [51].

Binol-derivative monodentate monophosphorothioate (**24**) and its iridium complex was covalently anchored to mesoporous silica supports (i.e., SBA-15, MCM-41, etc.). Among the catalysts investigated, iridium complex immobilized over SBA-15 (IrPSSBA-15) was found to be the ideal heterogeneous catalyst system for the enantioselective hydrogenation of itaconic acid derivatives. High conversions (99%) and excellent enantioselectivities (up to 96% ee) were observed under

**Scheme 3** Summary of the best results obtained in the Ir-catalyzed hydrogenation of imines using the combination of ligand **23** and triphenylphosphine



**Fig. 20** Monodentate monophosphorothioite ligand **24**

milder reaction conditions (Fig. 20) [52]. It was found that the catalytic activity and enantioselectivity were comparable to its homogenous analog. The catalyst was reused at least five times and the same activity and selectivity were maintained.

### 3.2 Monodentate Phosphines

Considering the most common ligand, triphenylphosphine iridium(I) cationic systems  $[\text{Ir}(\text{COD})(\text{PPh}_3)_2]\text{X}$  were used as catalysts, both homogeneous and heterogenized in smectite clay materials (montmorillonite MK-10, sodium bentonite and lithium hectorite). Both homogeneous and heterogenized  $[\text{Ir}(\text{COD})(\text{PPh}_3)_2]\text{X}$  ( $\text{X} = \text{PF}_6^-$ ,  $\text{BF}_4^-$ ) systems showed high catalytic activity in the hydrogenation of the imine *N*-benzylidene aniline. In particular, the immobilization of the organometallic complex  $[\text{Ir}(\text{COD})(\text{PPh}_3)_2]\text{BF}_4$  in the supports studied afforded catalytic systems that could be reused for a large number of runs, with negligible loss of the catalytic activity (>97% conversion) for at least 13 runs [53]. As described above (Sect. 3.1), achiral monophosphines as triphenylphosphine and tri-*ortho*-tolylphosphine have been combined with other monodentate chiral ligand as phosphoramidite, improving both the conversion and enantioselectivity [51].



## 4 Conclusions

Iridium complexes containing phosphorus ligands have shown to be active catalysts for hydrogenation reaction. Mono- and bidentate phosphines, phosphinites, phosphites, phosphoramidites, and a combination of these functionalities have been applied in the iridium-catalyzed hydrogenation. In the early days, diphosphines were the most widely used phosphorus ligands. However, they are generally difficult to synthesize and prone to oxidation. In recent years, chiral phosphites, phosphinites, phosphonites, and phosphoramidites have emerged as new types of ligands, which present several advantages and high catalytic activity and selectivity. Homogeneous and heterogeneous systems have been studied, allowing the recycling of catalysts. Slight modifications in the electronic and steric properties of the ligands influence the catalytic results, which indicate the potentiality of modular ligands. Imines, including unfunctionalized N–H imines and cyclic imines as quinolines and other substrates, have been transformed in the corresponding chiral amines with ee higher than 95% in many cases. Interestingly mild reaction conditions have been used for these hydrogenations that have been previously performed at high pressures. Binol-derived phosphoramidite constitute one of the most successful ligands in these asymmetric reaction, achieving, in some cases, practically total enantioselectivity.

**Acknowledgments** We thank the Spanish Government (Consolider Ingenio CSD2006-0003, 2008PGIR/07 to O. Pàmies, 2008PGIR/08 to M. Diéguez and ICREA Academia award to M. Diéguez) and the Catalan Government (2009SGR116) for financial support.

## References

1. Cornils B, Herrmann WA (eds) (2002) Applied homogeneous catalysis with organometallic compounds, 2nd edn. VCH, Weinheim
2. de Vries JG, Elsevier CJ (eds) (2007) The handbook of homogeneous hydrogenation. VCH, Weinheim
3. Oro LA, Claver C (eds) (2009) Iridium complexes in organic synthesis. VCH, Weinheim
4. Börner A (ed) (2008) Trivalent phosphorus compounds in asymmetric catalysis, synthesis and applications. VCH, Weinheim
5. Jacobsen EN, Pfaltz A, Yamamoto H (eds) (1999) Comprehensive asymmetric catalysis, vol 1. Springer, Berlin, Chapters 5 and 6
6. Blaser HU, Schmidt E (eds) (2004) Asymmetric catalysis on industrial scale: challenges, approaches and solutions. VCH, Weinheim
7. Ojima I (ed) (2010) Catalytic asymmetric synthesis, 3rd edn. VCH, Weinheim
8. Claver C, Fernandez E (2008) In: Anderson P (ed) Modern reduction methods. VCH, Weinheim, Chapter 3
9. Morimoto T, Nakajima N, Achiwa K (1995) Synlett 748
10. Sablong R, Osborn JA (1996) Tetrahedron: Asymmetry 7:3059
11. Margalef-Català R, Claver C, Salagre P, Fernandez E (2000) Tetrahedron: Asymmetry 11:1469
12. Moser H, Rihs G, Sauter HZ (1982) Naturforsch 37B:451
13. Ciba-Geigy Ltd (1972) US Patents 3,937,730 and 4,022,611

14. Xiao D, Zhang X (2001) *Angew Chem Int Ed* 40:3425
15. Hou G, Gosselin F, Li W, McWilliams JC, Sun Y, Weisel M, O'Shea PDC, Chen I, Davies W, Zhang X (2009) *J Am Chem Soc* 131:9882
16. Fryzuk MD, MacNeil PA, Rettig SJ (1987) *J Am Chem Soc* 109:2803
17. Espino G, Xiao L, Puchberger M, Mereiter K, Spindler F, Manzano BR (2009) *Dalton Trans* 2751
18. Dervisi A, Carcedo C, Ooi L (2006) *Adv Synth Catal* 348:175
19. Imamoto T, Iwadate N, Yoshida K (2006) *Org Lett* 8:2289
20. Wang X-B, Wang DW, Lu SM, Yu CB, Zhou YG (2009) *Tetrahedron: Asymmetry* 20:1040
21. Wang WB, Lu SM, Yang PY, Han XW, Zhou YG (2003) *J Am Chem Soc* 125:10536
22. Qiu L, Kwong FY, Wu J, Lam WH, Chan S, Yu W-Y, Li Y-M, Guo R, Zhou Z, Chan ASC (2006) *J Am Chem Soc* 128:5955
23. Katritzky AR, Rachwal S, Rachwal B (1996) *Tetrahedron* 52:15031
24. Wang DW, Wang XB, Wang DS, Lu SM, Zhou YG, Li YX (2009) *J Org Chem* 74:2780
25. Guimet E, Diéguez M, Ruiz A, Claver C (2004) *Tetrahedron: Asymmetry* 15:2247
26. Guiu E, Aghmiz M, Díaz Y, Claver C, Meseguer B, Militzer C, Castellón S (2006) *Eur J Org Chem* 627
27. Roucoux A, Schulz J, Patin H (2002) *Chem Rev* 102:3757
28. Gual A, Godard C, Philippot K, Chaudret B, Denicourt-Nowicki A, Roucoux A, Castellón S, Claver C (2009) *ChemSusChem* 2:769
29. Eggenstein M, Thomas A, Theuerkauf J, Franció G, Leitner W (2009) *Adv Synth Catal* 351:725
30. Jerphagnon T, Renaud J-L, Bruneau C (2004) *Tetrahedron: Asymmetry* 15:2101
31. Reetz MT (2003) *Chim Oggi* 21:5
32. van den Berg M, Feringa BL, Minnaard AJ (2007) In: de Vries JG, Elsevier CJ (eds) *The handbook of homogeneous hydrogenation*. VCH, Weinheim, p 995
33. de Vries JG (2005) In: Ager DJ (ed) *Handbook of chiral chemicals*, 2nd edn. CRC, Boca Raton, FL, p 269
34. Minnaard AJ, Feringa BL, Lefort L, de Vries JG (2007) *Acc Chem Res* 40:1267
35. Diéguez M, Pàmies O, Ruiz A, Claver C (2004) *Phosphite ligands in asymmetric hydrogenation*. In: Malhotra SV (ed) *Methodologies in asymmetric catalysis*, vol 880. ACS, Washington, DC, p 161
36. Lefort L, Boogers JAF, de Vries AHM, de Vries JG (2004) *Org Lett* 6:1733
37. Swennenhuis BHG, Chen R, van Leeuwen PWNM, de Vries JG, Kamer PCJ (2008) *Org Lett* 10:989
38. Mata Y, Diéguez M, Pàmies O, Woodward SJ (2006) *Org Chem* 71:8159
39. Mršić N, Lefort L, Boogers JAF, Minnaard AJ, Feringa BL, de Vries JG (2008) *Adv Synth Catal* 350:1081
40. Mršić N, Jerphagnon T, Minnaard AJ, Feringa BL, de Vries JG (2009) *Adv Synth Catal* 351:2549
41. Mršić N, Minnaard AJ, Feringa BL, de Vries JG (2009) *J Am Chem Soc* 131:8358
42. Murata S, Sugimoto T, Matsuura S (1987) *Heterocycles* 26:763
43. Bianchini C, Barbaro P, Scapacci G, Farnetti E, Graziani M (1998) *Organometallics* 17:3308
44. Erre G, Enthaler S, Junge K, Addis D, Beller M (2009) *Adv Synth Catal* 351:1437
45. Broady SD, Martin DMG, Lennon IC, Ramsden JA, Muir JC (2006) Patent WO2006067412 A1
46. Marie P, Deblon S, Breher F, Geier J, Böhrer C, Rüegger H, Schönberg H, Grützmacher H (2004) *Chem Eur J* 10:4198
47. Hou G-H, Xie J-H, Yan P-C, Zhou Q-L (2009) *J Am Chem Soc* 131:1366
48. Wu TR, Chong JM (2006) *J Am Chem Soc* 128:9646
49. Zhang Q, Tu G, Zhao Y, Cheng T (2002) *Tetrahedron* 58:6795
50. Knölker H-J, Agarwal S (2005) *Tetrahedron Lett* 46:1173
51. Reetz MT, Bondarev O (2007) *Angew Chem Int Ed* 46:4523
52. Sahoo S, Kumar P, Lefebvre F, Halligudi SB (2008) *J Catal* 254:91
53. Claver C, Fernández E, Margalef-Catalán R, Medina F, Salagre P, Sueiras JE (2001) *J Catal* 201:70



# Iridium-Catalyzed Asymmetric Hydrogenation of Olefins with Chiral N,P and C,N Ligands

David H. Woodmansee and Andreas Pfaltz

**Abstract** Asymmetric hydrogenation is one of the most important catalytic methods for the preparation of optically active compounds. For a long time, chiral rhodium- and ruthenium-diphosphine complexes were the catalysts of choice in this field. However, their application range in the asymmetric hydrogenation of olefins is limited to substrates bearing a coordinating group adjacent to the C=C bond. These limitations were surmounted with the advent of iridium complexes with chiral N,P-ligands. For a wide range of unfunctionalized olefins, excellent enantioselectivities can now be achieved with low catalyst loadings. Moreover, these catalysts also allow highly enantioselective hydrogenation of various functionalized olefins, heterocycles such as furans or indoles, and imines. Research in this field is productive with new catalysts giving high enantioselectivities for difficult substrates appearing at an ever more frequent rate. These methods greatly expand the repertoire of reactions available to synthetic chemists for the preparation of optically active compounds in a reliable and an atom-economical fashion.

**Keywords** 1,1'-Disubstituted alkenes, Asymmetric catalysis, Asymmetric hydrogenation, Imine, Iridium N,P complexes, Tetrasubstituted alkenes, Trisubstituted alkenes

## Contents

1	Introduction .....	32
2	Mechanistic Studies .....	33
2.1	Initial Studies: An Unexpected Anion Effect .....	33
2.2	NMR Investigations of Iridium PHOX Hydride Complexes .....	35
2.3	Computational Studies and Additional Experiments .....	36
3	Asymmetric Hydrogenation of Trisubstituted Alkenes .....	39
3.1	Asymmetric Hydrogenation of Standard Test Substrates .....	39

3.2	Asymmetric Hydrogenation of Purely Alkyl Substituted Olefins	46
3.3	Asymmetric Hydrogenation of Fluorinated Olefins	48
3.4	Asymmetric Reduction of Vinylboronates	49
3.5	Diastereoselective Reduction of Alkenes	50
3.6	Redox Rearrangement of Allylic Alcohols to Chiral Aldehydes	51
3.7	Conjugate Reduction	52
4	Asymmetric Hydrogenation of 1,1'-Disubstituted Olefins	56
5	Asymmetric Hydrogenation of Tetrasubstituted Olefins	59
6	Asymmetric Hydrogenation of Trisubstituted Olefins with Heteroatoms	59
6.1	Enol Esters and Ethers	61
6.2	Asymmetric Hydrogenation of Furans and Chromenes	64
6.3	Asymmetric Hydrogenation of Enamines and Indoles	65
6.4	Asymmetric Hydrogenation of Quinolines and Pyridines	68
7	Asymmetric Hydrogenation of Imines	69
	References	74

## 1 Introduction

The development of highly enantioselective rhodium–diphosphine catalysts in the early 1970s marked the beginning of a new era in asymmetric synthesis. For the first time, practically useful enantioselectivities could be obtained with synthetic chiral catalysts. The well-known *L*-Dopa process developed by Monsanto at that time [1] demonstrated that these catalysts can be applied on an industrial scale, and since then, hydrogenation has played a dominant role in industrial asymmetric catalysis [2]. Today, asymmetric hydrogenation remains a corner stone of the modern organic chemists' repertoire of reliable catalytic methods for the construction of optically active compounds [3, 4]. High enantioselectivity, low catalyst loadings, essentially quantitative yields, perfect atom economy, and mild conditions are attractive features of this transformation as evident in the ever growing list of publications using these methods.

A plethora of chiral phosphine ligands which induce very high enantioselectivity in rhodium- and ruthenium-catalyzed hydrogenations are known. However, the range of olefins that can be hydrogenated with high enantiomeric excess is still limited. Both rhodium and ruthenium catalysts require the presence of a coordinating functional group adjacent to the C=C bond, hydrogenation of dehydro amino acid derivatives or allylic alcohols being typical substrate classes. A caveat of this reactivity in regard to unfunctionalized olefins is that these catalysts generally display low reactivity and unsatisfactory enantioselectivity. Thus, their application has been restricted largely to certain classes of properly functionalized substrates.

Some years ago the Pfaltz group discovered a new class of chiral iridium N,P-ligand complexes which overcame the limitations of the rhodium- and ruthenium-based systems [5, 6]. Early transition metal metallocenes catalysts capable of asymmetric hydrogenation of a range of unfunctionalized olefins in

excellent enantioselectivities have been reported but under onerous conditions [7–9]. Moreover, iridium N,P-based systems showed exceptionally high activity in the hydrogenation of unfunctionalized tri- and even tetrasubstituted olefins. In this respect, they resembled the Crabtree catalyst,  $[\text{Ir}(\text{pyridine})(\text{Cy}_3\text{P})(\text{COD})]\text{PF}_6$  (Cy = cyclohexyl, COD = cyclooctadiene) [10], which provided the stimulus for this work. In addition, promising results were also obtained with certain functionalized alkenes for which no suitable catalysts were available. In this chapter, we discuss the special properties and scope of these catalysts with special emphasis on recent developments [11–14].


## 2 Mechanistic Studies

### 2.1 Initial Studies: An Unexpected Anion Effect

Initial studies with iridium complexes derived from chiral phosphinooxazolines (PHOX ligands) and (*E*)-1,2-diphenyl-1-propene as substrate gave encouraging results (Scheme 1) [5, 15]. With 4 mol% of catalyst ( $\text{X} = \text{PF}_6^-$ ) at 10–50 bar hydrogen pressure, up to 98% ee could be obtained. However, the turnover numbers were disappointingly low.

Kinetic studies demonstrated that with 4 mol% of catalyst in a 0.3 M solution of olefin at 7 bar hydrogen pressure, the reaction was extremely fast and reached completion within less than 1 min [16]. Lower catalyst loadings resulted in decreased conversion. Although the initial rate was still high at 1 mol% catalyst loading, a rapid and essentially complete deactivation of the catalyst was observed before 50% of the olefin was consumed. Deactivation is a known problem of the Crabtree catalyst, which is attributed to the formation of inactive hydride-bridged trinuclear complexes [10]. In the case of Ir(PHOX) complexes as well, NMR analysis of deactivated reaction mixtures suggested the presence of such hydride-bridged species. In subsequent studies, a trinuclear Ir(PHOX)-hydride complex was isolated and characterized by NMR and X-ray analysis [17]. This complex proved to be remarkably stable, and all experiments to convert it back into a catalytically active species failed.

Attempts to increase conversion by variation of the solvent, hydrogen pressure, or the catalyst and substrate concentration were unsuccessful. Coordinating solvents and additives such as amines, or coordinating anions such as halides, carboxylates,




**Scheme 1** Initial screening with PHOX


and even the very weakly coordinating triflate ion were found to deactivate the catalyst. The best results were obtained in anhydrous dichloromethane or 1,2-dichloroethane using cationic Ir-PHOX complexes with hexafluorophosphate as counterion. Rigorous exclusion of moisture and oxygen resulted in increased conversion. When the reaction was set up in carefully dried dichloromethane in a glove box, full conversion could be achieved with only 0.5 mol% of the catalyst. However, reactions at such low catalyst loadings were difficult to reproduce.

After extensive experimentation, a simple solution for avoiding catalyst deactivation was discovered, when testing an Ir-PHOX catalyst with tetrakis[3,5-bis(trifluoromethyl)phenyl]borate ( $\text{BAr}_F^-$ ) as counterion [5]. Iridium complexes with this bulky, apolar, and extremely weakly coordinating anion [18] did not suffer from deactivation, and full conversion could be routinely obtained with catalyst loadings as low as 0.02 mol% [19]. In addition, the  $\text{BAr}_F^-$  salts proved to be much less sensitive to moisture than the corresponding hexafluorophosphates. Tetrakis(pentafluorophenyl)borate and tetrakis(perfluoro-tert-butoxy)aluminate were equally effective with very high turnover frequency, whereas catalysts with hexafluorophosphate and tetrafluoroborate gave only low conversion while reactions with triflate were completely ineffective (Fig. 1).

How can these bulky, extremely weakly coordinating anions prevent catalyst deactivation? A comparative kinetic study of catalysts with different anions provided a plausible answer [19]. With  $\text{PF}_6^-$  as a counterion, the rate dependence on olefin concentration was first order, whereas the rate order observed for the corresponding  $\text{BAr}_F^-$  complex was close to zero. This striking difference may be explained by the stronger coordination of  $\text{PF}_6^-$  or formation of a tight anion pair, which slows down the addition of the olefin to the catalyst to such an extent that it becomes rate-limiting. In contrast, the essentially noncoordinating  $\text{BAr}_F^-$  ion does not interfere with olefin coordination, and the catalyst remains saturated with olefin even at low substrate concentration. The slower reaction of the  $\text{PF}_6^-$  salt with the olefin could



**Fig. 1** Order of reactivity of the complexes  $[\text{Ir}(\text{PHOX})(\text{COD})]\text{X}$  by TOF measured at 4°C with E- $\alpha$ -methylstilbene as substrate



**Fig. 2** Comparison of Crabtree catalyst with the  $\text{BAr}_F$  analog

explain its higher tendency to undergo deactivation. If we assume that deactivation is caused by the formation of hydride-bridged species leading to an inactive trinuclear complex, then the critical step in the catalytic cycle is the reaction of the Ir-hydride intermediate with the olefin. If olefin insertion is very fast, as in case of the  $\text{BAr}_F^-$  counterion, hydrogenation dominates over the deactivation pathway, whereas with the  $\text{PF}_6^-$  analog, the olefin reacts more slowly and deactivation becomes a significant competing process.

Virtually every iridium catalyst of the formula  $[\text{Ir}(\text{L}^*)(\text{COD})]^+ [\text{X}]^-$  for asymmetric olefin hydrogenation that has appeared after the initial counterion effect studies was based on  $\text{BAr}_F^-$  as the preferred anion [14]. The anion effect is broadly applicable in iridium-catalyzed reductions as experiments with a direct analog of the Crabtree catalyst of the formula  $[\text{Ir}(\text{pyridine})(\text{Cy}_3\text{P})(\text{COD})]\text{BAr}_F$  indicates (Fig. 2).


Hydrogenation of  $\delta$ -terpinene (Fig. 2) proceeded in higher conversion with Crabtree's catalyst with  $\text{BAr}_F$  counter ion rather than the normal  $\text{PF}_6$ . The  $\text{BAr}_F$  counter ion performed better in all instances where the more coordinating  $\text{PF}_6$  salt failed to reach complete hydrogenation [20].

## 2.2 NMR Investigations of Iridium PHOX Hydride Complexes

In an early work of Crabtree and coworkers, olefin dihydride intermediates formed during hydrogenation of cyclooctadiene using  $[\text{Ir}(\text{pyridine})(\text{PCy}_3)(\text{COD})]\text{PF}_6$  in dichloromethane at  $0^\circ\text{C}$  were detected by NMR spectroscopy [21]. In a more complementary study, Mazet et al. found that when  $[\text{Ir}(\text{PHOX})(\text{COD})]\text{BAr}_F$  complex **1** was treated with hydrogen at  $-40^\circ\text{C}$  for 5 min in  $[\text{D}_8]$ -THF, olefin dihydride intermediates were formed, which were characterized by NMR spectroscopy [22]. Two new signals appeared in the hydride region that were assigned to a single dihydride complex **2c** formulated as  $[\text{Ir}(\text{PHOX})(\text{H})_2(\text{COD})]\text{BAr}_F$  (Scheme 2).

The predominance of isomer **2c** over **2a** or **2b** is consistent with Crabtree's findings, which convincingly demonstrated that in the reaction of  $\text{H}_2$  with  $[\text{Ir}(\text{pyridine})(\text{PR}_3)(\text{COD})]\text{PF}_6$ , the formation of an Ir-H bond *trans* to the N ligand is electronically favored [23]. Highly selective formation of isomer **2c** results from  $\text{H}_2$  addition to the





**Scheme 2**  $^1\text{H}$  NMR study of Ir(**5a**) hydride formation

more sterically encumbered face of the starting complex because dihydrogen addition to the sterically more accessible face leading to isomer **2d** would build up steric strain between the chelating COD ligand and the isopropyl group in the oxazoline ring and the pseudoaxial P-phenyl group. When the solution containing complex **2c** was warmed to  $0^\circ\text{C}$  under hydrogen, a gradual consumption of isomer **2c** was observed accompanied by the appearance of two new hydride complexes **3c** and **3d** with concomitant formation of cyclooctane.

### 2.3 Computational Studies and Additional Experiments

DFT (density functional theory) calculations on the complete structures of complexes shown in Scheme 2 have been carried out by Mazet et al. [22, 24, 25]. The


fully minimized structures of the four possible *cis*-dihydrides formed by oxidative addition of H<sub>2</sub> to [Ir(PHOX)(COD)]<sup>+</sup> were calculated. The most stable structure corresponded to the reaction product **2c** that was shown to be formed exclusively in the NMR experiment. Isomers **2a** and **2d** were 10.6 and 4.9 kcal/mol higher in energy, whereas for isomer **2b**, no stable chelate structure could be located due to severe steric interactions, which prevent the formation of an Ir–N bond. The four possible [Ir(PHOX)(H)<sub>2</sub>(solvent)<sub>2</sub>]<sup>+</sup> complexes **3a–d** resulting from the hydrogenation of the cyclooctadiene ligand were also examined, and again, the two most stable structures corresponded to the isomers observed in the NMR experiments. These results show that steric interactions are very important and may dominate over electronic factors. Consequently, computational studies of potential reaction pathways should be based on full catalyst and substrate structures rather than simple model systems.

Unfortunately, attempts to observe and characterize intermediates under catalytic conditions have been unsuccessful so far. When considering which intermediates may be formed during catalysis, one of the first issues that becomes apparent is what ligands are coordinated to iridium during catalysis. An olefin dihydride iridium complex, which incorporates a bidentate N,P ligand, has a sixth coordination site available for an additional ligand. Whereas coordination of a second molecule of olefin seems highly unlikely due to steric hindrance, dihydrogen and dichloromethane may both be effective ligands for iridium.

Thus the two plausible catalytic cycles have been considered, one via an Ir dihydride complex **A** and the other via an IrH<sub>2</sub>(η<sup>2</sup>-H<sub>2</sub>) complex **B** (Fig. 3). The first is analogous to the well-established mechanism for rhodium diphosphine-catalyzed hydrogenation of olefins going through Ir(I) and Ir(III) intermediates [26–29].

Experimental support for an Ir(I)–Ir(III) mechanism was provided by Chen and Dietiker [30]. They reported an elegant experimental investigation of the hydrogenation of styrene with [Ir(PHOX)(COD)]BAR<sub>F</sub> in the gas phase by means of electrospray ionization tandem mass spectrometry. By means of reversible deuterium labeling, the investigators found masses corresponding only to intermediates with a mass corresponding to a dihydride complex with no presence of a trihydride species and concluded that no Ir(V) species with PHOX could be present in the catalytic reaction.


Based on DFT calculations Brandt et al. proposed a catalytic cycle via Ir(III) and Ir(V) intermediates, in which an additional dihydrogen molecule coordinated to an Ir-dihydride undergoes oxidative addition during migratory insertion [31]. However, since an extremely truncated model for the ligand and substrate (ethylene) was used which neglected the severe steric interactions present in the actual catalysts it



**Fig. 3** Possible Ir(III) and Ir(V) hydride intermediates (S solvent)

seems premature to rule out an Ir(I)–Ir(III) cycle. From subsequent calculations on the full catalyst and substrate structures, which were based on the postulated Ir(III)–Ir(V) cycle, a simple qualitative quadrant model was derived for rationalizing the observed enantioselectivities. Further studies by the same investigators on a complete complex with stereoelectronic contributions from both *trans*- $\alpha$ - and  $\beta$ -methylcinnamic esters were investigated and applied to the model (Fig. 4) [32]. The authors concluded that in the case of the *trans*- $\beta$ -methylcinnamic ester, electronic and steric factors cooperated in their model to give high enantioselectivity. In the case of the *trans- $\alpha$ -methylcinnamic ester, steric interactions placed the migrating hydride onto the alpha carbon, which would be electronically disfavored and thus lead to lower selectivities, both substrate selectivity observations being reflected by experimental results.*

Fan et al. also reported DFT calculations on the complete ligand and substrate structures with an iridium carbene-based complex for an Ir(III)–Ir(V) catalytic cycle, which reproduced the correct selectivity order for three different substrates [33]. Calculations on complete ligands and real substrates that reproduce experimentally determined enantioselectivities give plausible credence to these authors' computational experiments.



**Fig. 4** Andersson's quadrant model

Preliminary studies by Meuwly and Roseblade [24, 25] indicated that calculated pathways beginning from  $[\text{Ir}(\text{P}^{\wedge}\text{N})(\text{H})_2(\eta^2\text{-H}_2)(\eta^2\text{-olefin})]^+$  complexes lead to predicted enantioselectivities which are opposite to the experimentally observed values. While competing pathways involving solvated Ir(I)–Ir(III) intermediates were somewhat higher in energy (by a few kcal/mol), calculated transition state energies were in reasonable agreement with the experimentally observed enantioselectivities. Taking into account that  $\text{CH}_2\text{Cl}_2$  as the solvent is present in much higher concentration than  $\text{H}_2$ , the Ir(I)–Ir(III) and Ir(III)–Ir(V) cycles become energetically very similar, making it difficult to distinguish between them based on calculations alone.

Thus, additional experimental and computational studies will be needed to draw definitive conclusions regarding the mechanism of Ir-catalyzed asymmetric hydrogenation. The Ir(I)–Ir(III) and Ir(III)–Ir(V) cycles seem to be similar in energy, so it may well be that depending on the catalyst, substrate, and the hydrogenation conditions, one or the other pathway will be preferred or both cycles could operate in parallel.

### 3 Asymmetric Hydrogenation of Trisubstituted Alkenes

#### 3.1 *Asymmetric Hydrogenation of Standard Test Substrates*

Largely unfunctionalized trisubstituted olefins are the most commonly investigated hydrogenation substrates in iridium-catalyzed reductions [11–14]. As previously stated, strongly coordinating groups such as basic amines, alcohols, and strongly donating Lewis bases can in general slow the iridium catalysts down to a point where they are not effective. However, a number of olefins with moderately Lewis basic functionalities such as acids, alcohols, esters, ketones, ethers, halogens, and other similar groups adjacent or in proximity to the C=C bond have been successfully reduced with very high functional group tolerance and stereoselectivity. There is an increasing appearance of more coordinating functionalities such as enamines, indoles, imines, and even pyridines, all of which will be touched upon in the later sections of this chapter.

A survey of a list of typical ligands with reasonable asymmetric induction for the reduction of a commonly tested set of trisubstituted olefins (Table 1) reveals several common features (Fig. 5). Virtually, all of the ligands consist of a heterocycle with a  $\text{sp}^2$ -hybridized nitrogen atom as a hard  $\sigma$  donor and a strong soft donor moiety such as a trisubstituted phosphorus atom or an N-heterocyclic carbene forming the second portion of the chelating ligand. Six-membered metallacycles are the most investigated, with the exception of the seven-membered carbene ligand **9**, the seven-membered spirocycles SpinPHOX **15a–c**, and the rigid spiroindanes SiPHOX **16a–d**. Stereogenic units have been incorporated with varying degrees of success in all regions of the metallacycle backbone. Although some ligands display a wider

**Table 1** Asymmetric hydrogenation of standard substrates with chiral N,P and C,N iridium catalysts

Substrate	Ligand ee % <sup>a</sup>						
	<b>5</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>9</b>	<b>10</b>	<b>11</b>
 <b>25a</b>	<b>b</b> 97	<b>a</b> 99	<b>a</b> 98	<b>a-c</b> 99	99	<b>a-d</b> 99	<b>a</b> 99
 <b>25b</b>	<b>b</b> 81	<b>a</b> 99	<b>a</b> 91	<b>a-c</b> 99	97	<b>a-d</b> 99	<b>a</b> 90
 <b>25c</b>	<b>b</b> 63	<b>e</b> 92	<b>a</b> 89	<b>a</b> 98	80	–	–
 <b>25d</b>	<b>b</b> 72	<b>f</b> 95	<b>a</b> 95	<b>a</b> 92	–	<b>b</b> 55	<b>b</b> 94
 <b>25e</b>	<b>b</b> 96 (95)	<b>a</b> 92	<b>a</b> 97	<b>a</b> 97	93	<b>b, d</b> 97	<b>b</b> 98 (95)
 <b>25f</b>	<b>b</b> 84 (96)	<b>f</b> 96	<b>a</b> 94	<b>a</b> 99	–	<b>b, d</b> 98	<b>b</b> 93

*(continued)*

**Table 1** (continued)

Substrate	Ligand ee % <sup>a</sup>						
	<b>13</b>	<b>14</b>	<b>17</b>	<b>18</b>	<b>20</b>	<b>21</b>	<b>24</b>
 <b>25a</b>	<b>c</b> 98	<b>a</b> 98	99	<b>a-b</b> 99	<b>b,g</b> 99	<b>a</b> 94 (35)	78
 <b>25b</b>	<b>c</b> 99	<b>a</b> 89	99	<b>b</b> 99	<b>d</b> 94	<b>a</b> 94	–
 <b>25c</b>	–	<b>c</b> 96	92	<b>a</b> 95	<b>f</b> 88	<b>a</b> 43 (18)	–
 <b>25d</b>	<b>c</b> 95	<b>b</b> 96	96	–	<b>e</b> 83	<b>a</b> 97	89 (95)
 <b>25e</b>	–	<b>b</b> 96	93	<b>b</b> 92	–	<b>a</b> 45 (5)	99
 <b>25f</b>	<b>a</b> 88	<b>a</b> 95	99	<b>b</b> 99 (85)	<b>c</b> 98	<b>a</b> 55 (10)	82

<sup>a</sup>All conversions less than 99% are noted under the enantioselectivity in brackets

range of acceptable enantioselectivity, no single catalyst can be said to be generally applicable to all olefins. The current trend is toward good selectivity in a given class of substrates with modifications to many of the existing scaffolds in an attempt to improve selectivity for important targets.

Many similarities to the classic PHOX ligand **5a–c** are clearly visible from the given ligand list in Fig. 5. As indicated by NMR and computational studies (see Sect. 2.2 and 2.3), the coordinated olefin takes a position in a *trans* orientation from the soft donation moiety (i.e., phosphorus or carbene substituent) and adjacent to the nitrogen atom. In the PHOX system, this places the olefin into close proximity with the stereogenic center of the ligand to create a well-defined chiral environment. Ring size of the metallacycle plays an extremely important role in governing catalyst reactivity. Catalysts derived from ligands **8a–f** and **10a–d** are some of the most selective catalysts known for unfunctionalized trisubstituted olefins [32, 34–37].





Fig. 5 continued



**Fig. 5** Frequently used N,P and C,N ligands

While they have proven very selective for trisubstituted olefins, they fail to produce high selectivity or reactivity with the substrate classes of tetrasubstituted olefins, terminal olefins, and 1,3-dienes.

The catalyst derived from ligand **9** perform well with several classes of olefins albeit with slightly lower enantioselectivities in reduction of trisubstituted olefins, but are remarkably reactive toward 1,3-dienes [11, 33]. Further examples of the size



of the iridium metallacycle controlling selectivity and reactivity comes from the substrate profile of the five-membered iridium chelates from ligands **12a–e**, which reduce tetrasubstituted olefins under mild conditions and high conversions but give unacceptable enantioselectivity with less-substituted olefins [13].


On consultation of Table 1, it becomes apparent that small changes in overall geometry about the metal center as well as the donation capabilities to the metal center can have a drastic effect on the selectivity. PHOX ligand **5b** compared with the later permutation of SimplePHOX ligand **7a** has the same 4-tert-butyl-4,5-dihydrooxazole moiety as a stereo-defining group and nearly the same steric environment about the phosphorus atom, yet they give drastically different enantioselectivity. ThrePHOX **6a–f** ligands are highly selective and active catalysts which incorporate effective modifications on the PHOX progenitor. In comparison to both PHOX and the later derivative SimplePHOX, the stereochemistry of ThrePHOX has been moved to the center of the backbone and transmits chirality to both sides of the chelate [38–40].

Other features of note are observed when the phosphinite group of SimplePHOX is replaced by a phosphine in the newer NeophOX ligand **14a–c**, resulting in some small improvements in enantioselectivity but with the added advantages of a phosphine over the more reactive phosphinite [41]. The lack of overall changes to selectivity for these simple substrates may reflect that the alteration in overall geometry, most importantly the bite angle, matters more to reactivity than electronic changes to the soft donor group. Ligands **20a–b** provide reasonable selectivity but actually less than the parent SimplePHOX and at a cost of a more complicated synthesis, a strong indication that optimal interactions were disrupted by the excess chiral and steric encumbrance [42].


Other examples of adding too much bulk in the context of the general substrates come in the form of ligand **24** [43]. While the ligand maintains many of the stereochemical features of ligand **5c**, the added bulk leads to much lower enantioselectivities and conversion by disrupting interactions necessary for substrate binding. Similar effects are seen with ligands **21a–d**, which are closely related to the very active catalysts derived from **8a–c**, **10a–d**, and **11a–b**, with the exception of the deleterious addition of extra ring rigidity and steric encumbrance from the carbon bridge [44].

While Table 1 is an important tool for many of the studies, the current trend is to elaborate more the increasingly difficult substrates within a particular class of olefins. Andersson and coworkers have published several variations of the 1,2-diaryl substrate **25a** for the fairly more difficult 1,1'-diaryl class of trisubstituted olefins (Scheme 3) [45]. The authors note that the products form an important class of compounds that are pharmaceutically relevant and difficult to prepare by other available methods.

Reduction of all of the reported geminal diaryl substrates was completely under the control of the third stereo defining R-group. Particularly difficult were the sterically demanding **29** and the electron poor olefin **30**, both of which required heating to produce poor to moderate yields.



### Scheme 3 Asymmetric hydrogenation of geminal-diaryl substrates



### Scheme 4 Use of Ir N, P based asymmetric hydrogenation in total synthesis

The reduction of simple trisubstituted olefins derived from **25d** has found use in the enantioselective total synthesis of pseudopteroxazole **31c**, demethyl methoxy-calamenene **32c**, and both enantiomers of mutisianthol **33c** (Scheme 4) [41, 46, 47].

SimplePHOX **7a** proved a useful tool to force the diastereomeric reduction of olefin **31a** to pseudopteroxazole precursor **31b** in perfect diastereoselectivity and 90% yield with only trace amounts of over-reduced product. NeoPHOX catalyst from ligand **14b**, a closely related system to **7a**, furnished product **32b** in 93% ee, which was then easily recrystallized to enantiopure material with 58% recovery. The *R* enantiomer of **33b** was synthesized by the use of catalyst from ligand **8a** in 90% ee and 98% yield with the fully aromatized naphthalene as 2% byproduct. A higher catalyst loading of 2 mol% of catalyst from **7a** was used to produce the


*S* enantiomer in 80% ee with 13% of the fully aromatized naphthalene byproduct occurring even when using 50 bar of hydrogen pressure. Lower catalyst loadings gave increased amounts of aromatization byproduct and lower enantioselectivity.

Asymmetric hydrogenation of cyclic olefins is not the sole application of this methodology in synthesis, and the current trend in research is to use these catalysts in building more elaborately functionalized molecules.



### 3.2 Asymmetric Hydrogenation of Purely Alkyl Substituted Olefins

The development of chiral homogeneous catalysts with the capacity to reduce purely alkyl-substituted olefins with high asymmetric induction and yield has been a long-standing problem. Most iridium catalysts that perform well with nothing more coordinating than a phenyl group adjacent to the olefin fail to give good asymmetric induction in the absence of an aromatic substituent in close vicinity to the C=C bond [11–14].

As a first step to solving this problem, Bell et al. started an active screening project utilizing *E* and *Z* isomers of compound **34a** as a model (Scheme 5) [36]. Although **34a** contains an aromatic substituent for analytical purposes, the functional group has been moved three bonds away from the carbon–carbon double bond. If any coordination took place through this remote site the effect was expected to be weak. Out of the screening effort, pyridine phosphinite ligands **8a–c** were identified as excellent ligands giving high enantioselectivity and



**Scheme 5** Test substrates in the asymmetric hydrogenation of purely alkyl substituted alkenes

**Scheme 6** Asymmetric synthesis of  $\gamma$ -tocopherol from the triene**Scheme 7** Total stereocontrol in the asymmetric reduction of farnesol isomers

conversion. Both *E* and *Z* isomers gave acceptable enantioselectivity. As observed in other cases, the *E* and *Z* isomers were converted to opposite enantiomers. In order to rule out any small interactions from the distal aromatic ring, the completely alkyl-substituted olefin **35a** was reduced under standard conditions. Fittingly, ligand **8a** provided the branched chiral alkane **35b** in 92% enantioselectivity.

Because *cis* and *trans* olefins are converted to products of opposite configuration, it becomes possible to introduce two or more stereogenic centers with the desired relative and absolute configuration in a single step through hydrogenation of a di- or polyene by adjusting the geometry of the individual C=C bonds. This is demonstrated by the highly enantio- and diastereoselective preparation of  $\gamma$ -tocopherol **36b**, a component of vitamin E, from  $\gamma$ -tocotrienyl acetate **36a** (Scheme 6).


In a subsequent publication, the fully optimized stereospecific synthesis of each individual diastereomer and its corresponding enantiomer with catalyst loadings of **8c** from 0.1 to 0.25 mol% was reported (Scheme 7) [37]. Electron withdrawing protecting groups attached to the allylic alcohol slowed down the reaction rate and higher catalyst loadings were required to achieve full conversion. The authors conducted a grams scale synthesis of (*R,R*)-**37b** in comparable yield and enantioselectivity to the test reactions, illustrating the practicality of this approach.

### 3.3 Asymmetric Hydrogenation of Fluorinated Olefins


Chiral organofluorides are increasingly in demand as the pharmaceutical and materials industries seek to take advantage of the special properties these halides impart. A lack of methods to provide chiral monofluorides and trifluoromethyl groups has given the group of Andersson impetus to create new asymmetric hydrogenation routes to these valuable halides [48, 49].

A brief scope of  $\alpha$ -fluorocinnamic acid derivatives with both trisubstituted and tetrasubstituted C=C bonds were reduced with mixed results (Scheme 8) [48]. The vinyl fluorides proved extremely difficult for catalysts that usually perform superbly on more frequently studied substrates. Elevated temperatures and pressures were required to obtain moderate to good conversions with closely related substrates. A struggle with dehalogenation occurred with the harsher conditions required for this difficult reaction. The group noticed that phosphoramidate ligands **13a–c** gave lower amounts of dehalogenation and new phosphoramidate **26** was developed by the group to incorporate the best of both of their original ligands (Fig. 6). This catalyst gave improved enantioselectivity and less dehalogenation (Scheme 8).



Evidently, vinylfluorides represent a new challenge to iridium asymmetric hydrogenation. It is noteworthy that no examples of asymmetric reduction of



**Scheme 8** Asymmetric hydrogenation of vinyl fluorides



**Fig. 6** Scaffold morphing to improve dehalogenation profile



**Scheme 10** Asymmetric hydrogenation of vinyl boronic esters

vinyl fluorides without adjacent coordinating groups have been reported in the literature [14].

Trifluoromethyl-substituted olefins appear to also be problematic substrates for asymmetric hydrogenation as very long amounts of time and high pressures were required to achieve appreciable yields with ligand **10b** (Scheme 9) [49]. Nevertheless, useful yields and excellent enantioselectivities were obtained for most examples.

### 3.4 Asymmetric Reduction of Vinylboronates

Boronic esters have been used in a wide range of transformations. These useful reagents have been transformed into numerous functional groups and are essential reagents for several C–C bond-forming reactions. Transition metal-catalyzed hydroboration of olefins often leads to mixtures of branched and linear products. Several groups have reported asymmetric reductions of vinyl boronic esters [50–52] with chiral rhodium P,P complexes; however, the first iridium-catalyzed reduction was reported by Paptchikhine et al (Scheme 10) [53].

In contrast to normal trisubstituted aryl olefins, a strong pressure effect was observed in this case. Interestingly, for substrate **44a** catalysts' hydrogenation with Ir(**13d**) demonstrated opposite pressure-dependant enantioselectivity of substrate **45a** with catalyst Ir(**46**). Poor enantioselectivity was obtained for substrates that did not contain an aromatic ring adjacent to the olefin.


### 3.5 Diastereoselective Reduction of Alkenes

Although functional group-directed enantioselective hydrogenation tends to fall in the realm of rhodium and ruthenium catalysts, there are many examples of diastereoselective hydrogenations with Crabtree's catalyst that are controlled by coordination of the iridium center to a Lewis basic functional group (Fig. 7). When one considers that iridium catalysts do not require coordinating groups for hydrogenation activity and purely steric interactions suffice for enantioface discrimination, then protection of the directing group with a large noncoordinating moiety could reverse the direction of attack by the catalyst to create the opposite diastereomer-protected equivalent (Fig. 7).



Diastereoselective hydrogenations of this type have been reported by Burgess and coworkers [54–59] using chiral-protected and -unprotected allylic and homo allylic alcohols as substrates with their carbene catalyst Ir(**9**). Catalyst control was found to be dominant, but depending on the position and nature of the oxygen substituents, moderate to strong match/ mismatch effects were observed.

This approach has been used to synthesize a variety of valuable 1,3-deoxypolyketide chirons, 1,3-hydroxymethyl chirons, vicinal dimethyl chirons, and aldol-like products from trisubstituted olefins (Scheme 11). High diastereomeric ratios could be obtained often in greater than 90% for most cases.

A reasonably large difference in diastereomeric excess was observed between product **47b** with an adjacent methyl ester and **48b** with a primary alcohol in the equivalent position [57]. It was noted by the authors that in cases involving a 1,3 system, changing the pendant group from a primary allylic alcohol to a methyl ester caused a reversal of facial selectivity [54, 58]. The same effect was absent in the 1,2 systems **51b** and **52b** studied. The diastereomeric ratio in the latter case was attributed mainly to catalyst control [58].



**Fig. 7** Diastereoselective hydrogenation based on catalyst and substrate control

**Scheme 11** Diastereoselective hydrogenation with Ir(**9**)**Scheme 12** Rearrangement of allylic alcohols

### 3.6 Redox Rearrangement of Allylic Alcohols to Chiral Aldehydes

Mazet et al. have reported an efficient asymmetric isomerization reaction of allylic alcohols [60, 61]. In a preliminary report they utilized the BAR<sub>F</sub> analog of Crabtree's complex to efficiently catalyze a hydride transfer from the α position of the allylic alcohol to the β position of the olefin with a concomitant formation of a formyl group. A subsequent report detailed a remarkable enantioselective variant of this process catalyzed with Ir(**12g**) and (**12h**) (Scheme 12).

The iridium catalyst was activated by briefly purging the system with atmospheric pressure of hydrogen to remove the COD ligand followed by degassing the



system with vacuum and inert atmosphere to avoid over reduction. High enantioselectivities were obtained in moderate to good yields with a limited range of substrates. The system studied was very sensitive to ligand geometry the five-membered metallacycles were much more efficient than six-membered analogs and bulky substrates with aromatic substituents were required for high selectivity.

### 3.7 Conjugate Reduction

Catalytic asymmetric conjugate reductions represent one of the most venerable and investigated reaction types in catalysis. Conjugate reductions with rhodium and ruthenium diphosphine-based catalysts have been heavily investigated and substituted  $\alpha,\beta$ -unsaturated acids and esters are frequent substrates for these systems. Iridium N,P-based reductions are more recent and far less investigated but offer chemoselectivity advantages under certain circumstances [62]. Ester **25f** is the most regularly investigated conjugate reduction substrate and has been a general benchmark for evaluating the selectivity and reactivity of iridium N,P-based catalysts for many years. It has also been described that the *trans*- $\alpha$ -methylcinnamic esters have been particularly difficult substrates for reduction, and this has been rationalized by computational methods [32]. Nearly all systems that have been investigated are based on a cinnamic acid core with different substitution patterns about the  $\alpha$  and  $\beta$  carbons. Varying degrees of electrophilic groups have been incorporated to the conjugate system such as ketones, esters, acids, amides, diphenylphosphine oxides, and phosphonates [54–59, 62–67].

Iridium-catalyzed reduction of  $\alpha,\beta$ -unsaturated ketones has been investigated most recently by the groups of Bolm and Hou, respectively.

As Table 2 indicates, ligand **23c** proved effective for substrates with large, branched groups on the beta positions. The best selectivity was obtained for **54c**, which was substituted with aromatic groups on both sides of the substrate, and a large branched alkyl attached to the  $\beta$  position [62]. Entries **54e** and **54f** are encouraging and may indicate that this method could be extended to more broadly useful compounds. The investigators performed a solvent study and determined the reaction to be equally selective in toluene and dichloromethane; so the former was used presumably for its industrial attractiveness.

A broader range of substrates have been reduced with substitution at the  $\alpha$  position of  $\alpha,\beta$ -unsaturated ketones using ligand **5c** [63, 64].

Table 3 indicates that reduction of  $\alpha$  substituted unsaturated ketones is extremely efficient with PHOX **5c**. Unfortunately, substrates lacking any aromatic ring within close proximity to the olefin were reportedly not reduced [64]. However, a wide range of functional groups were shown to be tolerated by Bolm et al. [63]. Most impressively, nitroaromatic compounds were well tolerated under the conditions used, highlighting the extraordinary chemoselectivity that can be obtained with iridium N,P catalysts. Perhaps one of the most productive features of these studies is the ability to create a stereogenic center with high degrees of substitution and

**Table 2** Enantioselective hydrogenations of linear  $\alpha,\beta$ -unsaturated ketones

Substrate	Substrate	Yield %	ee %
<b>54a</b>		89	81
<b>54b</b>		94	89
<b>54c</b>		94	97
<b>54d</b>		86	92
<b>54e</b>		93	81
<b>54f</b>		70	79

perfect acyclic stereocontrol while maintaining an unencumbered site  $\alpha$  to the carbonyl group for further elaboration.

Substituted  $\alpha,\beta$ -unsaturated amides have also been reduced with high enantioselectivity and conversion with ligand **24** [65]. Here as well, the investigations relied heavily on substrates with adjacent aromatic rings and  $\alpha$  substitution. Catalyst loadings were moderately higher than with other systems with 2 mol% used. Table 4 indicates that ligand **24** can tolerate a wide range of substituents on the amide nitrogen but fairs best with the more sterically demanding groups such as isobutyl or benzyl. The use of more coordinating groups such as methoxyethyl **56e** or the versatile Weinreb amide **56f** results in a moderate to severe erosion of enantioselectivity.

**Table 3** Enantioselective hydrogenations of  $\alpha$ -substituted  $\alpha,\beta$ -unsaturated ketones

Substrate	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Yield (Conv) %	ee %
<b>55a</b>	Ph	Me	Me	91 <sup>a</sup> (100 <sup>b</sup> )	98 <sup>a</sup> (98 <sup>b</sup> )
<b>55b</b>	Ph	Me	Et	91 <sup>a</sup>	98 <sup>a</sup>
<b>55c</b>	Ph	Me	Ph	96 <sup>a</sup> (100 <sup>b</sup> )	99 <sup>a</sup> (97 <sup>b</sup> )
<b>55d</b>	Ph	Ph	Me	94 <sup>a</sup>	98 <sup>a</sup>
<b>55e</b>	4-MeOC <sub>6</sub> H <sub>4</sub>	Et	<i>n</i> -Pr	(100 <sup>b</sup> )	(98 <sup>b</sup> )
<b>55f</b>	Et	Me	Ph	89 <sup>a</sup>	87 <sup>a</sup>
<b>55g</b>	H	Bn	Ph	84 <sup>a</sup>	86 <sup>a</sup>
<b>55h</b>	2-MeC <sub>6</sub> H <sub>4</sub>	Me	Me	89 <sup>a</sup>	98 <sup>a</sup>
<b>55i</b>	2-MeOC <sub>6</sub> H <sub>4</sub>	Me	Me	90 <sup>a</sup>	99 <sup>a</sup>
<b>55j</b>	2-ClC <sub>6</sub> H <sub>4</sub>	Me	Me	89 <sup>a</sup>	98 <sup>a</sup>
<b>55k</b>	3-MeOC <sub>6</sub> H <sub>4</sub>	Me	Me	93 <sup>a</sup>	99 <sup>a</sup>
<b>55l</b>	3-ClC <sub>6</sub> H <sub>4</sub>	Me	Me	92 <sup>a</sup>	98 <sup>a</sup>
<b>55m</b>	3-NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	Me	Me	91 <sup>a</sup>	98 <sup>a</sup>
<b>55n</b>	4-MeOC <sub>6</sub> H <sub>4</sub>	Me	Me	97 <sup>a</sup>	98 <sup>a</sup>
<b>55o</b>	4-ClC <sub>6</sub> H <sub>4</sub>	Me	Me	92 <sup>a</sup>	99 <sup>a</sup>
<b>55p</b>	4-NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	Me	Me	88 <sup>a</sup>	99 <sup>a</sup>

<sup>a</sup>Hydrogenations were carried out in toluene [63]

<sup>b</sup>Hydrogenations were carried out in dichloromethane [64]

Some of the more sterically demanding substrates such as **56h** were reduced with a moderate loss of enantioselectivity.

Recently developed ligand SIPHOX **16a–d** proved to be a valuable tool in the asymmetric reduction of  $\alpha,\beta$ -unsaturated acids [66]. In a complete break with what is typically observed with iridium-based N,P ligands, SiPHOX catalysts were observed to reduce the strongly coordinating carboxylic acid triethylamine salt of  $\alpha$  methylcinnamic acid in excellent enantioselectivity and yield (Table 5). Equally surprising is the use of methanol as a solvent, which normally inhibits reductions mediated by iridium N,P catalysts.

Table 5 indicates an extremely efficient catalyst with very low catalyst loadings for such a highly coordinating environment. Functional group tolerance appears to be excellent with groups as reactive as arylbromides being converted in close to perfect yield.

Cesium carbonate was found to be a far more efficient additive for purely alkyl-derived substrates **57r–w**. Very useful enantioselectivities were obtained for all entries with the exception of a tertiary isobutyl-derived alkene **57u**, which was converted with a respectable 90% ee. The yields were slightly lower for purely alkyl-derived substrates, but full conversions were reported by the authors.

**Table 4** Asymmetric hydrogenation of  $\alpha$ -substituted  $\alpha,\beta$ -unsaturated amides with ligand **24**

Substrate	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	ee %
<b>56a</b>	Ph	Me	Ph	96
<b>56b</b>	Ph	Me	<i>n</i> -Bu	84
<b>56c</b>	Ph	Me	<i>i</i> -Bu	93
<b>56d</b>	Ph	Me	Bn	95
<b>56e</b>	Ph	Me	(CH <sub>2</sub> ) <sub>2</sub> OMe	90
<b>56f</b>	Ph	<i>t</i> -Bu	<i>i</i> -Bu	96
<b>56g</b>	Ph	Bn	<i>i</i> -Bu	97
<b>56h</b>	Ph	2-Napht	<i>i</i> -Bu	87
<b>56i</b>	3-ClC <sub>6</sub> H <sub>4</sub>	Et	<i>i</i> -Bu	98
<b>56j</b>	4-MeOC <sub>6</sub> H <sub>4</sub>	Et	<i>i</i> -Bu	96 <sup>a</sup>
<b>56k</b>	Ph	Et	<i>i</i> -Bu	97
<b>56l</b>	Ph	<i>i</i> -Pr	<i>i</i> -Bu	96
<b>56m</b>	Ph	<i>n</i> -Bu	<i>i</i> -Bu	95
<b>56n</b>	Ph	MeO	<i>i</i> -Bu	95
<b>56o</b>	<i>i</i> -Pr	H	Bn	93
<b>56p</b>	<i>n</i> -Pr	H	Bn	84
<b>56q</b>	<i>i</i> -Bu	H	Bn	87
<b>56r</b>	Me	H	Ph	95
<b>56s</b>	Ph	Me	OMe, Me	75 <sup>b</sup>

<sup>a</sup>3 Mol% of catalyst was used<sup>b</sup>Weinreb amide


Scheme 13 depicts an interesting application of this methodology in the synthesis of a crucial intermediate for the blood pressure-lowering drug Aliskiren with the less-encumbered ligand **16c**. In the presence of excess amounts of triethylamine, **58b** was produced in 96% yield and 98% ee. Catalyst loadings were optimized at 0.016 mol% or S/C ratio of 6,000. At S/C ratio of 10,000, a slight loss of enantioselectivity and yield was noticed.

Vinyl phosphine oxides and phosphonates are highly electron-deficient and can undergo conjugate reduction much like their carbonyl counter parts. The group of Andersson has reported an enantioselective reduction of both to yield chiral phosphine oxides and phosphonates (Table 6) [67].

The authors did not report on trisubstituted diphenylphosphine oxides, but apparently, the results are excellent with what is usually a difficult 1,1' unsubstituted pattern about the olefin. The investigation found near perfect selectivity and yields for most of the reported compounds when using **26** as a ligand. However, a very slow reaction was noted for the *E* olefin **59h** while hydrogenation of *E* configured olefin **59j** preceded smoothly, indicating a delicate balance between structure and reactivity. Other anomalous behavior of the substrates **59h–j** is that both *E*- and *Z*-configured olefins produce the same enantiomer.

**Table 5** Asymmetric hydrogenation of cinnamic and tiglic acid derivatives

Substrate	R <sup>1</sup>	R <sup>2</sup>	Yield %	ee %
57a	Ph	Me	99	99.2
57b	2-MeC <sub>6</sub> H <sub>4</sub>	Me	97	99
57c	3-MeC <sub>6</sub> H <sub>4</sub>	Me	98	99
57d	4-MeC <sub>6</sub> H <sub>4</sub>	Me	98	99
57e	2-MeOC <sub>6</sub> H <sub>4</sub>	Me	98	99
57f	3-MeOC <sub>6</sub> H <sub>4</sub>	Me	99	98
57g	4-MeOC <sub>6</sub> H <sub>4</sub>	Me	97	99
57h	2-ClC <sub>6</sub> H <sub>4</sub>	Me	97	96
57i	3-ClC <sub>6</sub> H <sub>4</sub>	Me	98	99
57j	4-ClC <sub>6</sub> H <sub>4</sub>	Me	97	98
57k	3-BrC <sub>6</sub> H <sub>4</sub>	Me	97	99
57l	4-BrC <sub>6</sub> H <sub>4</sub>	Me	97	98
57m	4-CF <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	Me	98	97
57n	2-naphthyl	Me	96	99
57o	furan-2-yl	Me	98	98
57p	Ph	<i>i</i> -Pr	97	99
57q	Ph	Ph	95	94
57r	Me	Me	92	99.1
57s	Et	Me	93	98
57t	<i>n</i> -Pr	Me	89	99
57u	<i>i</i> -Bu	Me	97	90
57v	<i>n</i> -Pr	Et	89	99.4
57w	Me	<i>n</i> -Pr	92	98


**Scheme 13** Asymmetric hydrogenation of an intermediate of aliskiren

## 4 Asymmetric Hydrogenation of 1,1'-Disubstituted Olefins

1,1'-Disubstituted olefins are a challenging substrate class when compared to the more widely investigated trisubstituted olefins. Although they react very quickly, obtaining high enantioselectivity with most ligands has been problematic. Blankenstein and Pfaltz reported the first iridium N,P-based catalyst capable of reducing

**Table 6** Asymmetric reduction of vinyl diphenylphosphine oxides and phosphonates

Substrate	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Conv %	ee %
<b>59a</b>	Ph	Ph	H	99	99
<b>59b</b>	Ph	4-MeOC <sub>6</sub> H <sub>4</sub>	H	99	99
<b>59c</b>	Ph	4-CF <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	H	93	99
<b>59d</b>	Ph	2-MeC <sub>6</sub> H <sub>4</sub>	H	99	99
<b>59e</b>	Ph	C <sub>6</sub> H <sub>11</sub>	H	99	99
<b>59f</b>	Ph	<i>t</i> -Bu	H	98	90
<b>59g</b>	Ph	CH <sub>2</sub> CH <sub>2</sub> OH	H	99	99
<b>59h</b>	OEt	Ph	<i>E</i> -CO <sub>2</sub> Et	10	90
<b>59i</b>	OEt	Ph	<i>Z</i> -CO <sub>2</sub> Et	99	99
<b>59j</b>	OEt	Bn	<i>E</i> -CO <sub>2</sub> Et	99	99
<b>59k</b>	OEt	Ph	H	99	99
<b>59l</b>	OEt	CH <sub>2</sub> CH <sub>2</sub> OAc	<i>E</i> -CO <sub>2</sub> Et	99	99

**Scheme 14** Reduction of 1,1' disubstituted alkenes

1,1'-disubstituted olefins in useful selectivity with SerPHOX **60** (Scheme 14) [38]. The authors discovered a very strong dependence on hydrogen pressure with the highest selectivity obtained at atmospheric pressure of hydrogen. The later developed catalyst Ir(**6b**) was found to give higher enantioselectivities and is now commercially available as the BAR<sub>F</sub> COD complex [68].

A large combined research effort of the three groups of Diéguez, Andersson, and Börner initiated a combinatorial study utilizing the SerPHOX and ThrePHOX backbones in combination with a chiral biarylphosphinite moiety, ligands **18** and

**Table 7** Asymmetric reduction of 1,1'-disubstituted olefins

Substrate	L	Ar	R	Conv %	ee %
<b>61a</b>	<b>19</b>	Ph	Et	100	99
<b>61b</b>	<b>19</b>	4-MeOC <sub>6</sub> H <sub>4</sub>	Et	100	99
<b>61c</b>	<b>19</b>	4-CF <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	Et	100	96
<b>61d</b>	<b>19</b>	Ph	<i>n</i> -Bu	100	94
<b>61e</b>	<b>19</b>	Ph	<i>i</i> -Bu	100	93
<b>61f</b>	<b>19</b>	Ph	neopentyl	100	90
<b>61g</b>	<b>19</b>	Ph	<i>i</i> -Pr	100	97
<b>61h</b>	<b>19</b>	Ph	cyclohexyl	100	97
<b>61i</b>	<b>19</b>	Ph	<i>t</i> -Bu	100	99
<b>61j</b>	<b>18a</b>	2-furyl	<i>n</i> -Bu	100	99
<b>61k</b>	<b>19</b>	2-thienyl	<i>n</i> -Bu	100	96
<b>61l</b>	<b>19</b>	2-pyridyl	Et	100	99
<b>61m</b>	<b>19</b>	2-pyridyl	<i>t</i> -Bu	100	99
<b>61n</b>	<b>18b</b>	Ph	2-naphthyl	100	99 <sup>a</sup>
<b>61o</b>	<b>18b</b>	Ph	2-MeC <sub>6</sub> H <sub>4</sub>	99	99 <sup>a</sup>
<b>61p</b>	<b>19</b>	4-CF <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	4-MeOC <sub>6</sub> H <sub>4</sub>	100	63 <sup>a</sup>
<b>61q</b>	<b>19</b>	Ph	CH <sub>2</sub> OH	100	95 <sup>a</sup>
<b>61r</b>	<b>19</b>	Ph	CH <sub>2</sub> OAc	100	91 <sup>a</sup>
<b>61s</b>	<b>19</b>	Ph	CH <sub>2</sub> TMS	100	96 <sup>a</sup>
<b>61t</b>	<b>19</b>	4-MeOC <sub>6</sub> H <sub>4</sub>	CF <sub>3</sub>	100	75 <sup>a</sup>

<sup>a</sup>Reaction was run at 50 bar hydrogen

**19** [69, 94]. A library of 96 possible phosphite oxazoline ligands was synthesized with systematic changes to the backbone and with special attention to the biaryl phosphite group.

Ligand **19** was found to be highly selective for the hydrogenation of 1-aryl, 1'-alkyl alkenes, and ligand **18b** was found to be selective for 1-aryl, 1'-hindered aryl alkenes (Table 7).

Ligand **19** performs excellently with the wide variety of 1,1'-disubstituted olefins reported. Substrates **61a–m** are efficiently reduced at 1 bar of hydrogen in high enantioselectivity with very little dependence on the bulk of the alkyl substituents. Strongly coordinating olefins such as **61l** and **61m** typically perform poorly in iridium-catalyzed hydrogenations, but reduction with **19** clearly breaks this rule and the substrates are reduced in excellent selectivity and yield.

The more sterically demanding **61n–o** were easily reduced by Ir(**18b**), perhaps due to a more open and accommodating catalyst structure. Other changes to note are the higher pressure (50 bar) required to reduce the chelating substrates **61q–r**, the hindered **61s**, and the electron poor **61t**. The diarylmethylene substrate **61p** reveals that enantioface discrimination is also possible based on electronic rather than steric effects. Both aryl groups have very similar steric demands but distinguish themselves by their  $\pi$  donor and  $\pi$  acceptor properties. The remarkable ee of

63% may be explained by the different interaction of the iridium center with the electron-poor and the electron-rich aryl groups and/or alignment of the catalyst dipole with the substrate dipole created by electron donor and acceptor groups.

## 5 Asymmetric Hydrogenation of Tetrasubstituted Olefins

Tetrasubstituted olefins remain a challenging class of substrate. Buchwald and coworkers have shown that chiral zirconocene complexes can catalyze the hydrogenation of tetrasubstituted olefins with high enantioselectivity [7, 8]. However, high catalyst loadings, long reaction times, and high pressures are a disadvantage of this system.

Schrems et al. identified a number of P,N ligands, which have given encouraging results for this class of substrate [70]. Surprisingly, the structurally simple readily accessible phosphinooxazoline **12a**, originally reported by Sprinz and Helmchen [71], and subsequent analogs **12b–e** proved to be the most efficient ligands for several substrates (Table 8).

For a few select cases, the cyclic alkene **62a–c** with simple primary alkyls as substituents were readily hydrogenated with SimplePHOX ligand **7a** and **7c**. Importantly, no epimerization at the benzylic position was observed, and hydrogenation gave entirely *cis* product with most substrates. Aromatization of the dihydronaphthalene substrates **63a–b** was a frequent side reaction, even at high pressures.

For some substrates, a dependence of enantioselectivity on pressure was observed with low to moderate pressures being optimal for many combinations. Ligand **12c** in combination with substrate **62a** gives an enantioselectivity of 92% at 50 bar of pressure and 97% at 1 bar, similar to the trend found in the early studies of the 1,1'-disubstituted substrates.

Yet even more important is the observation that ligands with identical configurations gave products of opposite configuration solely on the basis of the substituent on the stereogenic center. For instance, ligand **S-12a** gives the (–) reduction product of **63a** in 65% ee at 50 bar, and changing the ligand to **S-12f** under identical reaction conditions gives the (+) product in 39% ee. Clearly, the critical interactions with these catalysts and substrates are highly subject to the extremely crowded environment about the metal center and very small changes to either the substrate or ligand can lead to large changes in overall selectivity and yield.

Mastery of tetrasubstituted olefins greatly expands the tools available for the synthesis of chiral carbon skeletons and tetrasubstituted olefins have the added value of possibly creating two stereogenic centers simultaneously.

## 6 Asymmetric Hydrogenation of Trisubstituted Olefins with Heteroatoms

Asymmetric hydrogenation of a C=C bond with a heteroatom substituent leads to highly valued chiral centers which can often be elaborated into more complex functionalized molecules. Most asymmetric hydrogenation routes to these chiral centers have



**Table 8** Asymmetric reduction of tetrasubstituted olefins

Substrate	R	L	P (bar)	t (h)	ee %	Conv %	
<b>61a</b>	– MeO	<b>12c</b>	1	4	97	99	
<b>61b</b>	– F	<b>12d</b>	5	4	89	99	
Substrate	R <sup>1</sup>	R <sup>2</sup>	L	P (bar)	t (h)	ee (%)	Conv (%)
<b>62a</b>	Me	Me	<b>12a</b>	5	4	94	99
			<b>7c</b>	5	4	95	99
<b>62b</b>	Et	Me	<b>12a</b>	50	4	93	99
			<b>7a</b>	50	4	94	98
<b>62c</b>	<i>n</i> -Bu	Me	<b>12a</b>	50	4	94	99
			<b>7a</b>	50	4	95	97
<b>62d</b>	Ph	Me	<b>12a</b>	5	4	95	99
<b>62e</b>	Me	Ph	<b>12a</b>	50	4	88	20
<b>62f</b>	Et	Ph	<b>12a</b>	50	4	93	23
Substrate	R	L	P (bar)	t (h)	ee %	Conv %	
<b>63a</b>	Me	Me	<b>12a</b>	50	4	73	99
	Me	Me	<b>6a</b>	5	4	77	46
<b>63a</b>	Ph	Me	<b>12b</b>	5	4	91	32
Substrate	L	P (bar)	mol%	ee %	Conv %		
<b>64</b>	<b>5a</b>	50	2.0	94	99		
		50	1.0	93	99		
		50	0.5	93	99		
		50	0.1	90	99		
	<b>12e</b>	5	2.0	96	99		

relied on ruthenium or rhodium diphosphine complexes [1–4, 11]. Iridium N,P and C,N complexes are the newest addition to catalysts capable of asymmetric hydrogenation of heteroatom-substituted olefins, and in spite of the typical sensitivity of the complexes toward Lewis bases, many successful hydrogenations of such olefins have been reported in the recent literature. Indoles, enamines, enol esters, enol ethers, chromenes, furans, quinolines, and pyridines have all been successfully reduced with useful enantioselectivities using chiral iridium complexes and each shall be touched upon in the following section.

## 6.1 Enol Esters and Ethers

Zhu and Burgess have reported an asymmetric conjugate reduction of 1,3-enol ether esters (Table 9) and 1,3-enol ether alcohols (Table 10) [72]. Initial reaction conditions reached full conversion of *E*-1-methoxy-1-phenylethene using ligand **9** albeit with a very low enantioselectivity of 29%.

Asymmetric hydrogenation of **65a–h** with ligand **9** indicates a difficult reaction. Enantioselectivities are moderate to good with the best selectivity for the amide **65g**. Strong steric effects were observed with increasing the size of the ester giving an increase in ee. The steric crowding at the alkoxy bearing C atom was also observed to have a large effect on the enantioselectivity. Substituents larger than methoxy groups greatly eroded the selectivity as observed for the ethoxy ether **65e**. Secondary alkyl groups were completely deleterious to hydrogenation activity with an abysmal conversion of 15% for substrate **65f**.

Asymmetric hydrogenation of vinyl ether alcohols proceeded in better selectivity than the ester counterparts, but acid sensitivity was observed for **66a–d**, and a stoichiometric equivalent of potassium carbonate relative to the substrate was

**Table 9** Asymmetric hydrogenation of vinyl ether esters

Substrate	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Conv %	ee %
<b>65a</b>	Me	OMe	CO <sub>2</sub> H	99	63
<b>65b</b>	Me	OMe	CO <sub>2</sub> Me	85	60
<b>65c</b>	Me	OMe	CO <sub>2</sub> Et	99	78
<b>65d</b>	Me	OMe	CO <sub>2</sub> - <i>t</i> -Bu	99	88
<b>65e</b>	Me	OEt	CO <sub>2</sub> Et	99	66
<b>65f</b>	OMe	<i>i</i> -Pr	CO <sub>2</sub> Me	15	nd
<b>65g</b>	Me	OMe	CO <sub>2</sub> NEtOMe	90	90
<b>65h</b>	Me	OMe	COPh	5	nd

**Table 10** Asymmetric hydrogenation of vinyl ether alcohols

Substrate	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	ee %
<b>66a</b>	Me	OMe	H	96
<b>66b</b>	Me	OEt	H	98
<b>66c</b>	<i>n</i> -Bu	OMe	H	93
<b>66d</b>	<i>i</i> -Pr	OMe	H	91
<b>66e<sup>a</sup></b>	Me	OMe	TBDPS	89
<b>66f<sup>a</sup></b>	Me	OMe	Ac	92

<sup>a</sup>K<sub>2</sub>CO<sub>3</sub> was not required for this case

**Table 11** Ligand and substrate optimization studies

R	L 13a	L 13b	L 11a	L 10b
TMS	<sup>a</sup>	<sup>a</sup>	<sup>a</sup>	<sup>a</sup>
CH <sub>3</sub>	<sup>a</sup>	<sup>a</sup>	<sup>a</sup>	<sup>a</sup>
Ac	<sup>a</sup>	Full conv; rac	<sup>a</sup>	No conv
PO(OEt) <sub>2</sub>	Complex; mixture <sup>b</sup>	Full conv; 65% ee	<sup>a</sup>	No conv
POPPh <sub>2</sub>	30% conv; 82% ee	Full conv; 95% ee	41% conv; 63% ee	No conv


<sup>a</sup>A complex mixture was obtained

<sup>b</sup>Ethylbenzene was the major product

required to neutralize any acidic contaminants. Larger groups on either side of the olefin eroded enantioselectivity but not to the same degree as with substrates **65a–f**.

Cheruku et al. have investigated a series of enol esters [73, 95]. A preliminary experiment to determine the optimal protecting group with a small group of catalysts identified diphenylphosphinates as the optimal esters (Table 11).

Ligand **13b** in combination with diphenylphosphinate as the ester group gave the chiral phenylethyl diphenylphosphinate **67b** with full conversion and high enantioselectivity of 95%. Although it may seem that the diphenylphosphinate ester group is not ideal in terms of atom economy, it should be noted that it is an excellent leaving group and can be readily displaced with a number of good

**Table 12** Asymmetric hydrogenation of enol phosphinates

Substrate	R <sup>1</sup>	R <sup>2</sup>	Conv %	ee %
<b>68a</b>	Ph	H	99	95
<b>68b</b>	4-MeC <sub>6</sub> H <sub>4</sub>	H	97	96
<b>68c</b>	4- <i>t</i> -BuC <sub>6</sub> H <sub>4</sub>	H	93	94
<b>68d</b>	4-MeOC <sub>6</sub> H <sub>4</sub>	H	48 <sup>a</sup>	98
<b>68e</b>	4-BrC <sub>6</sub> H <sub>4</sub>	H	99	99
<b>68f</b>	4-CF <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	H	99	99
<b>68g</b>	4-NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	H	99	92
<b>68h</b>	β-naphthyl	H	99	85
<b>68i</b>	cyclohexyl	H	99	92
<b>68j</b>	<i>t</i> -Bu	H	99	99
<b>68k</b>	Ph	Me	99	96
<b>68l</b>	Ph	Et	90	92
<b>68m</b>	Ph	<i>i</i> -Pr	89	90
<b>68n</b>	Ph	CO <sub>2</sub> Et	99	99
<b>68o</b>	4-MeC <sub>6</sub> H <sub>4</sub>	CO <sub>2</sub> Et	99	99
<b>68p</b>	4-CF <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	CO <sub>2</sub> Et	99	99
<b>68q</b>	4-BrC <sub>6</sub> H <sub>4</sub>	CO <sub>2</sub> Et	99	98
<b>68r</b>	Me	CO <sub>2</sub> Et	99	99
<b>68s</b>	Et	CO <sub>2</sub> Et	99	99
<b>68t</b>	<i>i</i> -Pr	CO <sub>2</sub> Et	99	99
<b>68u</b>	ClCH <sub>2</sub>	CO <sub>2</sub> Et	95 <sup>b</sup>	93 <sup>b</sup>
<b>68v</b>	<i>t</i> -Bu	CO <sub>2</sub> Et	98	93
<b>68w</b>	<i>t</i> -Bu	Et	99	90
<b>68x</b>	<i>i</i> -Pr	Me	99	91

<sup>a</sup>Reaction was run with 2 mol% catalyst and 10 mg poly(vinylpyridine) resin

<sup>b</sup>Run at 2 mol% catalyst

nucleophiles. A study of both 1,1'-disubstituted vinyl phosphinates and trisubstituted vinyl phosphinates was conducted and the results are summarized in Table 12.

Asymmetric hydrogenation of both 1,1'-disubstituted vinyl phosphinates and the trisubstituted phosphinates with ligand **13b** was highly successful. Enantioselectivities were very high for many useful combinations including bulky alkyl groups on the stereogenic center. Substrate **68d** was very acid-sensitive, and an acid scavenger with increased amounts of catalyst was required to obtain reasonable results. Allylic chloride **68u** also required higher catalyst loading but was reduced efficiently and in high enantioselectivity. In many cases, the enantioselectivity was higher than what can be obtained from reduction of the corresponding ketones with ruthenium-based catalysts [95].

## 6.2 Asymmetric Hydrogenation of Furans and Chromenes

Chiral-reduced heterocycles containing only oxygen are common throughout organic synthesis and are highly sought after compounds. Asymmetric hydrogenation offers an atom-economical approach to these compounds in a single, clean step. However, hydrogenation of prochiral furans and dehydropyrans has proven difficult, mostly attributed to the increased stability of aromatic compounds and the electron-rich nature of these substrates [74]. For a long time, no suitable chiral catalysts were available until iridium N,P catalyst provided a solution for these problematic hydrogenation reactions.

ThrePHOX **6b** proved to be a valuable tool in the reduction of chromenes **71a–k**, all of which were reduced in greater than 90% enantioselectivity. The very electron-rich thiochromene **71h** required higher temperature, and lower conversion was observed, but the enantioselectivity remained in the useful range (Table 13) [96].


A series of furans and benzofurans were reduced in excellent enantioselectivity and conversion using iridium complexes derived from pyridine–phosphinite ligands. Tertiary butyl-derived phosphinites were the preferred donor groups with ligand **8a** giving the highest enantioselectivity most consistently. Higher temperatures and longer reaction times were required to obtain good conversion for all cases; most notably, **69e** reacted very sluggishly at 40°C (Fig. 8).

These results mark the very first successful catalytic reduction of furans in high enantioselectivity and greatly expand the methods available for the synthesis of enantiopure heterocycles.

**Table 13** Asymmetric hydrogenation of substituted 4H-chromenes

Substrate	R <sup>1</sup>	R <sup>2</sup>	X	Conv %	ee %
<b>71a</b>	Me	H	O	93	95
<b>71b</b>	4-MeC <sub>6</sub> H <sub>4</sub>	H	O	99	98
<b>71c</b>	2-FC <sub>6</sub> H <sub>4</sub>	H	O	99	99
<b>71d</b>	4-BrC <sub>6</sub> H <sub>4</sub>	H	O	95	91
<b>71e</b>	4-ClC <sub>6</sub> H <sub>4</sub>	6-Cl	O	90	97
<b>71f</b>	Ph	7-MeO	O	99	99
<b>71g</b>	2-furyl	H	O	99	97
<b>71h</b>	Ph	H	S	73 <sup>a</sup>	91 <sup>a</sup>
<b>71i</b>	Ph	H	O	99	99
<b>71j</b>	cyclohexyl	H	O	99	95
<b>71k</b>	Ph	7-(CH=CH) <sub>2</sub> -8	O	99	96

<sup>a</sup>Reaction was conducted at 40°C



**Fig. 8** Asymmetric hydrogenation results for substituted furans

### 6.3 Asymmetric Hydrogenation of Enamines and Indoles

Much like the enol systems discussed in Sect. 6.1, enamines are predictably difficult substrates for most iridium asymmetric hydrogenation catalysts. Both substrate and product contain basic functionalities which may act as inhibitors to the catalyst. Extended aromatic enamines such as indoles may be even more difficult substrates for asymmetric hydrogenation with an additional energetic barrier to overcome. Initial reports by Andersson indicated a very difficult reaction indeed (Table 14) [75]. Higher enantioselectivities were later reported by Baeza and Pfaltz (Table 14) [76].

Moderate enantioselectivities were obtained with ligand **13b** at room temperature. Simply changing from a diethyl amine moiety **72a** to a pyrrolidine **72g** caused a precipitous drop in enantioselectivity for both ligands **13b** and ThrePHOX **6b**. An electronic effect was observed for ligand **13b** with a drop from good selectivity with enamine **72b** with a methyl group in the para position of the aromatic ring to poor selectivity with the stronger  $\pi$  donating paramethoxy group with **72c**.

Steric interaction was also clearly an issue for all of the systems investigated. Most notable is the large difference in the reactivity between ThrePHOX **6b**- and PHOX **5c**-based catalysts with respect to the substituents bound to the nitrogen atom. ThrePHOX consistently performed best with *N*-methyl-*N*-benzyl enamines while PHOX preferred a *N*-methyl-*N*-phenyl derivative. Lowering the temperature had a favorable outcome on enantioselectivity but slowed the reaction and caused drops in conversion for **5c** and **6b**.

A variety of ligands were used to hydrogenate **73a**, **74a–b**, **75a–b**, and **76a** with moderate results. Steric interactions were magnified and that may explain why nearly all of the substrates in Scheme 15 give less than optimal results. *N*-methyl-*N*-phenyl-amine derived enamines **74a** and **75a** were resistant to hydrogenation where the analogous *N*-methyl-*N*-benzyl-amine-based enamines were completely converted with SimplePHOX ligand **77** with moderate enantioselectivity.

**Table 14** Asymmetric reduction of enamines

Substrate	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Conv %	ee %
<b>72a</b>	Et	Et	Ph	99 <sup>a</sup> (99 <sup>b</sup> )	84 <sup>a</sup> (54 <sup>b</sup> )
<b>72b</b>	Et	Et	4-MeC <sub>6</sub> H <sub>4</sub>	99 <sup>a</sup>	87 <sup>a</sup>
<b>72c</b>	Et	Et	4-MeOC <sub>6</sub> H <sub>4</sub>	99 <sup>a</sup>	64 <sup>a</sup>
<b>72d</b>	Et	Et	4-CF <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	99 <sup>a</sup>	77 <sup>a</sup>
<b>72e</b>	Et	Et	2-naphthyl	99 <sup>a</sup>	64 <sup>a</sup>
<b>72f</b>	Me	Ph	Ph	99 <sup>a</sup> (98 <sup>c</sup> )	79 <sup>a</sup> (91 <sup>c</sup> )
<b>72g</b>	<i>N,N'</i> -(CH <sub>2</sub> ) <sub>4</sub>		Ph	66 <sup>a</sup> (99 <sup>b</sup> )	33 <sup>a</sup> (8 <sup>b</sup> )
<b>72h</b>	<i>N,N'</i> -(CH <sub>2</sub> -CH <sub>2</sub> ) <sub>2</sub> O		Ph	75 <sup>a</sup>	30 <sup>a</sup>
<b>72i</b>	Me	4-MeOC <sub>6</sub> H <sub>4</sub>	Ph	99 <sup>c</sup>	90 <sup>c</sup>
<b>72j</b>	Me	4-ClC <sub>6</sub> H <sub>4</sub>	Ph	99 <sup>c</sup>	90.5 <sup>c</sup>
<b>72k</b>	Me	Bn	Ph	99 <sup>b,d</sup>	92.5 <sup>b,d</sup>
<b>72l</b>	Me	Bn	4-MeOC <sub>6</sub> H <sub>4</sub>	93 <sup>b</sup>	74 <sup>b</sup>
<b>72m</b>	Me	Bn	4-ClC <sub>6</sub> H <sub>4</sub>	93 <sup>b</sup>	74 <sup>b</sup>
<b>72n</b>	Me	Bn	2-furan	93 <sup>b,e</sup>	74 <sup>b,e</sup>
<b>72o</b>	Me	Bn	t-Bu	53 <sup>c</sup>	21 <sup>c</sup>

<sup>a</sup>Run with 0.5 mol% ligand **13b** at room temperature and 50 bar H<sub>2</sub> [75]

<sup>b</sup>Run with 0.5 mol% ligand **6b** at -20°C [76]

<sup>c</sup>Run with 0.5 mol% ligand **5c** at -20°C [76]


<sup>d</sup>MTBE was used as solvent

<sup>e</sup>Run at 0°C


It should also be mentioned that a very active and selective catalyst in the form of a monodentate phosphoramidite in combination with iridium was very successful in the reduction of cyclic enamines, but the discussion of this work is beyond the scope of this chapter (Scheme 16) [77]. Enantioselectivities were reported as excellent with atmospheric pressure sufficing for full conversion in most cases.

Baeza and Pfaltz also investigated indoles which reacted very sluggishly but excellent enantioselectivity could be obtained (Table 15) [78].

Hydrogenation of protected 2-indoles required higher catalyst loading and higher temperatures than used for enamines, but many substrates were hydrogenated in high enantioselectivity and yield. Both electron-donating groups and electron-withdrawing groups interfered with catalysis but the reaction with indole **79b** could be pushed to completion by changing the solvent to chlorobenzene and increasing the temperature to 110°C with very little loss in enantioselectivity. Nitroindole **79d** was completely unreactive and no hydrogenation products were observed. The boc-protecting group was exchanged for acyl or tosyl groups in a few cases and with a beneficial effect to conversion for some indoles. Indole **79i** was extremely difficult to reduce but useful levels of product and enantioselectivity



Scheme 15 Asymmetric hydrogenation of cyclic and acyclic enamines

Scheme 16 Asymmetric hydrogenation of enamines with Ir(**78**)

could be obtained with ligand **8a**. Indoles with substitution in the 3 position were very difficult to hydrogenate and reactions with esters in this position completely failed. However, simple methyl derivative **79k** was hydrogenated in excellent ee and yield.



**Table 15** Asymmetric hydrogenation of indoles


Substrate	mol% (L)	<i>T</i> (°C)	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	Conv % <sup>a</sup>	ee % <sup>a</sup>
<b>79a</b>	4 ( <b>8a</b> )	60	boc	Me	H	H	89	99
<b>79b</b>	4 ( <b>8a</b> )	60	boc	Me	H	MeO	68	99
		110 <sup>b</sup>					85 <sup>b</sup>	98.5 <sup>b</sup>
<b>79c</b>	4 ( <b>8a</b> )	60	boc	Me	H	F	64	98.5
<b>79d</b>	4 ( <b>8a</b> )	60	boc	Me	H	NO <sub>2</sub>	0	nd
<b>79e</b>	1 ( <b>6b</b> )	25	boc	Ph	H	H	94 <sup>c</sup>	88 <sup>c</sup>
<b>79f</b>	1 ( <b>6b</b> )	25	boc	CO <sub>2</sub> Et	H	H	66 <sup>c</sup>	92 <sup>c</sup>
							88 <sup>d</sup>	93 <sup>d</sup>
<b>79g</b>	2 ( <b>8a</b> )	60	Ac	CO <sub>2</sub> Et	H	H	97	99
<b>79h</b>	2 ( <b>80</b> )	25	Tosyl	Me	H	H	94	99
<b>79i</b>	4 ( <b>8a</b> )	110 <sup>b</sup>	Tosyl	Ph	H	H	70 <sup>b</sup>	98 <sup>b</sup>
<b>79j</b>	4 ( <b>8a</b> )	60	Tosyl	Me	H	H	95	99
<b>79k</b>	2.5 ( <b>8a</b> )	60	Tosyl	H	Me	H	97	98

<sup>a</sup>Reaction was run at 100 bar unless otherwise noted<sup>b</sup>Chlorobenzene as solvent, 110°C<sup>c</sup>Reaction was run at 75 bar<sup>d</sup>2 mol% of catalyst, DCM, 25°C

## 6.4 Asymmetric Hydrogenation of Quinolines and Pyridines

Asymmetric hydrogenation of nitrogen heterocycles is a relatively unexplored and open field. Only a handful of iridium N,P-based systems have been used successfully to hydrogenate substituted quinolines and the more stable pyridines. Lu et al. were the first to report a successful asymmetric hydrogenation of two substituted quinolines with ligand **24** (Table 16) [79]. A few reports using diphosphines and monophos variations followed until Lu and Bolm published a second iridium N,P-based asymmetric hydrogenation of similar quinolines with ligand **23a** (Table 16) [80].

Ligand **24** was surprisingly effective in the presence of 5 mol% iodine without the use of BAR<sub>F</sub> as a counterion. Reduction of sterically demanding substrates was a problem for both systems and aromatic substitution on the 2 position of the quinoline abolished the enantioselectivity and poor conversion was obtained. Electron-donating and -withdrawing groups also damaged the enantioselectivity although less for Zhou's system. Nevertheless, useful enantioselectivity and yield could be obtained for simple alkyl derivatives.

**Table 16** Asymmetric hydrogenation of quinolines


Substrate	R <sup>1</sup>	R <sup>2</sup>	Yield % <sup>a</sup> Conv % <sup>b</sup>	ee % <sup>a</sup> ee % <sup>b</sup>
<b>81a</b>	Me	H	95 <sup>a</sup> 95 <sup>b</sup>	90 <sup>a</sup> 87 <sup>b</sup>
<b>81b</b>	Et	H	95 <sup>a</sup> 62 <sup>b</sup>	91 <sup>a</sup> 77 <sup>b</sup>
<b>81c</b>	<i>n</i> -Pr	H	62 <sup>b</sup>	80 <sup>b</sup>
<b>81d</b>	<i>i</i> -Bu	H	53 <sup>b</sup>	75 <sup>b</sup>
<b>81e</b>	<i>n</i> -pentyl	H	94 <sup>a</sup> 90 <sup>b</sup>	92 <sup>a</sup> 65 <sup>b</sup>
<b>81f</b>	Me	Me	93 <sup>a</sup> 95 <sup>b</sup>	92 <sup>a</sup> 75 <sup>b</sup>
<b>81g</b>	Me	F	86 <sup>a</sup> 43 <sup>b</sup>	89 <sup>a</sup> 64 <sup>b</sup>
<b>81h</b>	Me	MeO	95 <sup>b</sup>	78 <sup>b</sup>
<b>81i</b>	Ph	H	45 <sup>a</sup>	3 <sup>a</sup>
<b>81j</b>	(CH <sub>2</sub> ) <sub>2</sub> Ph	H	92 <sup>a</sup>	72 <sup>a</sup>

<sup>a</sup>Reaction was conducted at 40 bar H<sub>2</sub>, 1 mol% [(**24**)Ir(COD)]Cl and 5 mol% I<sub>2</sub> [79]


<sup>b</sup>Reaction was conducted at 60 bar H<sub>2</sub>, 1 mol% [(**23a**)Ir(COD)]BAR<sub>F</sub> [80]

A single paper from Charette and coworkers details the catalytic asymmetric hydrogenation of *N*-iminopyridinium ylides to substituted piperdines using PHOX ligand **82** in combination with iodine (Table 17) [81].

Moderate to good enantioselectivities were obtained for nearly all examples, but the products from **83a–c** could be recrystallized to higher enantiomeric purity. Addition of iodine was critical for catalysis as was the use of a ligand with electron-poor *para*-fluorophenyl groups on the phosphorous atom. Substitution at the 3 position of the pyridine ring was described as being difficult for both the quinolines and pyridine systems. The resulting hydrazine derivatives could be easily converted to piperdines by reduction with Raney nickel or under Birch conditions.

## 7 Asymmetric Hydrogenation of Imines

Given the importance of chiral amines to synthetic chemistry as well as other fields asymmetric hydrogenation of imines has attracted wide interest but limited success compared to C=C and C=O bond reduction. The first asymmetric hydrogenation of imines was carried out in the seventies with ruthenium- and rhodium-based catalysts, followed later by titanium and zirconium systems [82]. Buchwald found that

**Table 17** Asymmetric hydrogenation of *N*-iminopyridinium ylides

Substrate	R	Yield %	ee %
<b>83a</b>	2-Me	90 (84 <sup>a</sup> )	90 (97 <sup>a</sup> )
<b>83b</b>	2-Et	96 (78 <sup>a</sup> )	83 (94 <sup>a</sup> )
<b>83c</b>	2- <i>n</i> Pr	98 (75 <sup>a</sup> )	84 (95 <sup>a</sup> )
<b>83d</b>	2-Bn	97	58
<b>83e</b>	2-CH <sub>2</sub> OBn	85	76
<b>83f</b>	2-(CH <sub>2</sub> ) <sub>3</sub> OBn	88	88
<b>83g</b>	2,3-dimethyl	91 (>95:5 <i>cis:trans</i> )	54
<b>83h</b>	2,5-dimethyl	92 (57:43 <i>cis:trans</i> )	86/84


<sup>a</sup>Numbers in parentheses are after a single recrystallization

Britzinger-type ansa titanocenes were highly selective for the reduction of cyclic imines, albeit at long reaction times, high catalyst loading, and with high pressures [83]. Crabtree found that pyridine phosphine complexes of iridium were highly active in the reduction of imines [10]. The asymmetric synthesis of Metolachlor, an important agrochemical used in the protection of maize, is the first example of an industrially useful enantioselective reduction of imines catalyzed by an iridium complex with a Josiphos type ligand [84]. The first use of iridium N,P complexes in the asymmetric hydrogenation of imines was reported by Pfaltz and coworkers [85].

Many of the systems studied required high catalyst loadings, very high pressures, elevated temperatures, and long reaction times. Several problems can be encountered with imines such as hydrolysis to starting materials, dimer and trimer formation, and the presence of enamine and syn/anti isomers [82]. Additionally, with the exception of *N*-aryl imines, the resulting amines are far more basic than the imine substrates and may cause a larger degree of product inhibition of the catalyst. Many recent advances have been made with iridium systems other than N,P ligands such as diphosphines and monodentate phosphoramidites [86, 87].

Initial studies with PHOX indicated a lack of an anion effect, replacement of PF<sub>6</sub><sup>-</sup> by SbF<sub>6</sub><sup>-</sup>, or BF<sub>4</sub><sup>-</sup> as counter ion had no apparent effect on the overall reaction. However, nearly all reported catalysts within the last decade utilize BAR<sub>F</sub> as a counterion, perhaps for the higher catalyst stability, easier handling, and general reactivity toward a broader range of substrates.

Andersson reported the use of ligand **13a** in the asymmetric hydrogenation of substituted acetophenone-based *N*-aryl imines (Table 18) [88, 89]. New ligands **84** and **85** were reported by the groups of Bolm and Knochel, respectively, for


**Table 18** Asymmetric hydrogenation of acetophenone based *N*-aryl ketimines

Imine	R <sup>1</sup>	R <sup>2</sup>	Ligand ee%							
			5a <sup>a</sup>	13a <sup>b</sup>	14d <sup>c</sup>	15a <sup>d</sup>	16a <sup>e</sup>	77 <sup>f</sup>	84 <sup>g</sup>	85 <sup>h</sup>
86a	H	H	90	90	94	91	93	96	–	84
86b	H	4-MeO	96	89	94	90	–	95	96	85
86c	H	Ar <sup>i</sup>	–	–	–	–	–	–	–	94
87a	2-Me	H	–	83	–	–	–	–	–	–
87b	2-Me	4-MeO	–	–	–	–	–	–	94	–
87c	2-Me	Ar <sup>i</sup>	–	–	–	–	–	–	–	94
88a	3-Me	4-MeO	–	–	–	–	–	–	93	–
88b	3-Me	Ar <sup>i</sup>	–	–	–	–	–	–	–	93
91a	4-Me	H	–	–	–	88	94	–	–	–
91b	4-Me	4-MeO	–	–	–	–	–	–	96	–
92	4-AcO	Ar <sup>i</sup>	–	–	–	–	–	–	–	94
93	4-MeO	H	–	86	90	–	94	94	–	–
94a	4-MeO	4-MeO	–	86	–	–	–	–	94	–
94b	3-MeO	4-MeO	–	–	–	–	–	–	96	–
94c	2-MeO	4-MeO	–	–	–	–	–	–	90	–
95a	3-Cl	H	–	–	–	93	93	–	–	–
95b	3-Br	H	–	–	–	93	92	–	–	–
96a	4-Cl	H	96	–	95	93	90	96	–	–
96b	4-Cl	4-MeO	–	89	–	91	–	–	95	–
96c	4-Cl	Ar <sup>i</sup>	–	–	–	–	–	–	–	92
97a	4-Br	H	–	–	–	91	91	–	–	–
97b	3-Br	H	–	–	–	93	92	–	–	–
98a	4-CF <sub>3</sub>	H	–	–	–	92	–	–	–	–
98b	4-CF <sub>3</sub>	Ar <sup>i</sup>	–	–	–	–	–	–	–	89
99a	H	4-Cl	89	–	92	–	97	94	–	–
99b	H	4-Br	–	–	–	89	96	–	–	–
100	3,4-Xy	H	–	–	–	95	94	–	–	–

Condition <sup>a</sup>0.5 mol% cat, DCM, 5 bar H<sub>2</sub>, 0°CCondition <sup>b</sup>0.5 mol% cat, DCM, 20 bar H<sub>2</sub>Condition <sup>c</sup>0.5 mol% cat, DCM, 5 bar H<sub>2</sub>, –20°CCondition <sup>d</sup>1.0 mol% cat, DCE, 1 atm H<sub>2</sub>, 10°CCondition <sup>e</sup>1.0 mol% cat, MTBE, 1 atm H<sub>2</sub>, 10°C, 4 Å MSCondition <sup>f</sup>0.5 mol% cat, DCM, 5 bar H<sub>2</sub>, –20°CCondition <sup>g</sup>0.5 mol% cat, 2 mol% I<sub>2</sub>, toluene, 20 bar H<sub>2</sub>Condition <sup>h</sup>1.0 mol% cat, 4:1 toluene: methanol, 10 bar H<sub>2</sub>Condition <sup>i</sup>Ar = 4-MeO-3,5-Me<sub>2</sub>-C<sub>6</sub>H<sub>2</sub>

asymmetric hydrogenation of imines with improved enantioselectivities and yields (Fig. 9) [90, 91]. Two similar chiral spirocycle-based ligands SpinPHOX **15** and SiPHOX **16** proved highly effective in the asymmetric hydrogenation of these difficult substrates [92, 93]. Baeza and Pfaltz reported three structurally similar

**Fig. 9** Ligands for asymmetric hydrogenation of imines



catalysts based on PHOX in an expanded look at the earlier work and found excellent selectivity and activity at low catalyst loadings [97].

Essentially, all studies have used acetophenone-based *N*-aryl ketimines as substrates or derivatives thereof. Some of the *N*-aryl groups are easily deprotected to yield branched  $\alpha$ -aryl primary amines, which can be further functionalized.

Several authors have conducted investigations of the *N*-aryl moiety to optimize interactions for each individual catalyst. All of the catalysts in Table 18 function well at 1 mol% catalyst loading and with ambient to moderate pressures of hydrogen. Noncoordinating solvents were preferred with the exception of Knochel's ligand **85**, which gave the highest enantioselectivity in a mixture of 25% methanol in toluene.

Although all of the substrates tested in Tables 18 and 19 have a very high degree of similarity, some useful trends can be identified for each individual catalyst. Different conditions were used for different catalyst systems, possibly reflecting different mechanisms for each individual system. For instance, addition of iodine was found to give inferior results with PHOX and related ligands, yet Bolm's ligand **84** requires iodine for catalysis, a result more in agreement with iridium diphosphine complexes [82]. Other interesting deviations are the optimal conditions reported by Knochel with ligand **85**, which calls for 25% methanol in toluene as a solvent. This result stands in stark contrast to Zhou's ligand **16a**, which is inhibited by methanol in imine hydrogenations.

PHOX **5a**, SimplePHOX **77**, and NeoPHOX **14d** are very closely related in bite angle and overall geometry however these catalysts differ in the electronic properties on the phosphine moiety. All of these ligands gave excellent enantioselectivity in close agreement with each other, reflecting the importance of geometry and bite angle over electronic properties. The authors also reported a strong temperature effect on selectivity with lower temperature giving higher enantioselectivity. Catalysts loadings as low as 0.1 mol% were effective for several substrate and catalyst combinations but with an increased pressure requirement to reach useful conversions.

In comparison, ligand **13a** generally gave moderate to good enantioselectivities over a limited range of substrates, and larger substituents were not tolerated well with imine **87a** reaching only 52% conversion albeit at 0.5 mol% catalyst loading. This system appeared to be the most sensitive to the size of the substrate, changing a methyl group such as in imine **86a** with an ee of 90% for an ethyl group as in **101a** decreased the enantioselectivity to 78%.

Ligand **84** gave excellent enantioselectivity for most of the reported substrates but some sensitivity to steric hindrance was also clear. Changing from  $\alpha$ -naphthyl

**Table 19** Asymmetric hydrogenation of imines

Imine	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Ligand ee%				
				<b>13a</b> <sup>a</sup>	<b>14d</b> <sup>b</sup>	<b>15b</b> <sup>c</sup>	<b>84</b> <sup>d</sup>	<b>85</b> <sup>e</sup>
<b>101a</b>	Ph	Et	Ph	78	82	92	–	–
<b>101b</b>	Ph	Et	4-MeO-C <sub>6</sub> H <sub>4</sub>	–	–	–	92	–
<b>101c</b>	Ph	Et	Ar <sup>f</sup>	–	–	–	–	94
<b>102</b>	α-naphth	Me	4-MeO-C <sub>6</sub> H <sub>4</sub>	–	–	–	98	–
<b>103a</b>	β-naphth	Me	Ph	91	–	–	–	–
<b>103b</b>	β-naphth	Me	4-MeO-C <sub>6</sub> H <sub>4</sub>	–	–	–	69	–
<b>103c</b>	β-naphth	Me	Ar <sup>f</sup>	–	–	–	–	93
<b>103c</b>	β-naphth	Me	Bn	–	–	93	–	–
<b>104a</b>			Ph	90	–	–	–	–
<b>104b</b>			4-MeO-C <sub>6</sub> H <sub>4</sub>	–	–	–	91	–
<b>104c</b>			Bn	–	–	96	–	–
<b>104d</b>			Me	–	–	98	–	–
<b>104e</b>			i-Bu	–	–	96	–	–

Condition <sup>a</sup>0.5 mol% cat, DCM, 20 bar H<sub>2</sub>

Condition <sup>b</sup>0.5 mol% cat, DCM, 5 bar H<sub>2</sub>, –20°C

Condition <sup>c</sup>1.0 mol% cat, DCM, 20 atm H<sub>2</sub>, 10°C

Condition <sup>d</sup>0.5 mol% cat, 2 mol% I<sub>2</sub>, toluene, 20 bar H<sub>2</sub>, 20 h

Condition <sup>e</sup>1.0 mol% cat, 4:1 toluene: methanol, 10 bar H<sub>2</sub>, 25°C, 20 h

Condition <sup>f</sup>Ar = 4-MeO-3,5-Me<sub>2</sub>-C<sub>6</sub>H<sub>2</sub>

methyl ketimine **102** with an ee of 98% for β-naphthyl-derived **103b** drops the enantioselectivity to 69%.

Inferior results with *N*-benzyl compared to *N*-aryl imines have been reported for many iridium-based systems [93]. Han et al. compensated for this same negative effect with SpinPHOX **15a** by replacing the pendant isopropyl group with a bulkier *tert*-butyl group in **15b**, demonstrating that the control of the system can be obtained by simple ligand modification. The added bulk of **15b** made it possible to reduce the very challenging *N*-alkyl-substituted imines **104d–e** in excellent enantioselectivity.

Knochel took a different approach by optimizing the substrate *N*-aryl group while investigating ligand **85**. Introduction of two meta-methyl groups in the 4-methoxyphenyl substituent increased the enantioselectivity from 84% with imine **86a** to 94% with **86c**.

Electronic effects seem to have very little influence on the overall enantioselectivity with both electron-donating and -withdrawing groups giving very small changes to the enantioselectivity when all other variables were the same. Substrate geometry seems to play a much more influential role with this particular class of substrates.

## References

1. Knowles WS (2002) *Angew Chem Int Ed* 41:1998
2. Blaser HU, Schmidt E (eds) (2004) *Asymmetric catalysis on industrial scale*. VCH, Weinheim
3. Brown JM (1999) Hydrogenation of functionalized carbon-carbon double bonds. In: Jacobsen EN, Pfaltz A, Yamamoto Y (eds) *Comprehensive asymmetric catalysis*, vol I. Springer, Berlin, Chapter 5.1
4. Noyori R (2002) *Angew Chem Int Ed* 41:2008
5. Lightfoot A, Schnider P, Pfaltz A (1998) *Angew Chem Int Ed* 37:2897
6. Pfaltz A, Blankenstein J, Hilgraf R, Hörmann E, McIntyre S, Menges F, Schönleber M, Smidt SP, Wüstenberg B, Zimmermann N (2002) *Adv Synth Catal* 345:33
7. Broene RD, Buchwald SL (1993) *J Am Chem Soc* 115:12569
8. Troutman MV, Appella DH, Buchwald SL (1999) *J Am Chem Soc* 121:4916
9. Conticello VP, Brard L, Giardello MA, Tsuji Y, Sabat M, Stern CL, Marks TJ (1992) *J Am Chem Soc* 114:2761
10. Crabtree RH (1979) *Acc Chem Res* 12:331
11. Cui X, Burgess K (2005) *Chem Rev* 105:3272
12. Källström K, Munslow I, Andersson PG (2006) *Chem Eur J* 12:3194
13. Roseblade SJ, Pfaltz A (2007) *Acc Chem Res* 40:1402
14. Church TL, Andersson PG (2008) *Coord Chem Rev* 252:513
15. Helmchen G, Pfaltz A (2000) *Acc Chem Res* 33:336
16. Blackmond DG, Lightfoot A, Pfaltz A, Rosner T, Schnider P, Zimmermann N (2000) *Chirality* 12:442
17. Smidt SP, Pfaltz A, Martínez-Viviente E, Pregosin PS, Albinati A (2003) *Organometallics* 22:1000
18. Krossing I, Raabe I (2004) *Angew Chem Int Ed* 43:2066
19. Smidt SP, Zimmermann N, Studer M, Pfaltz A (2004) *Chem Eur J* 10:4685
20. Wüstenberg B, Pfaltz A (2008) *Adv Synth Catal* 350:174
21. Crabtree RH, Demou PC, Eden D, Mihelcic JM, Parnell CA, Quirk JM, Morris GE (1982) *J Am Chem Soc* 104:6994
22. Mazet C, Smidt SP, Meuwly M, Pfaltz A (2004) *J Am Chem Soc* 126:14176
23. Burk MJ, McGrath MP, Wheeler R, Crabtree RH (1988) *J Am Chem Soc* 110:5034
24. Meuwly M, Roseblade SJ, Pfaltz A. University of Basel, Unpublished results
25. Roseblade SJ, Pfaltz AC (2007) *R Chimie* 10:178
26. Gridnev ID, Imamoto T (2004) *Acc Chem Res* 37:633
27. Halpern J (1982) *Science* 217:401
28. Brown JM, Chaloner PA (1978) *J Chem Soc Chem Commun*: 321
29. Feldgus S, Landis CR (2000) *J Am Chem Soc* 122:12714
30. Dietiker R, Chen P (2004) *Angew Chem Int Ed* 43:5513
31. Brandt P, Hedberg C, Andersson PG (2003) *Chem Eur J* 9:339
32. Hedberg C, Källström K, Brandt P, Hansen LK, Andersson PG (2006) *J Am Chem Soc* 128:2995
33. Fan Y, Cui X, Burgess K, Hall MB (2004) *J Am Chem Soc* 126:16688
34. Kaiser S, Smidt SP, Pfaltz A (2006) *Angew Chem Int Ed* 45:5194
35. Liu QB, Yu CB, Zhou YG (2006) *Tetrahedron Lett* 47:4733
36. Bell S, Wüstenberg B, Kaiser S, Menges F, Netscher T, Pfaltz A (2006) *Science* 311:642
37. Wang A, Wüstenberg B, Pfaltz A (2008) *Angew Chem Int Ed* 47:2298
38. Blankenstein J, Pfaltz A (2001) *Angew Chem Int Ed* 40:4445
39. Menges F, Pfaltz A (2002) *Adv Synth Catal* 344:40
40. Smidt SP, Menges F, Pfaltz A (2004) *Org Lett* 6:2023
41. Schrems MG, Pfaltz A (2009) *Chem Commun* 41:6210
42. Schönleber M, Hilgraf R, Pfaltz A (2008) *Adv Synth Catal* 350:2033

43. Li X, Li Q, Wu X, Gao Y, Xu D, Kong L (2007) *Tetrahedron Asymmetry* 18:629
44. Veredel JJ, Andersson PG (2007) *Dalton Trans* 47:5603
45. Tolstoy PI, Engman M, Paptchikhine A, Bergquist J, Church TL, Leung AWM, Andersson PG (2009) *J Am Chem Soc* 131:8855
46. Harmata M, Hong X (2005) *Org Lett* 7:3581
47. Bianco GG, Ferraz HMC, Costa AM, Costa-Lotufo LV, Pessoa CU, de Moraes MO, Schrems MG, Pfaltz A, Silva LF (2009) *J Org Chem* 74:2561
48. Engman M, Diesen JS, Paptchikhine A, Andersson PG (2007) *J Am Chem Soc* 129:4536
49. Engman M, Cheruku P, Tolstoy P, Bergquist J, Völker SF, Andersson PG (2009) *Adv Synth Catal* 351:375
50. Ueda M, Saitoh A, Miyaura NJ (2002) *Organomet Chem* 642:145
51. Morgan JB, Morken JP (2004) *J Am Chem Soc* 126:15338
52. Moran WJ, Morken JP (2006) *Org Lett* 8:2413
53. Paptchikhine A, Cheruku P, Engman M, Andersson PG (2009) *Chem Commun* 40:5996
54. Zhou J, Ogle JW, Fan Y, Banphavichit V, Zhu Y, Burgess K (2007) *Chem Eur J* 13:7162
55. Zhou J, Burgess K (2007) *Angew Chem Int Ed* 46:1129
56. Zhou J, Zhu Y, Burgess K (2007) *Org Lett* 9:1391
57. Zhu Y, Burgess K (2008) *J Am Chem Soc* 130:8894
58. Zhao J, Burgess K (2009) *Org Lett* 11:2053
59. Zhao J, Burgess K (2009) *J Am Chem Soc* 131:13236
60. Mantilli L, Mazet C (2009) *Tetrahedron Lett* 50:4141
61. Mantilli L, Gérard D, Torche S, Besnard C, Mazet C (2009) *Angew Chem Int Ed* 48:5143
62. Lu SM, Bolm C (2008) *Chem Eur J* 14:7513
63. Lu SM, Bolm C (2008) *Angew Chem Int Ed* 47:8920
64. Lu WJ, Chen YW, Hou XL (2008) *Angew Chem Int Ed* 47:10133
65. Lu WJ, Hou XL (2009) *Adv Synth Catal* 351:1224
66. Li S, Zhu SF, Zhang CM, Song S, Zhou QL (2008) *J Am Chem Soc* 130:8584
67. Cheruku P, Paptchikhine A, Church TL, Andersson PG (2009) *J Am Chem Soc* 131:8285
68. McIntyre S, Hörmann E, Menges F, Smidt SP, Pfaltz A (2005) *Adv Synth Catal* 347:282
69. Mazuela J, Veredel JJ, Coll M, Schäffner B, Börner A, Andersson PG, Pàmies O, Diéguez M (2009) *J Am Chem Soc* 131:12344
70. Schrems MG, Neumann E, Pfaltz A (2007) *Angew Chem Int Ed* 46:8274
71. Sprinz J, Helmchen G (1993) *Tetrahedron Lett* 34:1769
72. Zhu Y, Burgess K (2008) *Adv Synth Catal* 350:979
73. Cheruku P, Gohil S, Andersson PG (2007) *Org Lett* 9:1659
74. Zhou YG (2007) *Acc Chem Res* 40:1357
75. Cheruku P, Church TL, Trifonova A, Wartmann T, Andersson PG (2008) *Tetrahedron Lett* 49:7290
76. Baeza A, Pfaltz A (2009) *Chem Eur J* 15:2266
77. Hou GH, Xie JH, Yan PC, Zhou QL (2009) *J Am Chem Soc* 131:1366
78. Baeza A, Pfaltz A (2009) *Chem Eur J* 16:2036
79. Lu SM, Han XW, Zhou YG (2004) *Adv Synth Catal* 346:909
80. Lu SM, Bolm C (2008) *Adv Synth Catal* 350:1101
81. Legault CY, Charette AB (2005) *J Am Chem Soc* 127:8966
82. Blaser HU, Spindler F (1999) Hydrogenation of imino groups. In: Jacobsen EN, Pfaltz A, Yamamoto Y (eds) *Comprehensive asymmetric catalysis*, vol I. Springer, Berlin, Chapter 6.2
83. Willoughby CA, Buchwald SL (1994) *J Am Chem Soc* 116:11703
84. Blaser HU (2002) *Adv Synth Catal* 344:17
85. Schneider P, Koch G, Pretot R, Wang G, Bohnen FM, Kruger C, Pfaltz A (1997) *Chem Eur J* 3:887
86. Xiao D, Zhang X (2001) *Angew Chem Int Ed* 40:3425
87. Mršić N, Minnaard AJ, Feringa BL, Vries JG (2009) *J Am Chem Soc* 131:8358
88. Trifonova A, Diesen JS, Chapman CJ, Andersson PG (2004) *Org Lett* 6:3825



89. Trifonova A, Diesen JS, Andersson PG (2006) *Chem Eur J* 12:2318
90. Moessner C, Bolm C (2005) *Angew Chem Int Ed* 44:7564
91. Cheemala MN, Knochel P (2007) *Org Lett* 9:3089
92. Zhu SF, Xie JB, Zhang YZ, Li S, Zhou QL (2006) *J Am Chem Soc* 128:12886
93. Han Z, Wang Z, Zhang X, Ding K (2009) *Angew Chem Int Ed* 48:5345
94. Diéguez M, Mazuela J, Pàmies O, Verendel JJ, Andersson PG (2008) *J Am Chem Soc* 130:7208
95. Cheruku P, Diesen J, Andersson PG (2008) *J Am Chem Soc* 130:5595
96. Valla C, Baeza A, Menges F, Pfaltz A (2008) *Synlett* 3167
97. Baeza A, Pfaltz A (2010) *Chem Eur J* 16:4003

# Iridium-Catalyzed Hydrogen Transfer Reactions

Ourida Saidi and Jonathan M. J. Williams

**Abstract** This chapter describes the application of iridium complexes to catalytic hydrogen transfer reactions. Transfer hydrogenation reactions provide an alternative to direct hydrogenation for the reduction of a range of substrates. A hydrogen donor, typically an alcohol or formic acid, can be used as the source of hydrogen for the reduction of carbonyl compounds, imines, and alkenes. Heteroaromatic compounds and even carbon dioxide have also been reduced by transfer hydrogenation reactions. In the reverse process, the oxidation of alcohols to carbonyl compounds can be achieved by iridium-catalyzed hydrogen transfer reactions, where a ketone or alkene is used as a suitable hydrogen acceptor. The reversible nature of many hydrogen transfer processes has been exploited for the racemization of alcohols, where temporary removal of hydrogen generates an achiral ketone intermediate. In addition, there is a growing body of work where temporary removal of hydrogen provides an opportunity for using alcohols as alkylating agents. In this chemistry, an iridium catalyst “borrows” hydrogen from an alcohol to give an aldehyde or ketone intermediate, which can be transformed into either an imine or alkene under the reaction conditions. Return of the hydrogen from the catalyst provides methodology for the formation of amines or C–C bonds where the only by-product is typically water.

**Keywords** Alcohols · Alkenes · Asymmetric transfer hydrogenation · C-alkylation · Imines · Ketones · *N*-alkylation · Oxidation · Reduction · Transfer hydrogenation

## Contents

1	Introduction .....	78
2	Reduction of Ketones and Aldehydes by Transfer Hydrogenation .....	78
3	Reduction of Other Substrates by Transfer Hydrogenation .....	82
4	Asymmetric Transfer Hydrogenation of Ketones .....	85

5	Asymmetric Transfer Hydrogenation of Other Substrates .....	89
6	Oxidation of Alcohols .....	91
7	Tishchenko Reaction .....	93
8	Dynamic Kinetic Resolution Processes .....	94
9	Alkylation of Amines with Alcohols .....	96
10	C-Alkylation Reactions of Alcohols .....	100
	References .....	103

## 1 Introduction

Iridium complexes are amongst the most important for catalyzing hydrogen transfer reactions. Hydrogen transfer from a donor alcohol to an acceptor ketone has seen considerable attention, although other hydrogen donors and acceptors have also been used. In the presence of suitable enantiomerically pure ligands, many of these hydrogen transfer reactions have been performed asymmetrically, with high levels of enantioselectivity achieved in many cases. More recently, iridium catalysts have proved to be popular for the activation of alcohols by the temporary removal of hydrogen, which has allowed alcohols to be used as alkylating agents for nitrogen- and carbon-based nucleophiles.


## 2 Reduction of Ketones and Aldehydes by Transfer Hydrogenation

The reduction of ketones and aldehydes into the corresponding secondary and primary alcohols is one of the fundamental functional group transformations. The reaction is usually performed with either a stoichiometric reducing agent such as  $\text{NaBH}_4$  or by metal-catalyzed reduction. Whilst direct hydrogenation with  $\text{H}_2$  is completely atom efficient, transfer hydrogenation from a suitable hydrogen donor means that the use of potentially explosive  $\text{H}_2$  gas can be avoided. In many cases, 2-propanol is used as the hydrogen donor, and by using it as the solvent, it is present in sufficient excess that the equilibrium is driven towards the formation of the product.


There have been many reports of the use of iridium-catalyzed transfer hydrogenation of carbonyl compounds, and this section focuses on more recent examples where the control of enantioselectivity is not considered. In particular, recent interest has been in the use of iridium *N*-heterocyclic carbene complexes as active catalysts for transfer hydrogenation. However, alternative iridium complexes are effective catalysts [1, 2] and the air-stable complex **1** has been shown to be exceptionally active for the transfer hydrogenation of ketones [3]. For example, acetophenone **2** was converted into the corresponding alcohol **3** using only 0.001 mol% of this

catalyst as shown in Scheme 1. The availability of the hydrogen on the nitrogen donor was believed to be important for reactivity with concerted hydrogen transfer from both the NH and IrH to the ketone.

*N*-Heterocyclic carbene complexes of Ir(I) and Ir(III) have also demonstrated high reactivity in transfer hydrogenation reactions of ketones (Scheme 2) [4]. Complex **4** catalyzed the reduction of a range of ketones into the corresponding alcohols, including the reduction of pinacolone **7** into alcohol **8** with a low catalyst loading and short reaction time [5]. The chelating bis(*N*-heterocyclic carbene) complex **5** was shown to catalyze the reduction of ketones, and in the case of the reduction of benzophenone **9** to alcohol **10**, the reaction was complete within 4 min [6].



**Scheme 1** A highly active iridium catalyst for transfer hydrogenation




**Scheme 2** Use of *N*-heterocyclic carbene complexes of iridium

The same catalyst has also been used for the reduction of aldehydes to primary alcohols [7]. Several other iridium *N*-heterocyclic carbene complexes have been shown to be successful as catalysts for the transfer hydrogenation of ketones [8–12], including the interesting complex **6**, where the cyclopentadienyl ring is tethered to the *N*-heterocyclic carbene. Complex **6** was employed at low catalyst loading for the reduction of a range of ketones including the conversion of cyclohexanone **11** into cyclohexanol **12** [13].


The use of water as a solvent can have environmental advantages for some processes, and iridium-catalyzed transfer hydrogenation of carbonyl compounds has been achieved using water as the medium. The water-soluble hydride complex [Cp\*Ir(bipy)H]PF<sub>6</sub> (Cp\* = pentamethylcyclopentadienyl, bipy = 2,2'-bipyridine) catalyzes the reduction of ketones with formic acid as the hydrogen donor [14], where the fastest reactions occurred at pH 2–3. Xiao and coworkers have demonstrated that complex **13** formed from [Cp\*IrCl<sub>2</sub>]<sub>2</sub> and monotosylated ethylenediamine provides a highly active catalyst for the transfer hydrogenation of aldehydes using sodium formate [15, 16]. The reactions were described as being “on water and in air” as the catalyst is not soluble in water, and the presence of air did not appear to cause any problems. For example, the reduction of benzaldehyde **14** to give benzyl alcohol **15** was achieved with only 0.02 mol% catalyst in under an hour (Scheme 3). A polymer-bound variant of catalyst **13** was developed where the transfer hydrogenation of aldehydes occurred at the interface of emulsion droplets [17].

The mechanism for the iridium-catalyzed hydrogen transfer reaction between alcohols and ketones has been investigated, and there are three main reaction pathways that have been proposed (Scheme 4). Pathway (a) involves a direct hydrogen transfer where hydride transfer takes place between the alkoxide and ketone, which is simultaneously coordinated to the iridium center. Computational studies have given support to this mechanism for some iridium catalysts [18].

Alternative reaction pathways, which involve iridium hydride complexes, have also been presented. Bäckvall and coworkers have developed deuterium labeling experiments to probe whether hydrogen transfer reactions proceed through a monohydride or dihydride mechanism. In the case of Ir(I) complexes, the monohydride mechanism (b) is favored [19, 20]. Once an iridium hydride species is formed, the reaction pathway proceeds by coordination of the incoming ketone and hydride transfer. Ligand exchange with the alcohol acting as the hydrogen donor occurs and hydride transfer regenerates the iridium hydride species [21]. The concerted hydrogen transfer pathway (c) is available when the ligand contains an available hydrogen, as seen for complex **1** in Scheme 1. The iridium hydride transfers one hydrogen atom to the carbon of the incoming carbonyl compound, while the associated amine NH



**Scheme 3** Transfer hydrogenation of an aldehyde in water


**Scheme 4** Mechanistic pathways for transfer hydrogenation

provides the second hydrogen atom, which is transferred to the oxygen of the carbonyl compound. The catalyst is then restored to the iridium hydride by the reverse process of hydrogen delivery from the donor alcohol.

It seems likely that all three reaction pathways are feasible and the nature of the ligands determines which pathway is favored by a particular iridium catalyst.

### 3 Reduction of Other Substrates by Transfer Hydrogenation

Iridium catalysts have been used for the transfer hydrogenation of alkenes using alcohols as the hydrogen donor (Scheme 5). Even though the first report of iridium-catalyzed reduction of alkenes by hydrogen transfer appeared as early as 1967 [22], there has been much less attention given to this process in comparison with the reduction of carbonyl compounds.  $\text{IrCl}(\text{CO})(\text{PPh}_3)_2$  has been used to catalyze hydrogen transfer from formic acid to  $\alpha,\beta$ -unsaturated ketones [23], and a polymer-supported variant of this catalyst has also been shown to be effective [24]. Ishii and coworkers have developed the catalyst combination  $[\text{Ir}(\text{cod})\text{Cl}]_2/\text{dppp}/\text{Cs}_2\text{CO}_3$  [cod = 1,5-cyclooctadiene, dppp = 1,3-bis(diphenylphosphino)propane] for the transfer hydrogenation of  $\alpha,\beta$ -unsaturated carbonyl compounds [1].




**Scheme 5** Reduction of alkenes by transfer hydrogenation

The catalyst is also effective for the reduction of styrenes, ketones, and aldehydes. Cyclohexenone **16** was reduced to cyclohexanone **11** by transfer hydrogenation, and using a higher catalyst loading, styrene **17** was reduced to ethylbenzene **18**. The elaboration of  $[\text{Ir}(\text{cod})\text{Cl}]_2$  into the triazole-derived iridium carbene complex **19** provided a catalyst, which was used to reduce alkene **20** by transfer hydrogenation [25].

The transfer hydrogenation of imines has been achieved using iridium catalysts, with representative examples shown in Scheme 6. The same catalyst **19** was also effective for the transfer hydrogenation of imine **22** to give *N*-benzylaniline **23** after 30 min [25]. The reaction was also successful when the amine and aldehyde were added as separate components and imine formation taking place in situ prior to reduction. An alternative reductive amination procedure using iridium-catalyzed transfer hydrogenation has been achieved using ammonium formate as the nitrogen source and reducing agent. Catalyst **24** was used to convert a range of  $\alpha$ -keto acids into  $\alpha$ -amino acids, including the conversion of keto acid **26** into alanine **27** in water [26]. Corberan and Peris have shown that catalyst **25** is active for transfer hydrogenation reactions and have applied this to one example of imine reduction, which is achieved at room temperature [27].

Remarkably, complex **25** was also able to reduce  $\text{CO}_2$  by transfer hydrogenation in 2-propanol [28]. While there have been many reports using  $\text{H}_2$  to reduce  $\text{CO}_2$ , the work of Peris and coworkers is the only example of a hydrogen transfer reaction to reduce  $\text{CO}_2$  with 2-propanol [29]. The reduction is run in the presence of  $\text{KOH}$ ,



**Scheme 6** Reduction of imines by transfer hydrogenation






which allows the reduction product to be formed as the stable potassium formate salt (Scheme 7).


The catalytic reduction of nitro groups is usually achieved using heterogeneous catalysts, although the iridium complex **28** has been shown to be effective for the reduction of *p*-nitroanisole **29** to the corresponding aniline **30** using isopropanol as the hydrogen donor (Scheme 8) [30]. In the reduction of some nitroarenes, azo compounds (Ar–N=N–Ar) could be formed as by-products or as the major product by variation of the reaction conditions.

Fujita, Yamaguchi, and coworkers have used iridium-catalyzed transfer hydrogenation for the partial reduction of a range of quinolines (Scheme 9) [31]. For the parent quinoline **31**, [Cp\*IrCl<sub>2</sub>]<sub>2</sub> catalyzed the reduction to give 1,2,3,4-tetrahydroquinoline **32** in good yield. In some cases, perchloric acid was used as an additive to accelerate the reaction by protonation of the nitrogen atom. Crabtree and coworkers have identified a cationic iridium *N*-heterocyclic carbene complex as an alternative catalyst for partial reduction of quinolines using either 2-propanol or formic acid as the reducing agent [32].

**Scheme 7** Reduction of carbon dioxide by transfer hydrogenation



**Scheme 8** Reduction of a nitroarene by transfer hydrogenation




**Scheme 9** Partial reduction of quinoline by transfer hydrogenation

## 4 Asymmetric Transfer Hydrogenation of Ketones


The control of enantioselectivity in the reduction of carbonyl compounds provides an opportunity for obtaining the product alcohols in an enantiomerically enriched form. For transfer hydrogenation, such reactions have been dominated by the use of enantiomerically pure ruthenium complexes [33, 34], although Pfaltz and coworkers had shown by 1991 that high levels of enantioselectivity could be obtained using iridium(I) bis-oxazoline complexes [35].

In the late 1990s, two classes of enantiomerically pure iridium complexes were shown to be effective for the asymmetric transfer hydrogenation of aromatic ketones. Inoue and coworkers used the Ir(I) precursor  $[\text{Ir}(\text{cod})\text{Cl}]_2$  in conjunction with ligand **33** for the reduction of a range of ketones at room temperature with good enantioselectivity [36]. For example, propiophenone **36** was converted into the corresponding alcohol **37** with 93% ee (Scheme 10). Two groups independently reported that the combination of  $[\text{Cp}^*\text{IrCl}_2]_2$  or the analogous rhodium complex with amino sulfonamide ligands was capable of achieving very high enantioselectivities in the transfer



**Scheme 10** Asymmetric reduction of ketones with iridium catalysts







**Scheme 12** Further examples of asymmetric transfer hydrogenation with  $[\text{Ir}(\text{cod})\text{Cl}]_2$

There has been a growing interest in the development of aqueous phase asymmetric transfer hydrogenation of ketones [57, 58]. Water-soluble enantiomerically pure ligands **54** and **55** have been prepared [59, 60]. Ligand **54**, although highly successful with  $[\text{Cp}^*\text{RhCl}_2]_2$ , gave low conversions using  $[\text{Cp}^*\text{IrCl}_2]_2$  but still with good enantioselectivity. Sulfonated ligand **55** was used in a mixture of water and 2-propanol to reduce aromatic ketones including ketone **56** (Scheme 13). The presence of water was found to accelerate the rate of reduction, even though the concentration of 2-propanol was lowered.

Gao and coworkers have developed a tetra-sulfonated version of their ligand **42** and demonstrated that this can be used in 2-propanol/water (2:1) with similar levels of enantioselectivity to those obtained with the nonsulfonated ligand [61]. Xiao and coworkers have demonstrated that it is not necessary for the TsDPEN ligand to be derivatized for asymmetric transfer hydrogenation reactions to be successful in water [62]. They used unmodified TsDPEN **58**, which was precomplexed to  $[\text{Cp}^*\text{IrCl}_2]_2$  and then used to catalyze the reduction of over twenty aryl ketones using sodium formate as the reducing agent. Especially high enantioselectivities




**Scheme 13** Further examples of asymmetric transfer hydrogenation with  $[Cp^*IrCl_2]_2$




**Scheme 14** Asymmetric transfer hydrogenation in water

were obtained in the reduction of tetralone **38** and 2-acetylfuran **59** (Scheme 14). However, in all cases, the analogous rhodium catalysts were found to be more reactive and more selective.

Ward and coworkers have developed an interesting approach to the asymmetric transfer hydrogenation of ketones by designing a series of artificial metalloenzymes [63]. They prepared a range of metal complexes, including the iridium catalyst **61**, which were tethered to the biotin group. The biotin group has a high affinity for the protein Streptavidin, and once the catalyst was docked into the protein, the chiral environment around the iridium was exploited for the asymmetric transfer hydrogenation of ketones. In the case of the iridium complex **61**, a mutant of the wild type Streptavidin, S112Y Sav, was found to give a good level of asymmetric induction for the reduction of bromoacetophenone **62** to the corresponding alcohol **63** (Scheme 15).



**Scheme 15** Use of a protein to provide an asymmetric environment



**Scheme 16** Asymmetric reduction of imines with iridium catalysts


## 5 Asymmetric Transfer Hydrogenation of Other Substrates

The asymmetric transfer hydrogenation of imines using iridium catalysts has received less attention than with ruthenium catalysts, and the enantioselectivities appear to be lower for iridium catalysts. Haraguchi and coworkers have used the polymer-supported ligand **64** for the reduction of imine **65** using  $\text{Et}_3\text{N-HCO}_2\text{H}$  (2:5 azeotrope) as the hydrogen source (Scheme 16) [64]. When the catalyst precursor was  $[\text{Cp}^*\text{IrCl}_2]_2$ , the product **66** was formed with 77% ee, whereas a

higher ee could be obtained using  $[\text{RuCl}_2(p\text{-cymene})]_2$  as the catalyst precursor. There has been a report of a dynamic kinetic resolution reaction for the reduction of imines using iridium complex **34** [65]. Imine **67** is stereochemically labile under the reaction conditions due to ready tautomerization to the achiral enamine. Iridium complex **34** catalyzed the reduction of imine **67** to give the amine **68** with reasonable enantioselectivity. However, when the isoelectronic  $\text{Ru}(p\text{-cymene})$  complex was used, enantioselectivities of up to 98% ee could be obtained for related substrates.

In spite of the success of asymmetric iridium catalysts for the direct hydrogenation of alkenes, there has been very limited research into the use of alternative hydrogen donors. Carreira and coworkers have reported an enantioselective reduction of nitroalkenes in water using formic acid and the iridium aqua complex **69** [66]. For example, the reduction of nitroalkene **70** led to the formation of the product **71** in good yield and enantioselectivity (Scheme 17). The use of other aryl substrates afforded similar levels of enantioselectivity.


The asymmetric reduction of quinolines has been achieved using Hantzsch esters as the hydrogen donor [67]. Loss of hydrogen from Hantzsch ester **72** leads to the corresponding pyridine, which provides a strong driving force for hydrogen donation. The iridium complex  $[\text{Ir}(\text{cod})\text{Cl}]_2$  is complexed with (*S*)-Segphos **73** and activated with iodine and then used as the catalyst for the transfer hydrogenation of 2-methylquinoline **74** (Scheme 17). The asymmetric reduction of substrate **74** has also been reported using sodium formate as the reducing agent with an in situ catalyst formed from  $[\text{Cp}^*\text{IrCl}_2]_2$  and TsDPEN, although the product **75** was formed with only 11% ee [68]. This contrasts with the corresponding rhodium catalyst where the product was formed with 96% ee.



**Scheme 17** Asymmetric reduction of other substrates with iridium catalysts

## 6 Oxidation of Alcohols

The use of metal-catalyzed hydrogen transfer reactions for alcohol oxidation provides an alternative approach to the use of noncatalytic stoichiometric oxidants. Various alcohols and diols have been oxidized using iridium catalysts in the presence of a ketone as a hydrogen acceptor (Scheme 18). The Oppenauer-type oxidation of primary and secondary alcohols using  $[\text{Cp}^*\text{IrCl}_2]_2$  in the presence of  $\text{K}_2\text{CO}_3$  has been shown to be effective [69]. Even at room temperature, benzyl alcohol **15** could be converted into benzaldehyde **14** using acetone as both the solvent and hydrogen acceptor (Scheme 18). No over-oxidation to carboxylic acids or esters was observed. Secondary alcohols including 1-phenethyl alcohol **3** were readily converted into the corresponding ketones. Modifications of the  $[\text{Cp}^*\text{IrCl}_2]_2$  complex have been found to give improvements in turnover number in alcohol oxidation reactions. The *N*-heterocyclic carbene complex **76** could be used with




**Scheme 18** Oxidation of alcohols by hydrogen transfer




lower catalyst loadings than for the parent complex [70, 71]. The iridium complex **77** has also been used as a catalyst for the oxidation of alcohols (Scheme 18). Using either butanone or acetone as the solvent and hydrogen acceptor, primary alcohols were selectively oxidized to aldehydes. Electron-rich benzylic alcohols were particularly good substrates, with aliphatic alcohols performing less well. The oxidation of cinnamyl alcohol **78** gave cinnamaldehyde **79** in good yield [72], while the oxidation of diols led to the formation of lactones via intermediate hydroxy aldehydes [73]. Cyclization to the lactol and further oxidation led to formation of the lactone. For example, diol **80** was oxidized to give lactone **81** under mild reaction conditions.

Asymmetric variants of alcohol oxidation reactions have been reported, including the desymmetrization of meso diols, to give lactones using  $[\text{Cp}^*\text{IrCl}_2]_2$  in the presence of an enantiomerically pure amino alcohol. In the best example reported, meso diol **82** was converted into the enantiomerically enriched lactone **83** using  $[\text{Cp}^*\text{IrCl}_2]_2$ /ligand **85**/*t*-BuOK as the catalyst [74]. Related chemistry involving the desymmetrization of meso diol **85** gave the hydroxy ketone **86** with over 99% ee, using the preformed catalyst (*R,R*)-**34** (Scheme 19) [75].


Complex **77** has also been reported to catalyze the oxidative dimerization of alcohols to esters when the reactions are performed in the presence of base [76]. The presence of base presumably encourages the reversible attack of the alcohol onto the initially formed aldehyde to give a hemiacetal, which is further oxidized to give the ester product. Alcohols **87** and **15** were converted into esters **88** and **89** with good isolated yields (Scheme 20). Alternative iridium catalysts have been used for related oxidative dimerization reactions, and the addition of base is not always a requirement for the reaction to favor ester formation over aldehyde formation [77, 78].



**Scheme 19** Asymmetric oxidation of diols by desymmetrization



**Scheme 20** Oxidative dimerization of alcohols to give esters




**Scheme 21** Iridium catalyzed conversion of an alcohol to a primary amide


The iridium-catalyzed oxidation of benzylic alcohols has been coupled with an in situ oxime formation. The iridium catalyst then fulfills a second role by catalyzing the rearrangement of the oxime into an amide. For example, benzyl alcohol **15** underwent oxidation to benzaldehyde using styrene as the hydrogen acceptor in the presence of  $[\text{Cp}^*\text{IrCl}_2]_2$ , as shown in Scheme 21. Addition of hydroxylamine hydrochloride to the reaction mixture rapidly generated benzaldehyde oxime, which after, further heating, rearranged to give benzamide **90** in good isolated yield [79]. The reaction of 2-aminobenzyl alcohol with enolizable ketones has been catalyzed by iridium complexes to give quinolines as the oxidized products [80]. Two equivalents of ketone were used in order to drive the oxidation process.

## 7 Tishchenko Reaction

The dimerization of aldehydes to form esters is a completely atom-efficient process known as the Tishchenko reaction, which involves no net oxidation or reduction. Suzuki, Katoh, and coworkers have used complex **77** to catalyze the Tishchenko reaction of a range of aldehydes, including dihydrocinnamaldehyde **91** and benzaldehyde **14** (Scheme 22) [81]. The same catalyst has been used for an intramolecular variant of the reaction, where keto-aldehyde **92** isomerizes to lactone **93** via an intramolecular Tishchenko reaction. The oxidized product is formed as a by-product,



**Scheme 22** Tishchenko reaction catalysed by iridium complexes



**Scheme 23** Oxidation of an aldehyde to an ester

and by running the reaction in the presence of an additional aldehyde as a hydrogen acceptor, the oxidized species **94** is formed as the sole product [82].


A related report has demonstrated that aldehydes can be converted into esters in the presence of allylic alcohols [83]. Dihydrocinnamaldehyde **91** and allyl alcohol **95** undergo a cross-condensation reaction to give ester **96**. The allyl alcohol **95** also serves as a hydrogen acceptor, perhaps via an isomerization/reduction sequence (Scheme 23).

## 8 Dynamic Kinetic Resolution Processes


The reversibility of hydrogen transfer reactions has been exploited for the racemization of alcohols and amines. By coupling the racemization process with an enantioselective enzyme-catalyzed acylation reaction, it has been possible to achieve dynamic kinetic resolution reactions. The combination of lipases or

esterases with various ruthenium catalysts has been widely employed for these reactions [84], although there are also a few reports of combining enzymes with iridium catalysts to achieve dynamic kinetic resolution reactions [85–88]. The general principle of the dynamic resolution process is outlined in Scheme 24. Interconversion of alcohol (*R*)-**97** with its enantiomer (*S*)-**97** can be achieved by the temporary removal of hydrogen via the achiral ketone **98**. The enzyme selectively acylates one enantiomer of the alcohol to give the enantiomerically enriched ester (*R*)-**99**, which is inert to racemization under the reaction conditions.

Marr and coworkers have used the combination of the iridium *N*-heterocyclic carbene complex **100** (0.1 mol%) with the lipase Novozyme 435 to convert racemic alcohols **101** into enantiomerically enriched esters **102** with excellent yield and enantioselectivities (Scheme 25) [85]. Peris and coworkers have used complex **25**



**Scheme 24** Dynamic kinetic resolution strategy



**Scheme 25** Examples of iridium catalyzed dynamic kinetic resolution


to effect the racemization of a branched primary alcohol and used this in combination with an enzyme-catalyzed acylation in order to perform a dynamic kinetic resolution [27].

$[\text{Cp}^*\text{IrI}_2]_2$  has been used by Blacker and coworkers as an efficient catalyst for the racemization of amines [86, 87]. The catalyst reversibly converts the amine into an achiral imine in a similar way to that seen for alcohol racemization. Racemic amine **103** underwent dynamic kinetic resolution to form the carbamate **104** with high enantioselectivity in the presence of *Candida rugosa* lipase and the iridium catalyst (Scheme 25). Feringa, deVries, and coworkers have reported that the reaction of  $[\text{CpIrI}_2]_2$  with benzylic amines generates iridacyclic complexes, which are also competent catalysts for the racemization of alcohols and amines [88].


## 9 Alkylation of Amines with Alcohols

The ability of iridium complexes to interconvert alcohols and carbonyl compounds has been developed to allow alcohols to act as alkylating agents. Alcohols are generally less toxic than the corresponding alkyl halides, and this methodology provides a more environmentally benign approach to the alkylation of amines. The mechanism is believed to proceed via the temporary removal of hydrogen from the alcohol **106** to give an intermediate aldehyde **107** (Scheme 26). The intermediate aldehyde **107** is more electrophilic than the original alcohol and will readily react with an amine **108** to form an imine **109**. Whether the imine formation occurs within the coordination sphere of the metal is debatable and may depend on the particular catalyst that is being used. Return of the hydrogen from the catalyst reduces the imine into the amine **110** with no net oxidation or reduction. The overall process has been termed “borrowing hydrogen,” and the area has been reviewed recently [89–93].

The most widely used iridium catalyst for the *N*-alkylation of amines with alcohols is  $[\text{Cp}^*\text{IrCl}_2]_2$ , which was identified and developed by Fujita, Yamaguchi, and coworkers [94]. Aniline **111** has been selectively alkylated by a range of




**Scheme 26** Alkylation of amines with alcohols by borrowing hydrogen




**Scheme 27** Examples of amine alkylation reactions

alcohols including benzylic alcohols and primary and secondary alcohols (Scheme 27)[95]. Good isolated yields were generally obtained, including the formation of products **112** and **113** from the reaction of aniline **111** with either benzyl alcohol **15** or cyclohexanol **12**. Secondary amines could also be alkylated under similar reaction conditions. Amine **114** was alkylated with benzyl alcohol **15**, and pyrrolidine **116** was alkylated with octanol **117** to give the corresponding tertiary amine products **115** and **118** [96].

Several cyclizations of amino alcohols have been reported (Scheme 28) [97], including the ring closure of aniline **119** to give tetrahydroisoquinoline **120** [98]. Interestingly, when the reaction was run on the substrate with one less methylene group in the tether, the five-membered ring product was formed as the indole with loss of hydrogen. Cyclization reactions involving diols and either amines or diamines have been reported. Nordstrøm and Madsen cyclized diamine **121** with ethylene glycol **122** to give the bicyclic piperazine **123** [99]. The reactions could be run in water, and although the presence of base was beneficial for these reactions, it was not essential. In a separate study, Williams and coworkers showed that base was generally not needed for amination reactions catalyzed by  $[\text{Cp}^*\text{IrCl}_2]_2$  when they were run in water or ionic liquids [100]. Cyclization reactions of primary amines with diols provide an interesting approach to the synthesis of *N*-heterocycles. In one example, the formation of morpholine **126** was achieved by the reaction of amine **124** with diol **125** [101, 102].



**Scheme 28** The use of diols to give cyclic amines




**Scheme 29** Alkylation of sulfonamides with alcohols

Recently, the *N*-alkylation of sulfonamides using  $[\text{Cp}^*\text{IrCl}_2]_2$  as the catalyst has been reported using either water [100] or toluene [103] as the solvent (Scheme 29). *p*-Toluenesulfonamide **127** was alkylated with a range of over 20 alcohols. When benzylic alcohols were used, it was possible to use a low catalyst loading in order to achieve reaction. The catalytic species formed in the reactions was a bridged iridium dimer complex,  $[(\text{Cp}^*\text{Ir})(\mu\text{-NTs})_2]$ . Eisenstein, Crabtree, and coworkers briefly examined the alkylation of *p*-toluenesulfonamide **127** in the context of a mechanistic investigation [104]. Computational studies had shown that while alcohol and imine complexation to iridium were requirements for *N*-alkylation reactions, the basic amines coordinate more strongly, lowering the rate of reaction. The faster reaction observed using *p*-toluenesulfonamide, which is less basic than conventional amines, provided experimental support for the theoretical calculations.


Under more forcing reaction conditions,  $[\text{Cp}^*\text{IrCl}_2]_2$  has been reported to convert primary amines into secondary amines with loss of ammonia [105]. At 170°C, benzylamine **129** was converted into dibenzylamine **130**, with only trace amounts

of tertiary amines being formed (Scheme 30). The cross-coupling of two amines has been achieved with high selectivity where the iodide complex  $[\text{Cp}^*\text{IrI}_2]_2$  was found to be more reactive [106]. Thus, benzylamine **129** reacted with diisopropylamine **131** to give the secondary amine **132**. These reactions are believed to proceed via oxidation of the diisopropylamine **131** to an imine, which is then attacked by the primary amine with loss of isopropylamine and subsequent reduction to give the amine product.

Complexation of  $[\text{Cp}^*\text{IrCl}_2]_2$  with *N*-heterocyclic carbenes has led to complexes such as **25**, developed by Peris and coworkers [107, 108], and **133**, developed by Crabtree and coworkers [12]. Complex **24** is activated by the addition of silver triflate and is effective for the *N*-alkylation of amines with alcohols and for the *N*-alkylation of anilines with primary amines. Complex **25** has also been shown to couple benzyl alcohol **15** with a range of alcohols, including ethanol **134**, to give ether products such as ether **135** (Scheme 31). Complex **133** was an active hydrogen transfer catalyst for the reduction of ketones and imines, using 2-propanol as the hydrogen source. It was also an effective catalyst for the *N*-alkylation of amines




**Scheme 30** Iridium catalyzed amine coupling



**Scheme 31** Conversion of alcohols into ethers





**Scheme 32** Further examples of amine formation from alcohols

with alcohols and the  $\beta$ -alkylation of secondary alcohols with primary alcohols (see Sect. 10).


Other iridium complexes that do not contain the  $\text{Cp}^*$  ligand have also been found to be capable of catalyzing *N*-alkylation reactions (Scheme 32). Liu and coworkers have reported that *N*-heterocyclic carbene complexes of the type  $(\text{NHC})\text{Ir}(\text{CO})(\text{PPh}_3)\text{Cl}$  are effective catalysts [109]. The *N*-alkylation of amines with alcohols has been catalyzed by  $[\text{Ir}(\text{cod})\text{Cl}]_2$  in the presence of dpfp (bis(diphenylphosphino)ferrocene) [110]. The same catalyst combination was also used for the formation of alkylated anilines using iminophosphorane **136** as the nitrogen source [111]. In the reaction with benzyl alcohol **15**, the reaction was believed to proceed by oxidation to benzaldehyde followed by imine formation via an aza-Wittig reaction with iminophosphorane **136** and return of hydrogen to generate the amine product **112**. Kempe and coworkers have found that ligand **137** greatly enhanced the activity of  $[\text{Ir}(\text{cod})\text{Cl}]_2$  and was an effective catalyst even at  $70^\circ\text{C}$  for the alkylation of aniline **111** with benzyl alcohol **15** to give *N*-benzyl aniline **112** [112]. In particular, the combination of  $[\text{Ir}(\text{cod})\text{Cl}]_2$  with ligand **137** was effective for the monoalkylation of 2-aminopyridine **138** with a range of alcohols, including benzyl alcohol **15**, at low catalyst loading.

## 10 C-Alkylation Reactions of Alcohols


The activation of alcohols by iridium-catalyzed borrowing hydrogen reactions has also been applied to the formation of C–C bonds [113]. Reactions proceed by temporary removal of hydrogen from an alcohol to give an aldehyde (or ketone), which is transformed into an alkene with subsequent return of the hydrogen to provide a new C–C bond. Ishii and coworkers have used  $[\text{Ir}(\text{cod})\text{Cl}]_2$  with

triphenylphosphine and potassium hydroxide to catalyze the  $\alpha$ -alkylation of ketones with alcohols [114]. Typical examples include the reaction of ketones **140** and **38** with 1-butanol **141** to give the butylated products **142** and **143** (Scheme 33). For the reaction with ketone **140**, the alkylation was completely selective for reaction on the less substituted side of the ketone. Related iridium-catalyzed  $\alpha$ -alkylation of ketones with alcohols has been reported where the product ketone is reduced in situ by transfer hydrogenation with an enantiomerically pure ruthenium complex [115].

Fujita, Yamaguchi, and coworkers have coupled secondary alcohols with primary alcohols using  $[\text{Cp}^*\text{IrCl}_2]_2$  as the catalyst [116]. The reaction occurs by removal of hydrogen from the secondary alcohol **3** and the primary alcohol **141** to give ketone **2** and aldehyde **144** intermediates (Scheme 34). An aldol condensation of these intermediates leads to the  $\alpha,\beta$ -unsaturated ketone **145**, which is reduced to the saturated alcohol **146** by the return of two equivalents of hydrogen.




**Scheme 33** Formation of C-C bonds using alcohols




**Scheme 34** Alcohol coupling reactions by borrowing hydrogen

similar reaction conditions [117]. A selfcondensation of primary alcohols has also been reported using either  $[\text{Ir}(\text{cod})\text{Cl}]_2$  or  $[\text{Cp}^*\text{IrCl}_2]_2$  [118].

A variety of other carbon nucleophiles have been alkylated with alcohols including malonate esters, nitroalkanes, ketonitriles [119, 120], barbituric acid [121], cyanoesters [122], arylacetonitriles [123], 4-hydroxycoumarins [124], oxindoles [125], methylpyrimidines [126], indoles [127], and esters [128]. Selected examples are given in Scheme 35. Thus, benzyl alcohol **15** could be alkylated with nitroethane **147**, 1,3-dimethylbarbituric acid **148**, phenylacetonitrile **149**, methylpyrimidine **150**, and even *t*-butyl acetate **151** to give the corresponding alkylated products **152–156**.



**Scheme 35** Further examples of alcohols as alkylating agents



**Scheme 36** Enantioselective variant of borrowing hydrogen

The formation of C–C bonds from alcohols using iridium-catalyzed borrowing hydrogen chemistry has also been reported using Wittig reagents and Horner Wadsworth Emmons reagents for the alkene-forming step in the reaction pathway [129, 130]. In the example in Scheme 36, the reaction was performed using an enantiomerically pure iridium catalyst, which was used to couple phosphonium ylid, **157** with benzyl alcohol **15** to give the product **158** with reasonably good control of enantioselectivity [131].

## References

1. Sakaguchi S, Yamaga T, Ishii Y (2001) *J Org Chem* 66:4710–4712
2. Tsoureas N, Owen GR, Hamilton A, Orpen AG (2008) *Dalton Trans* 6039–6044
3. Clarke ZE, Maragh PT, Dasgupta TP, Gusev DG, Lough AJ, Abdur-Rashid K (2006) *Organometallics* 25:4113–4117
4. Normand AT, Cavell KJ (2008) *Eur J Inorg Chem* 2781–2800
5. Hillier AC, Lee HM, Stevens ED, Nolan SP (2001) *Organometallics* 20:4246–4252
6. Albrecht M, Miecznikowski JR, Samuel A, Faller JW, Crabtree RH (2002) *Organometallics* 21:3596–3604
7. Miecznikowski JR, Crabtree RH (2004) *Organometallics* 23:629–631
8. Kownacki I, Kubicki M, Szubert K, Marciniak B (2008) *J Organomet Chem* 693:321–328
9. Türkmen H, Pape T, Hahn FE, Çetinkaya B (2008) *Eur J Inorg Chem* 5418–5423
10. Sun J-F, Chen F, Dougan BA, Xu H-J, Cheng Y, Li Y-Z (2009) *J Organomet Chem* 694:2096–2105
11. Zinner SC, Rentsch CF, Herdtweck E, Herrmann WA, Kühn FE (2009) *Dalton Trans* 7055–7062
12. Gnanamgari D, Sauer ELO, Schley ND, Butler C, Incarvito CD, Crabtree RH (2009) *Organometallics* 28:321–325
13. da Costa PA, Viciano M, Sanaú M, Merino S, Tejada J, Peris E, Royo B (2008) *Organometallics* 27:1305–1309
14. Abura T, Ogo S, Watanabe Y, Fukuzumi S (2003) *J Am Chem Soc* 125:4149–4154
15. Wu X, Liu J, Li X, Zanotti-Gerosa A, Hancock F, Vinci D, Ruan J, Xiao J (2006) *Angew Chem Int Ed* 45:6718–6722
16. Wu X, Corcoran C, Yang S, Xiao J (2008) *ChemSusChem* 1:71–74
17. Li J, Zhang Y, Han D, Jia G, Gao J, Zhong L, Li C (2008) *Green Chem* 10:608–611
18. Handgraaf J-W, Reek JNH, Meijer EJ (2003) *Organometallics* 22:3150–3157
19. Pàmies O, Bäckvall J-E (2001) *Chem Eur J* 7:5052–5058
20. Samec JSM, Bäckvall J-E, Andersson PG, Brandt P (2006) *Chem Soc Rev* 35:237–248
21. Zhao J, Hesslink H, Hartwig JF (2001) *J Am Chem Soc* 123:7220–7227
22. Trocha-Grimshaw J, Henbest HB (1967) *Chem Commun* 544
23. Blum J, Sasson Y, Iflah S (1972) *Tetrahedron Lett* 1015

24. Azran J, Buchman O (1984) *J Org Chem* 49:1327–1333
25. Gnanamgari D, Moores A, Rajaseelan E, Crabtree RH (2007) *Organometallics* 26:1226–1230
26. Ogo S, Uehara K, Abura T, Fukuzumi S (2004) *J Am Chem Soc* 126:3020–3021
27. Corberán R, Peris E (2008) *Organometallics* 27:1954–1958
28. Sanz S, Benítez M, Peris E (2010) *Organometallics* 29:275–277
29. Himeda Y (2007) *Eur J Inorg Chem* 3927–3941
30. Wang C-Y, Fu C-F, Liu Y-H, Peng S-M, Liu S-T (2007) *Inorg Chem* 46:5779–5786
31. Fujita K-I, Kitatsuji C, Furukawa S, Yamaguchi R (2004) *Tetrahedron Lett* 45:3215–3217
32. Voutchkova AM, Gnanamgari D, Jakobsche CE, Butler C, Miller SJ, Parr J, Crabtree RHJ (2008) *Organomet Chem* 693:1815–1821
33. Palmer MJ, Wills M (1999) *Tetrahedron Asymmetry* 10:2045–2061
34. Wang C, Wu X, Xiao J (2008) *Chem Asian J* 3:1750–1770
35. Müller D, Umbrecht G, Weber B, Pfaltz A (1991) *Helv Chim Acta* 74:232
36. Inoue S-I, Nomura K, Hashiguchi S, Noyori R, Izawa Y (1997) *Chem Lett* 957–958
37. Blacker AJ, Mellor B WO9842643B1, Avecia Ltd., filed 26/03/97
38. Mashima K, Abe T, Tani K (1998) *Chem Lett* 1199–1200
39. Mashima K, Abe T, Tani K (1998) *Chem Lett* 1201–1202
40. Murata K, Ikariya T (1999) *J Org Chem* 64:2186–2187
41. Noyori R, Hashiguchi S (1997) *Acc Chem Res* 30:97–102
42. Li X, Blacker J, Houson I, Wu X, Xiao J (2006) *Synlett* 1155–1160
43. Ohkuma T, Utsumi N, Watanabe M, Tsutsumi K, Arai N, Murata K (2007) *Org Lett* 9:2565–2567
44. Furegati M, Rippert AJ (2005) *Tetrahedron Asymmetry* 16:3947–3950
45. Chen J-S, Li Y-Y, Dong Z-R, Li B-Z, Gao J-X (2004) *Tetrahedron Lett* 45:8415–8418
46. Li Y-Y, Zhang H, Chen J-S, Liao X-L, Dong Z-R, Gao J-X (2004) *J Mol Catal* 218:153–156
47. Dong Z-R, Li Y-Y, Chen J-S, Li B-Z, Xing Y, Gao J-X (2005) *Org Lett* 7:1043–1045
48. Debono N, Besson M, Pinel C, Djakovitch L (2004) *Tetrahedron Lett* 45:2235–2238
49. Paredes P, Díez J, Gamasa MP (2008) *Organometallics* 27:2597–2607
50. McManus HH, Barry SM, Andersson PG, Guiry PJ (2004) *Tetrahedron* 60:3405–3416
51. Dyson G, Frison J-C, Whitwood AC, Douthwaite RE (2009) *Dalton Trans* 7141–7151
52. Fuentes JA, France MB, Slawin AMZ, Clarke ML (2009) *New J Chem* 33:466–470
53. Maire P, Breher F, Schönberg H, Grützmacher H (2005) *Organometallics* 24:3207–3218
54. He W, Zhang B-L, Jiang R, Liu P, Sun X-L, Zhang S-Y (2006) *Tetrahedron Lett* 47:5367–5370
55. Lundgren RJ, Stradiotto M (2008) *Chem Eur J* 14:10388–10395
56. Petra DGI, Kamer PCJ, Spek AL, Schoemaker HE, Van Leeuwen PWNM (2000) *J Org Chem* 65:3010–3017
57. Wu X, Xiao J (2007) *Chem Commun* 2449–2466
58. Wu X, Wang C, Xiao J (2010) *Platinum Metals Rev* 54:3–19
59. Li L, Wu J, Wang F, Liao J, Zhang H, Lian C, Zhu J, Deng J (2007) *Green Chem* 9:23–25
60. Thorpe T, Blacker J, Brown SM, Bubert C, Crosby J, Fitzjohn S, Muxworthy JP, Williams MJM (2001) *Tetrahedron Lett* 42:4041–4043
61. Li B-Z, Chen J-S, Dong Z-R, Li Y-Y, Li Q-B, Gao J-X (2006) *J Mol Catal Chem* 258:113–117
62. Wu X, Li X, Zanotti-Gerosa A, Pettman A, Liu J, Mills AJ, Xiao J (2008) *Chem Eur J* 14:2209–2222
63. Letondor C, Pordea A, Humbert N, Ivanova A, Mazurek S, Novic M, Ward TR (2006) *J Am Chem Soc* 128:8320–8328
64. Haraguchi N, Tsuru K, Arakawa Y, Itsuno S (2009) *Org Biomol Chem* 7:69–75
65. Ros A, Magriz A, Dietrich H, Ford M, Fernández R, Lassaletta JM (2005) *Adv Synth Catal* 347:1917–1920
66. Soltani O, Ariger MA, Carreira EM (2009) *Org Lett* 11:4196–4198

67. Wang D-W, Zeng W, Zhou Y-G (2007) *Tetrahedron Asymmetry* 18:1103–1107
68. Wang C, Li C, Wu X, Pettman A, Xiao J (2009) *Angew Chem Int Ed* 48:6524–6528
69. Fujita K-I, Furukawa S, Yamaguchi R (2002) *Organomet Chem* 649:289–292
70. Hanasaka F, Fujita K-I, Yamaguchi R (2005) *Organometallics* 24:3422–3433
71. Hanasaka F, Fujita K-I, Yamaguchi R (2006) *Organometallics* 25:4643–4647
72. Suzuki T, Morita K, Tsuchida M, Hiroi K (2003) *J Org Chem* 68:1601–1602
73. Suzuki T, Morita K, Tsuchida M, Hiroi K (2002) *Org Lett* 4:2361–2363
74. Suzuki T, Morita K, Matsuo Y, Hiroi K (2003) *Tetrahedron Lett* 44:2003–2006
75. Suzuki T, Ghazati K, Katoh T, Sasai H (2009) *Org Lett* 11:4286–4288
76. Suzuki T, Matsuo T, Watanabe K, Katoh T (2005) *Synlett* 9:1453–1455
77. Izumi A, Obora Y, Sakaguchi S, Ishii Y (2006) *Tetrahedron Lett* 47:9199–9201
78. Yamamoto N, Obora Y, Ishii Y (2009) *Chem Lett* 38:1106–1107
79. Owston NA, Parker AJ, Williams MJ (2007) *Org Lett* 9:73–75
80. Taguchi K, Sakaguchi S, Ishii Y (2005) *Tetrahedron Lett* 46:4539–4542
81. Suzuki T, Yamada T, Matsuo T, Watanabe K, Katoh T (2005) *Synlett* 1450–1452
82. Suzuki T, Yamada T, Watanabe K, Katoh T (2005) *Bioorg Med Chem Lett* 15:2583–2585
83. Kiyooka S-I, Ueno M, Ishii E (2005) *Tetrahedron Lett* 46:4639–4642
84. Paines O, Bäckwall J-E (2003) *Chem Rev* 103:3247–3262
85. Marr AC, Pollock CL, Saunders GC (2007) *Organometallics* 26:3283–3285
86. Blacker AJ, Stirling MJ, Page MI (2007) *Org Process Res Dev* 11:642–648
87. Stirling MJ, Blacker AJ, Page MI (2007) *Tetrahedron Lett* 48:1247–1250
88. Jerphagnon T, Gayet AJA, Berthiol F, Rittleng V, Mršić N, Meetsma A, Pfeffer M, Minnaard AJ, Feringa BL, De Vries JG (2009) *Chem Eur J* 15:12780–12790
89. Lamb GW, Williams JM (2008) *J Chimica Oggi* 26(3):17–18
90. Hamid MHSA, Slatford PA, Williams MJ (2007) *Adv Synth Catal* 349:1555–1575
91. Nixon TD, Whittlesey MK, Williams MJ (2009) *Dalton Trans* 753–762
92. Dobereiner GE, Crabtree RH (2010) *Chem Rev* 110:681–703
93. Guillena G, Ramón DJ, Yus M (2010) *Chem Rev* 110:1611–1641
94. Fujita K-I, Yamaguchi R (2005) *Synlett* 4:560–571
95. Fujita K-I, Li Z, Ozeki N, Yamaguchi R (2003) *Tetrahedron Lett* 44:2687–2690
96. Fujita K-I, Enoki Y, Yamaguchi R (2008) *Tetrahedron* 64:1943–1954
97. Eary CT, Clausen D (2006) *Tetrahedron Lett* 47:6899–6902
98. Fujita K-I, Yamamoto K, Yamaguchi R (2002) *Org Lett* 4:2691–2694
99. Nordström LU, Madsen R (2007) *Chem Commun* 5034–5036
100. Saidi O, Blacker AJ, Lamb GW, Marsden SP, Taylor JT, Williams MJ (2010) *Org Process Res Dev* 14:1046–1049
101. Fujita K-I, Fujii T, Yamaguchi R (2004) *Org Lett* 6:3525–3528
102. Yamaguchi R, Kawagoe S, Asai S, Fujita K-I (2008) *Org Lett* 10:181–184
103. Zhu M, Fujita K-I, Yamaguchi R (2010) *Org Lett* 12:1336–1339
104. Balcells D, Nova A, Clot E, Gnanamgari D, Crabtree RH, Eisenstein O (2008) *Organometallics* 27:2529–2535
105. Lorentz-Petersen LLR, Jensen P, Madsen R (2009) *Synthesis* 4110–4112
106. Saidi O, Blacker AJ, Farah M, Marsden SP, Williams MJ (2009) *Angew Chem Int Ed* 48:7375
107. Prades A, Corberán R, Poyatos M, Peris E (2008) *Chem Eur J* 14:11474–11479
108. Corberán R, Mas-Marzá E, Peris E (2009) *Eur J Inorg Chem* 1700–1716
109. Chang Y-H, Fu C-F, Liu Y-H, Peng S-M, Chen J-T, Liu S-T (2009) *Dalton Trans* 861–867
110. Cami-Kobeci G, Slatford PA, Whittlesey MK, Williams MJ (2005) *Bioorg Med Chem Lett* 15:535–537
111. Cami-Kobeci G, Williams MJ (2004) *Chem Commun* 1072–1073
112. Blank B, Michlik S, Kempe R (2009) *Chem Eur J* 15:3790–3799
113. Guillena G, Ramón DJ, Yus M (2007) *Angew Chem Int Ed* 46:2358–2364

114. Taguchi K, Nakagawa H, Hirabayashi T, Sakaguchi S, Ishii Y (2004) *J Am Chem Soc* 126:72–73
115. Onodera G, Nishibayashi Y, Uemura S (2006) *Angew Chem Int Ed* 43:3819–3822
116. Fujita K-I, Asai C, Yamaguchi T, Hanasaka F, Yamaguchi R (2005) *Org Lett* 7:4017–4019
117. Gnanamgari D, Leung CH, Schley ND, Hilton ST, Crabtree RH (2008) *Org Biomol Chem* 6:4442–4445
118. Matsu-ura T, Sakaguchi S, Obora Y, Ishii Y (2006) *J Org Chem* 71:8306–8308
119. Black PJ, Cami-Kobeci G, Edwards MG, Slatford PA, Whittlesey MK, Williams MJJ (2006) *Org Biomol Chem* 4:116–125
120. Morita M, Obora Y, Ishii Y (2007) *Chem Commun* 2850–2852
121. Löfberg C, Grigg R, Keep A, Derrick A, Sridharan V, Kilner C (2006) *Chem Commun* 5000–5002
122. Grigg R, Löfberg C, Whitney S, Sridharan V, Keep A, Derrick A (2009) *Tetrahedron* 65:849–854
123. Löfberg C, Grigg R, Whittaker MA, Keep A, Derrick A (2006) *J Org Chem* 71:8023–8027
124. Grigg R, Whitney S, Sridharan V, Keep A, Derrick A (2009) *Tetrahedron* 65:7468–7473
125. Grigg R, Whitney S, Sridharan V, Keep A, Derrick A (2009) *Tetrahedron* 65:4375–4383
126. Blank B, Kempe R (2010) *J Am Chem Soc* 132:924–925
127. Whitney S, Grigg R, Derrick A, Keep A (2007) *Org Lett* 9:3299–3302
128. Iuchi Y, Obora Y, Ishii Y (2010) *J Am Chem Soc* 132:2536–2537
129. Edwards MG, Williams MJJ (2002) *Angew Chem Int Ed* 41:4740–4743
130. Black PJ, Edwards MG, Williams MJJ (2006) *Eur J Org Chem* 4367–4378
131. Shermer DJ, Slatford PA, Edney DD, Williams MJJ (2007) *Tetrahedron Asymmetry* 18:2845–2848

# Formation of C–C Bonds via Iridium-Catalyzed Hydrogenation and Transfer Hydrogenation

John F. Bower and Michael J. Krische

**Abstract** The formation of C–C bonds via catalytic hydrogenation and transfer hydrogenation enables carbonyl and imine addition in the absence of stoichiometric organometallic reagents. In this review, iridium-catalyzed C–C bond-forming hydrogenations and transfer hydrogenations are surveyed. These processes encompass selective, atom-economic methods for the vinylation and allylation of carbonyl compounds and imines. Notably, under transfer hydrogenation conditions, alcohol dehydrogenation drives reductive generation of organoiridium nucleophiles, enabling carbonyl addition from the aldehyde or alcohol oxidation level. In the latter case, hydrogen exchange between alcohols and  $\pi$ -unsaturated reactants generates electrophile–nucleophile pairs en route to products of hydro-hydroxyalkylation, representing a direct method for the functionalization of carbinol C–H bonds.

**Keywords** Iridium, Enantioselective, Hydrogenation, Transfer hydrogenation, Allylation, Vinylation

## Contents

1	Introduction .....	108
2	Iridium-Catalyzed Hydrogenation for C–C Bond Formation .....	110
2.1	Alkynes as Vinylmetal Surrogates via Iridium-Catalyzed Hydrogenation .....	110
2.2	Allenes as Allylmetal Surrogates via Hydrogenation .....	115
3	Iridium-Catalyzed Transfer Hydrogenation for C–C Bond Formation .....	119
3.1	Allenes as Allylmetal Surrogates via Transfer Hydrogenation .....	119
3.2	Dienes as Allylmetal Surrogates via Transfer Hydrogenation .....	121

---

J.F. Bower

Department of Chemistry and Biochemistry, University of Texas at Austin, 1 University Station – A5300, Austin, TX 78712-0165, USA

and

Chemistry Research Laboratory, University of Oxford, 12 Mansfield Road, Oxford 1 3TA, UK

M.J. Krische (✉)

Department of Chemistry and Biochemistry, University of Texas at Austin, 1 University Station – A5300, Austin, TX 78712-0165, USA

e-mail: mkrische@mail.utexas.edu



3.3 Alkynes as Allylmetal Surrogates via Transfer Hydrogenation .....	123
3.4 Allylic Acetates as Allylmetal Surrogates via Transfer Hydrogenation .....	124
4 Conclusion .....	130
References .....	131

## 1 Introduction

The majority of chemical commodities (plastics, foams, pharmaceuticals, agrochemicals) are made from rapidly depleting petrochemical feedstocks. Consequently, there is a growing need to develop catalytic methods for the direct manufacture of chemical products from abundant renewable resources in the absence of stoichiometric byproducts [1]. In the specific case of carbonyl and imine addition, a departure from the use of premetallated nucleophiles represents a particularly important focus. Progress in this area will depend largely upon the discovery of new chemical reactivity.

Catalytic hydrogenation is the cleanest and most cost-effective method available for the reduction of organic molecules. Accordingly, chemical processes such as the Haber–Bosch reaction<sup>1</sup> [2, 3], alkene hydroformylation<sup>2</sup> [4–7], and enantioselective hydrogenation<sup>3</sup> [8–10], all of which provide materials essential to humanity, are practiced on a vast scale. The industrialization of these processes is assisted by the fact that such hydrogen-mediated transformations do not generate stoichiometric byproducts, which facilitate product purification and waste stream management. Interestingly, although catalytic hydrogenation and the Grignard reaction are both reductive transformations, a union between catalytic hydrogenation and carbonyl addition chemistry failed to materialize after the emergence of alkene hydroformylation, which may be viewed as the prototypical C–C bond-forming hydrogenation.

The collective studies from our laboratory establish metal-catalyzed hydrogenation and transfer hydrogenation as general methods for C–C bond formation.<sup>4,5,6</sup>

---

<sup>1</sup>The catalytic hydrogenation of atmospheric nitrogen, accounts for the annual production of over 100,000,000 metric tons of ammonia, which is the limiting nutrient in terrestrial plant growth. The Haber–Bosch process is estimated to sustain one-third of the Earth’s population. Approximately half the nitrogen in our bodies is nitrogen fixed through the Haber–Bosch reaction.

<sup>2</sup>The prototypical C–C bond forming hydrogenation, hydroformylation combines basic feedstocks ( $\alpha$ -olefins, carbon monoxide, and hydrogen) with perfect atom economy and accounts for the production of over 10 million metric tons of aldehyde annually, making it the largest volume application of homogeneous metal catalysis.


<sup>3</sup>The asymmetric hydrogenation of C=X  $\pi$ -bonds (X = O, NR) is estimated to account for over half the chiral drugs manufactured industrially, withstanding physical and enzymatic resolution.

<sup>4</sup>For selected reviews on C–C bond forming hydrogenation and transfer hydrogenation, see [11–17].

<sup>5</sup>Prior to our systematic studies, two isolated reports of hydrogen mediated C–C coupling were reported, see [18, 19].

<sup>6</sup>Side products of reductive C–C bond formation have been observed in catalytic hydrogenation on rare occasions, see [20, 21].

**Scheme 1** Catalytic C–C coupling via hydrogenation and transfer hydrogenation (X=O, NR<sub>2</sub>)



These processes exploit  $\pi$ -unsaturated reactants as surrogates to moisture sensitive organometallic reagents, enabling carbonyl and imine addition under essentially neutral conditions in the absence of stoichiometric organometallic reagents and resulting metallic byproducts. Remarkably, under transfer hydrogenation conditions, hydrogen exchange between alcohols and  $\pi$ -unsaturated reactants generates electrophile–nucleophile pairs en route to products of hydro-hydroxyalkylation, representing a direct method for the functionalization of carbinol C–H bonds.<sup>7,8,9</sup> For such “carbonyl additions from the alcohol oxidation level,” preactivation is not required as both the nucleophile and electrophile are generated catalytically from simple and stable precursors (Scheme 1). In this review, iridium-catalyzed C–C bond-forming hydrogenations and transfer hydrogenations are surveyed. Related hydrogen-mediated C–C couplings employing rhodium and ruthenium catalysts

<sup>7</sup>The alcohol-unsaturate couplings developed in our laboratory provide products of carbonyl addition. In contrast, related hydrogen auto-transfer processes provide products of alcohol substitution via pathways involving oxidation–condensation–reduction and the use of preactivated nucleophiles. For recent reviews, see [22–25].

<sup>8</sup>Processes that enable direct catalytic C–C functionalization of carbinol C–H bonds are highly uncommon. Rh-catalyzed alcohol-vinylarene C–C coupling has been described. The requirement of BF<sub>3</sub> and trends in substrate scope suggest these processes involve alcohol dehydrogenation-reductive Prins addition: [26–29].

<sup>9</sup>For radical mediated C–C functionalization of carbinol C–H bonds, see [30, 31].

are not covered in this account but are described in several recent monographs (see footnote 4, [11–17]).

## 2 Iridium-Catalyzed Hydrogenation for C–C Bond Formation

### 2.1 Alkynes as Vinylmetal Surrogates via Iridium-Catalyzed Hydrogenation

Allylic alcohols and allylic amines are highly versatile building blocks for organic synthesis, and methods for their preparation in enantiomerically enriched form are the subject of intensive investigation.<sup>10,11</sup> The majority of established protocols rely upon metal-catalyzed allylic substitution employing *O*- and *N*-nucleophiles (For reviews on metal catalyzed allylic amination and alkoxylation, see: [37–43]). An alternative approach to allylic alcohols and allylic amines involves the catalytic enantioselective vinylation of aldehydes and imines. Based upon the seminal studies of Oguni [44] and Noyori [45] on the enantioselective addition of dialkylzinc reagents to aldehydes (For reviews encompassing catalytic enantioselective addition of organozinc reagents to carbonyl compounds, see: [46–51]), Oppolzer (see footnote 12, [52]) devised the first catalytic enantioselective aldehyde vinylations.<sup>12,13</sup> In this process, vinylzinc reagents are generated *in situ* by alkyne hydroboration followed by transmetallation from boron to zinc using ZnMe<sub>2</sub>. A conceptually related approach was developed subsequently by Wipf, in which hydrozirconation-transmetallation is used to generate the vinylzinc reagent (see footnote 12, [55]). These studies, along with catalytic enantioselective alkyl- and aryl-zinc-mediated ketone additions described by Yus [73, 74] and Fu [75], set the stage for catalytic enantioselective ketone vinylations, as described by Walsh (For catalytic enantioselective ketone vinylation using organozinc reagents, see: [76–78]). Catalytic methods for carbonyl vinylation continue to emerge, as demonstrated by recent reports on rhodium-catalyzed aldehyde additions employing vinylboron reagents [79].

Enantioselective vinylzinc-mediated imine additions reveal additional challenges (For reviews encompassing the catalytic enantioselective addition of organozinc reagents to imines, see: [80–84]). After the seminal report of Soai [85], several enantioselective catalysts for the addition of organozinc reagents to imines were developed. However, highly activated *N*-acyl and *N*-(diphenylphosphinoyl) imines are required (For enantioselective catalytic addition of organozinc reagents

---

<sup>10</sup>For reviews encompassing the synthesis of allylic alcohols, see [32, 33].

<sup>11</sup>For reviews encompassing the synthesis of allylic amines, see [34–36].

<sup>12</sup>For enantioselective catalytic addition of vinylzinc reagents to aldehydes, see [52–70].

<sup>13</sup>For reviews encompassing catalytic enantioselective aldehyde vinylation using organozinc reagents, see [71, 72].

to imines, see: [85–96]). To address the issue of reactivity, Tomioka (see footnote 14, [97]) disclosed an enantioselective copper catalyst for organozinc-mediated imine addition.<sup>14</sup> In addition to late transition metals (Cu, Rh) (see footnote 14) [117, 118], early transition metal catalysts (Ti, Zr, Hf) (For enantioselective zirconium, titanium and hafnium catalyzed addition of organozinc reagents to imines, see: [119–124]) are found to promote the highly enantioselective addition of organozinc reagents to imines. The enantioselective addition of organolithium reagents to imines promoted or catalyzed by chiral chelating agents also has been described (For reviews encompassing catalytic enantioselective addition of organolithium reagents to carbonyl compounds and imines, see: [125–132]). Finally, under the conditions of rhodium catalysis, organotin, organotitanium, and organoboron reagents participate in catalytic asymmetric imine addition (For enantioselective rhodium catalyzed addition of organometallic reagents to imines, see: Organotin reagents [133–134], Organotitanium reagents [135] and Organoboron reagents [136]). Despite these efforts, highly enantioselective imine additions employing vinylmetal reagents remain elusive (Catalyzed addition of vinylzirconocenes to imines is known, but enantioselective variants have not been developed: [147, 148]).<sup>15</sup>


The direct metal-catalyzed reductive coupling of alkynes to carbonyl compounds or imines provides an alternative to the use of discrete vinylmetal reagents (For a compilation of reviews on the subject of metal catalyzed reductive C–C coupling, see: [150]).<sup>16</sup> Catalytic intramolecular carbonyl vinylation was first reported by Ojima (1994), who employed a rhodium catalyst in combination with a silane as the terminal reductant [159]. Related cyclizations using titanium or nickel catalysts later were reported by Crowe [160] (For an aligned study, see: [161]) and Montgomery [162–167], respectively. Finally, in an adaptation of Montgomery's cyclization protocol, intermolecular nickel-catalyzed alkyne–aldehyde reductive couplings were reported by Jamison [168–171]. Subsequently, improved nickel catalysts for this process were devised by Takai [172] and Montgomery [165]. However, while direct alkyne–carbonyl reductive couplings circumvent the use of discrete vinylmetal reagents, they nevertheless employ terminal reductants such as hydrosilanes, hydrostannanes, organozinc reagents, organoboron reagents or chromium(II) chloride, which generate stoichiometric quantities of byproduct.

Under the conditions of iridium-catalyzed hydrogenation, alkyne–carbonyl and alkyne–imine reductive coupling occurs in the absence of stoichiometric byproducts. For example, iridium-catalyzed hydrogenation of nonconjugated alkynes in the presence of  $\alpha$ -ketoesters delivers the corresponding  $\alpha$ -hydroxy esters in


<sup>14</sup>For enantioselective copper catalyzed addition of organozinc reagents to imines, see [97–116].

<sup>15</sup>Enantioselective Ni-catalyzed alkyne, imine, triethylborane 3-component coupling has been reported, but modest selectivities (51–89% ee) are observed. For this method, vinylation is accompanied by ethyl transfer: [149]

<sup>16</sup>For selected reviews encompassing intra- and intermolecular direct reductive coupling of alkynes to carbonyl partners, see [151–158]



**Scheme 2** Iridium-catalyzed hydrogenation of alkynes in the presence of α-ketoesters to furnish α-hydroxy esters





**Scheme 3** Alkyne-carbonyl reductive coupling via iridium-catalyzed hydrogenation under a deuterium atmosphere

excellent yield [173]. Notably, nonsymmetric alkynes engage in highly regioselective reductive coupling (Scheme 2).

Deuterium labeling studies are consistent with a mechanism involving alkyne-carbonyl oxidative coupling to furnish iridacycle **A** followed by Brønsted acid-assisted hydrogenolysis of this species to deliver the product. Deuterium at oxygen is removed through exchange in the course of chromatographic isolation but is observed via EI-MS analysis of reaction mixtures. Although mechanisms involving alkyne hydrometallation cannot be excluded on the basis of deuterium labeling studies alone, regioselectivity trends observed in response to variation of the steric features of both the alkyne and carbonyl partners strongly suggest that oxidative coupling pathways are operative (Scheme 3).


Cationic rhodium complexes also catalyze alkyne-carbonyl and alkyne-imine reductive coupling under hydrogenation conditions [174–178]. The rhodium-catalyzed processes are applicable to the reductive coupling of conjugated alkynes (1,3-enynes, 1,3-diyne) to activated ketones [174], aldehydes [175–177], or imines [178]. The rhodium-catalyzed reductive coupling of acetylene to carbonyl compounds or imines also is described [179–182]. For reactions of conjugated alkynes, the observed trends in substrate scope suggest π-backbonding at the stage of the

**Scheme 4** Significance of  $\pi$ -backbonding at the stage of the alkyne–metal complex**Fig. 1** Carboxylic acid-assisted hydrogenolysis of oxa-iridacyclic intermediates that arise via alkyne–carbonyl oxidative coupling

metal–alkyne complex facilitates alkyne–C=X (X=O, NR) oxidative coupling [183–185]. As rhodium is a relatively weak  $\pi$ -donor, conjugated alkynes and activated carbonyl and imine partners, which embody lower lying LUMOs, are required for oxidative coupling. Due to relativistic effects, iridium is a stronger  $\pi$ -donor than rhodium [186–189] and, therefore, can promote oxidative coupling of nonconjugated alkynes, which have higher lying LUMOs (Scheme 4).<sup>17</sup>

Regarding the role of the Brønsted acid cocatalyst, computational studies by Musashi and Sakaki suggest that carboxylic acid additives accelerate coupling by circumventing highly energetic four-centered transition structures for hydrogenolysis of metallacyclic intermediates **A** [195]. In the presence of carboxylic acid, protonolysis of oxa-iridacyclic intermediates **A** provide iridium carboxylates, which engage in hydrogenolysis through lower energy six-centered transition structures **C**. Protonolysis itself may occur through six-centered transition structures **B**. This interpretation is consistent with recently reported ESI–MS and DFT studies on the rhodium-catalyzed reductive coupling of acetylene to carbonyl and imine partners under hydrogenation conditions (Fig. 1) [182].

<sup>17</sup>Alkyne complexation by iridium(I) results in substantial deviation from linearity, as revealed by single crystal X-ray diffraction analysis: [190–194].




**Scheme 5** Enantioselective iridium-catalyzed hydrogenation of alkynes in the presence of *N*-arylsulfonyl imines to furnish trisubstituted allylic amines

Cationic iridium complexes modified by chiral phosphine ligands promote highly enantioselective alkyne–imine reductive coupling under hydrogenation conditions to furnish trisubstituted allylic amines as single geometrical isomers [196, 197]. Carboxylic acid cocatalysts are required to enforce high levels of conversion. Additionally, Na<sub>2</sub>SO<sub>4</sub> facilitates coupling by serving as a desiccant, thus mitigating imine hydrolysis by adventitious water. A remarkable feature of these transformations involves high levels of contrastric regioselectivity observed in reductive couplings of nonsymmetric alkynes (Scheme 5).

As corroborated by deuterium labeling, the observed contrastric regiocontrol appears to be a consequence of the oxidative coupling pathway. Specifically, formation of aza-iridacycle **A** via alkyne–imine oxidative coupling occurs such that nonbonded interactions between the large alkyne substituent and the sterically demanding iridium center are minimized. Carboxylic acid-assisted hydrogenolysis (via **B** and **C**) delivers the allylic amine. Formation of the iridacyclic intermediate **A** is the enantiodetermining event. Here, square planar iridium(I) binds the alkyne and imine at adjacent coordination sites. In accordance with the Dewar–Chatt–Duncanson model [183–185], the bound alkyne exhibits iridacyclopropene character (Alkyne complexation by iridium(I) results in substantial deviation from linearity, as revealed by single crystal X-ray diffraction analysis: (see footnote 17) [190–194]). Bidentate coordination of the *N*-arylsulfonyl imine to the iridium center is anticipated<sup>18</sup> [198]. Insertion of the imine into the iridium–carbon bond of the iridacyclopropene then delivers the indicated aza-iridacyclopentene **A**. Minimization of nonbonded interactions between the *N*-arylsulfonyl group and the phenylphosphino moieties accounts for the observed sense of stereoinduction (Scheme 6).

<sup>18</sup>The bidentate κ<sub>2</sub>-mode of binding has been observed by single crystal X-ray diffraction analysis for a related palladium *N*-arylsulfonamidate complex.




**Scheme 6** Proposed mechanism for hydrogenative imine vinylation and proposed enantiodetermining transition states

## 2.2 Allenes as Allylmetal Surrogates via Hydrogenation

Enantioselective carbonyl allylation is one of the most broadly utilized transformations in synthetic organic chemistry (For reviews on enantioselective carbonyl allylation, see: [199–204]). Shortly after the first reports of carbonyl allylation employing isolable allylboron and allylsilicon reagents by Mikhailov and Bubnov [205] and Hosomi and Sakurai [206], respectively, enantioselective carbonyl allylations were reported by Hoffmann (see footnote 19, [207]). While the design of increasingly effective chiral allylmetal reagents continues<sup>19</sup> [207–220], this

<sup>19</sup>For selected examples of chirally modified allyl metal reagents, see [207–220].






**Fig. 2** Selected examples of chiral allylmetal reagents for enantioselective carbonyl allylation

approach only can be viewed as an interim solution as the use of stoichiometric chiral-inducing elements invariably leads to molar quantities of byproduct, diminishing prospects for large-volume applications (Fig. 2).

This limitation is addressed, in part, by the groundbreaking work of Yamamoto [221] and subsequent studies by Umani-Ronchi [222] and Keck [223] on chiral Lewis acid catalysts for enantioselective carbonyl allylation. Chiral Lewis base-catalyzed enantioselective carbonyl allylations soon followed, as reported by Denmark [224].<sup>20</sup> Although these methods avoid the stoichiometric use of chiral modifiers, they employ molar quantities of allylmetal reagent and, consequently, generate stoichiometric quantities of metallic byproducts. Furthermore, preparation of the allylmetal reagent itself is often beset by excessive waste production. For example, allylstannanes employed in the Umani-Ronchi–Keck process are prepared from the corresponding organomagnesium reagents, which, in turn, are prepared from the allylic halides [222, 223]. Thus, preactivation in the form of three stoichiometric operations is required in advance of C–C coupling (Scheme 7).

<sup>20</sup>For selected examples of catalytic asymmetric carbonyl allylation employing allylmetal reagents, see [221–224].



**Scheme 7** Preactivation often accompanies the use of stoichiometric organometallic reagents

The reductive generation of transient allylmetal species from allylic alcohols, allylic carboxylates,<sup>21,22</sup> or allylic halides<sup>23,24</sup> represents an approach to carbonyl allylation that avoids discrete allylmetal reagents. However, with a few exceptions, (see footnote 21, [235–238]) such processes require stoichiometric quantities of metallic reductants, such as  $\text{SmI}_2$ ,  $\text{SnCl}_2$ , and  $\text{Et}_2\text{Zn}$ . Finally, the carbonyl–ene reaction enables carbonyl allylation in the absence of stoichiometric byproducts by employing simple olefins as allyl donors; however, enantioselective variants are limited to highly activated electrophiles, and general protocols remain a distant goal.<sup>25,26</sup>

Carbonyl allylation in the absence of metallic reagents is achieved under the conditions of iridium-catalyzed hydrogenation or transfer hydrogenation employing 1,2-dienes, 1,3-dienes, or allyl acetates as precursors to transient allylmetal species (see footnote 4, [16]). Here, hydrogen or isopropanol may serve as reductant. For transfer hydrogenative processes, isopropanol dehydrogenation drives reductive generation of allylmetal nucleophiles from unsaturates. Of greater significance is the fact that hydrogen exchange between reactant alcohols and  $\pi$ -unsaturated partners occurs to generate electrophile–nucleophile pairs en route to products of C–C coupling. In this way, carbonyl allylation is achieved directly from the alcohol oxidation level, availing a method for the hydro-hydroxyalkylation of unsaturates via functionalization of carbinol C–H bonds (Scheme 8) (see footnotes (7, 8, 9), [22–31]).

Hydrogen-mediated carbonyl allylation was first demonstrated under the conditions of iridium-catalyzed hydrogenation in studies of allene–carbonyl reductive coupling [269]. Using a cationic iridium catalyst, hydrogenation of commercially available 1,1-dimethylallene in the presence of carbonyl electrophiles delivers products of carbonyl *tert*-prenylation as single regioisomers in good to excellent

<sup>21</sup>For selected examples of carbonyl allylations employing nucleophilic  $\pi$ -allyls derived from allylic acetates and carboxylates, see: *Palladium* [225–233], *Rhodium* [234, 235], *Ruthenium* [236–238], *Iridium* [239–241].


<sup>22</sup>For selected reviews on carbonyl allylation via umpolung of  $\pi$ -allyls, see [242–247].

<sup>23</sup>For catalytic enantioselective carbonyl allylation and crotylation via Nozaki–Hiyama coupling, see [248–257].


<sup>24</sup>For recent reviews of catalytic Nozaki–Hiyama coupling, see [258–261].

<sup>25</sup>For reviews on carbonyl–ene reactions, see [262–265].

<sup>26</sup>For Nickel catalyzed carbonyl–ene reactions, see [266–268].




**Scheme 8** Carbonyl allylation via hydrogenation and transfer hydrogenation



**Scheme 9** Catalytic carbonyl addition via iridium-catalyzed hydrogenative coupling of dimethylallene

yield. Note that functional groups typically considered “hydrogen-labile,” such as aryl halides, benzylic ethers, alkenes, and nitroarenes, are tolerated under these conditions. Unlike conventional allylation protocols, all reactant atoms, including hydrogen, are incorporated into the product and so no stoichiometric byproducts are generated (Scheme 9).

Reductive coupling of 1,1-dimethylallene and 5-nitro-2-furancarboxaldehyde under a deuterium atmosphere provides the product of *tert*-prenylation incorporating deuterium at the interior vinylic position (80%  $^2\text{H}$ ). This result is consistent with a mechanism involving allene–aldehyde oxidative coupling. However, alternate pathways involving allene hydrometallation to furnish allyliridium species cannot be excluded on the basis of these data (Scheme 10).



**Scheme 10** Iridium-catalyzed reductive coupling of 1,1-dimethylallene to an aldehyde under an atmosphere of deuterium


### 3 Iridium-Catalyzed Transfer Hydrogenation for C–C Bond Formation

#### 3.1 Allenes as Allylmetal Surrogates via Transfer Hydrogenation


Although carbonyl *tert*-prenylation occurs efficiently upon iridium-catalyzed hydrogenation of 1,1-dimethylallene in the presence of aldehydes, attempted crotylations and allylations employing methylallene or allene gas, respectively, suffer from competing conventional hydrogenation of the olefinic C–C coupling product. Corresponding iridium-catalyzed transfer hydrogenations overcome this limitation [270]. Under nearly identical conditions, by simply substituting isopropanol (200 mol%) for hydrogen as the terminal reductant, 1,1-dimethylallene reductively couples to aldehydes to furnish products of *tert*-prenylation in good to excellent yield across a broad range of substrates. Remarkably, it was found that alcohol reactants serve dually as reductants and aldehyde precursors, enabling formation of identical products. Thus, carbonyl addition occurs with equal facility from the aldehyde or alcohol oxidation levels. This chemistry is also applicable to carbonyl crotylation and allylation employing methylallene or allene gas, respectively. However, the volatility of these reagents contributes to diminished efficiencies (Scheme 11).

The coupling of 1,1-dimethylallene to benzaldehyde employing  $d_3$ -isopropanol as reductant delivers the product of *tert*-prenylation, incorporating deuterium primarily at the internal vinylic position (85%  $^2\text{H}$ ). Similar results are obtained when the reaction is conducted from the alcohol oxidation level employing  $d_2$ -benzyl alcohol. Further mechanistic insight comes from crossover experiments in which 1,1-dimethylallene is exposed to equimolar quantities of *p*-nitrobenzyl alcohol and benzaldehyde. The *p*-nitrophenyl- and phenyl-containing adducts are obtained in a 4:1 ratio, respectively. An identical product distribution is obtained upon exposure of 1,1-dimethylallene to equimolar quantities of *p*-nitrobenzaldehyde and benzyl alcohol under otherwise identical conditions. These data are consistent with fast and reversible alcohol dehydrogenation to form nonmetal-bound aldehyde in advance of C–C coupling (Scheme 12).

Highly enantioselective variants of these processes employ a cyclometallated iridium *C,O*-benzoate derived from allyl acetate, *m*-nitrobenzoic acid, and




**Scheme 11** Carbonyl *tert*-prenylation, crotylation, and allylation from the aldehyde or alcohol oxidation level under the conditions or iridium-catalyzed transfer hydrogenation




**Scheme 12** Isotopic labeling and crossover experiments in iridium-catalyzed couplings of 1,1-dimethylallene under transfer hydrogenation conditions

(*S*)-SEGPHOS [271, 272]. Using this complex as a precatalyst, transfer hydrogenation of 1,1-dimethylallene in the presence of diverse aldehydes mediated by isopropanol delivers products of *tert*-prenylation in good to excellent yield and with excellent levels of enantioselectivity. In the absence of isopropanol, enantioselective carbonyl reverse prenylation is achieved directly from the alcohol oxidation level to furnish an equivalent set of adducts. Notably, enantioselective *tert*-prenylation is achieved under mild conditions (30–50°C) in the absence of stoichiometric metallic reagents. Indeed, for reactions conducted from the alcohol oxidation level, stoichiometric byproducts are completely absent (Scheme 13).



**Scheme 13** Enantioselective carbonyl *tert*-prenylation from the alcohol or aldehyde oxidation level via iridium-catalyzed C–C bond-forming transfer hydrogenation



**Scheme 14** Proposed catalytic mechanism for carbonyl *tert*-prenylation from the alcohol oxidation level via iridium-catalyzed C–C bond-forming transfer hydrogenation


Our collective mechanistic studies are consistent with the indicated catalytic cycle. Notably, the catalyst engages primary alcohols in rapid and reversible dehydrogenation, yet the coupling products, which are homoallylic alcohols, are not subject to oxidation as coordination of the homoallylic olefin to the catalyst provides a hexa-coordinate 18-electron complex lacking an open coordination site for  $\beta$ -hydride elimination (Scheme 14).

### 3.2 Dienes as Allylmetal Surrogates via Transfer Hydrogenation


Under the conditions of iridium-catalyzed transfer hydrogenation employing isopropanol as reductant, 1,3-cyclohexadiene couples to aryl aldehydes to provide

products of carbonyl cyclohexenylation in good to excellent yield and with high levels of diastereocontrol. Under nearly identical conditions, but in the absence of isopropanol, corresponding alcohols couple directly to 1,3-cyclohexadiene to furnish identical products of carbonyl cyclohexenylation (Scheme 15) [273].

More recently, using the cyclometallated iridium *C,O*-benzoate derived from allyl acetate, 4-methoxy-3-nitrobenzoic acid and BIPHEP, catalytic carbonyl crotylation employing 1,3-butadiene from the aldehyde, or alcohol oxidation was achieved under transfer hydrogenation conditions [274]. Carbonyl addition occurs with roughly equal facility from the alcohol or aldehyde oxidation level. However, products are obtained as diastereomeric mixtures. Stereoselective variants of these processes are under development. It should be noted that under the conditions of ruthenium-catalyzed transfer hydrogenation, conjugated dienes, including butadiene, couple to alcohols or aldehydes to provide either products of carbonyl crotylation or  $\beta,\gamma$ -enones (Scheme 16) [275, 276].



**Scheme 15** Iridium-catalyzed couplings of 1,3-cyclohexadiene under transfer hydrogenation conditions (ratio refers to 1,4-olefinic versus 1,5-olefinic alcohols)




**Scheme 16** Iridium-catalyzed C–C couplings of 1,3-butadiene under transfer hydrogenation conditions


### 3.3 Alkynes as Allylmetal Surrogates via Transfer Hydrogenation

As demonstrated in recent work by Obora and Ishii, alkynes serve as allyl donors in carbonyl allylations from the alcohol oxidation level [277]. Specifically, upon exposure to an iridium catalyst generated in situ from  $[\text{Ir}(\text{OH})(\text{cod})]_2$  and  $\text{P}(n\text{-Oct})_3$ , 1-aryl-2-methylalkynes couple to primary alcohols to furnish homoallylic alcohols with complete branched regioselectivity and excellent levels of diastereoselectivity (Scheme 17).

For alkyne-mediated allylation, alcohol dehydrogenation generates an iridium hydride that hydrometallates the alkyne to form a vinylmetal species, which, in turn, isomerizes to generate the primary  $\sigma$ -allyliridium intermediate. Carbonyl addition through a closed six-centered transition state accounts for the observed *anti*-diastereoselectivity (Scheme 18). Interestingly, under the conditions of ruthenium-catalyzed transfer hydrogenation, alkynes couple to alcohols or aldehydes to provide products of carbonyl vinylation [278]. Here, isomerization of the vinyl-ruthenium intermediate to the  $\sigma$ -allylruthenium intermediate is not observed.



**Scheme 17** Carbonyl aryallylation from the alcohol oxidation level via iridium-catalyzed transfer hydrogenation employing alkynes as allyl donors



**Scheme 18** Plausible stereochemical model for *anti*-diastereoselective iridium-catalyzed allylation of alcohols employing alkynes as allyl donors




### 3.4 Allylic Acetates as Allylmetal Surrogates via Transfer Hydrogenation

Using allyl acetate as the allyl donor, enantioselective variants of the parent carbonyl allylation process are achieved under the conditions of iridium-catalyzed transfer hydrogenation employing the cyclometallated iridium *C,O*-benzoate derived from allyl acetate, 3-nitrobenzoic acid, and a chiral *bis*-phosphine ligand, which is generated in situ [279, 280]. Allyl acetate couples to diverse alcohols to furnish products of carbonyl allylation with exceptional levels of enantioselectivity. Identical products of carbonyl allylation are produced from the corresponding aldehydes upon use of isopropanol as terminal reductant. Thus, enantioselective carbonyl allylation is achieved from the alcohol or aldehyde oxidation level in the absence of metallic reagents or metallic reductants. The reactivity embodied by these processes is remarkable in light of the fact that iridium-catalyzed allylic substitution (*O*-allylation) employing alcohol nucleophiles is a well-established mode of reactivity (Scheme 19) [281–283].<sup>27</sup>


The results of deuterium labeling are consistent with intervention of a symmetric iridium  $\pi$ -allyl intermediate or rapid interconversion of  $\sigma$ -allyl haptomers through the agency of a symmetric  $\pi$ -allyl (Scheme 20) [280]. Competition experiments akin to those previously described (see Scheme 12) again demonstrate rapid and reversible dehydrogenation of the carbonyl partner in advance of C–C coupling.

The BINAP derivative of the *ortho*-cyclometallated iridium catalyst has been characterized by single crystal X-ray diffraction analysis [280]. Remarkably, although the reaction sequence depends upon oxidation of either the reactant alcohol or isopropanol, the enantiomeric purity of the homoallylic alcohol product




**Scheme 19** Enantioselective carbonyl allylation from the alcohol or aldehyde oxidation level via iridium-catalyzed C–C bond-forming transfer hydrogenation

<sup>27</sup>Under the conditions of ruthenium catalysis, alcohols and allylic acetates couple to form enones, see [284].




**Scheme 20** Carbonyl allylation from the alcohol oxidation level employing isotopically labeled allyl acetate [R = *p*-(CO<sub>2</sub>Me)C<sub>6</sub>H<sub>4</sub>]



**Scheme 21** Proposed catalytic mechanism for carbonyl allylation from the alcohol oxidation level via iridium-catalyzed C–C bond-forming transfer hydrogenation

is not subject to erosion through reversible dehydrogenation as the pendant alkene of the product prevents dehydrogenation by occupying the single remaining vacant coordination site on iridium, as required for  $\beta$ -hydride elimination (Scheme 21).

A model accounting for the observed sense of absolute stereoselection is based upon the coordination mode revealed in the crystal structure of the cyclometallated *C,O*-benzoate complex [280]. It is postulated that aldehyde binding by the  $\sigma$ -allyl haptomer occurs such that the allyl moiety is placed between the naphthyl and phenyl moieties of the ligand, allowing the aldehyde to reside in a more open environment. In the favored mode of addition, the aldehyde C–H bond projects into




**Fig. 3** Proposed stereochemical model accounting for the observed sense of absolute stereoinduction (chiral ligand = (*S*)-Cl,MeO-BIPHEP)

the  $\pi$ -face of a phenyl moiety of the ligand, potentially resulting in a weakly attractive aldehyde C–H– $\pi$  interaction. The disfavored mode of addition requires the aldehyde substituent to project into the  $\pi$ -face of the ligand phenyl moiety, resulting in a severe nonbonded interaction (Fig. 3).


Direct asymmetric carbonyl allylation from the alcohol oxidation level potentially streamlines synthesis by circumventing discrete alcohol oxidation as well as enabling carbonyl allylation processes that are unattainable using conventional allylmetal reagents. For example, two-directional chain elongation via double allylation of 1,3-dialdehydes has not been achieved due to the instability of the requisite 1,3-dicarbonyl precursors (For a review on two-directional chain synthesis, see: [285]). However, propane-1,3-diol and related glycols are highly tractable and participate in bidirectional allylation via sequential generation and capture of transient monoaldehydes to provide  $C_2$ -symmetric adducts in good yield [286]. In these processes, the minor enantiomer of the monoallylated intermediate is transformed to the *meso*-stereoisomer of the product, thus amplifying the level of enantiomeric enrichment (Scheme 22) (This mechanism for enantiomeric enrichment is documented by Eliel and Midland: [287, 288]).

Iterative two-directional chain elongation enables rapid construction of 1,3-polyols (Scheme 23). For example, the polyol substructure that appears in the oxopolyene macrolide (+)-roxaticin is available in nine steps from propane-1,3-diol upon three iterations of double carbonyl allylation [286]. Notably, high levels of catalyst-directed stereoselectivity are observed (For selected examples of catalyst directed diastereoselectivity, see: [289–296]). Analogous 1,3-polyol syntheses via iterative homologation in one direction also are reported [297].

This carbonyl allylation strategy is extended to crotylation through the use of  $\alpha$ -methyl allyl acetate as a crotylmetal surrogate [298]. Here, after screening several *ortho*-cyclometallated *C,O*-benzoates, the iridium complex derived from 4-cyano-3-nitrobenzoic acid and (*S*)-SEGPPOS [272] was found to provide good levels of *anti*-diastereoselectivity accompanied by uniformly high levels of enantioselectivity across a broad range of substrates. Notably, complete branch-regioselectivity is observed in all cases. The observed regio- and *anti*-diastereoselectivity suggest that




**Scheme 22** 1,*n*-Glycols as dialdehyde equivalents in iridium-catalyzed enantioselective carbonyl allylation from the alcohol oxidation level




**Scheme 23** Synthesis of the 1,3-polyol substructure of the polyene macrolide (+)-roxaticin

aldehyde addition occurs from the primary (*E*)- $\sigma$ -allyliridium haptomer. As before, carbonyl crotylation is achieved with equal facility from the aldehyde or alcohol oxidation level (Scheme 24).




**Scheme 24** Enantioselective carbonyl crotylation from the alcohol or aldehyde oxidation level via iridium-catalyzed C–C bond-forming transfer hydrogenation



**Scheme 25** Diastereo- and enantioselective *anti*-(hydroxy)allylation employing allylic *gem*-dicarboxylates as allyl donors via iridium-catalyzed transfer hydrogenation

Allylic *gem*-dibenzoates derived from acrolein serve as (alkoxy)allylmetal surrogates under the conditions of iridium-catalyzed transfer hydrogenation [299]. In this case, generation of the catalytic complex *in situ* failed to provide the desired products in good yield, requiring use of the isolated complex prepared from  $[\text{Ir}(\text{cod})\text{Cl}]_2$ , 4-cyano-3-nitrobenzoic acid, allyl acetate, and the chiral phosphine ligand (*R*)-SEGPHOS [272]. To achieve high levels of *anti*-diastereoselectivity, efficient partitioning of transient (*E*)- and (*Z*)- $\sigma$ -allyliridium intermediates is required. It was found that the allylic *gem*-dibenzoate derived from acrolein is unique in its ability to combine high levels of *anti*-diastereoselectivity and favorable reactivity. Under the reported conditions, iridium-catalyzed aldehyde *anti*-(hydroxy)allylation occurs under transfer hydrogenation conditions to provide *anti*-1-ene-3,4-diols with excellent relative and absolute stereocontrol. Corresponding carbonyl additions from the alcohol oxidation level are currently under development (Scheme 25).

New catalytic allylation methodologies continue to emerge. For example, iridium-catalyzed transfer hydrogenation of  $\alpha$ -(trimethylsilyl)allyl acetate in the presence of aldehydes mediated by isopropanol and employing the iridium catalyst




**Scheme 26** Enantioselective carbonyl (trimethylsilyl)allylation and (hydroxymethyl)allylation from the alcohol or aldehyde oxidation level via iridium-catalyzed C–C bond-forming transfer hydrogenation

modified by (*R*)-SEGPHOS [272] delivers products of (trimethylsilyl)allylation in good isolated yields and with exceptional levels of *anti*-diastereo and enantio-control [300]. In the absence of isopropanol, carbonyl (trimethylsilyl)allylation is achieved directly from the alcohol oxidation level to furnish an equivalent set of adducts (Scheme 26, top). Similarly, using the enantiomeric iridium complex, the indicated cyclic carbonate engages in carbonyl (hydroxymethyl)allylation from the alcohol or aldehyde oxidation level with good *anti*-diastereoselectivities and exceptional levels of enantiocontrol [301]. Notably, stereoselective (hydroxymethyl)allylation protocols employing conventional allylmetal reagents are unknown (Scheme 26, bottom).<sup>28,29,30</sup>

<sup>28</sup>For (hydroxymethyl)allylation via palladium catalyzed reductive coupling of allylic carboxylates, see [302–304].

<sup>29</sup>For (hydroxymethyl)allylation via palladium catalyzed reductive coupling of vinyl epoxides, see [305, 306].

<sup>30</sup>For catalytic enantioselective (hydroxymethyl)allylation, see [307, 308].




**Scheme 27** Enantioselective allylation, crotylation, and *tert*-prenylation of substituted isatins via iridium-catalyzed transfer hydrogenation

Enantioselective ketone addition is one of the many challenging extensions in scope that remains. In a preliminary step toward this goal, the allylation, crotylation, and *tert*-prenylation of substituted isatins was examined using the cyclometallated iridium *C,O*-benzoate generated in situ from [Ir(cod)Cl]<sub>2</sub>, 4-chloro-3-nitrobenzoic acid and CTH-(*R*)-P-PHOS [309]. Isopropanol-mediated transfer hydrogenation of allyl acetate,  $\alpha$ -methyl allyl acetate or 1,1,-dimethylallylene as allyl donors, in the presence of the substituted isatins delivered the anticipated products of allylation, crotylation, and *tert*-prenylation, respectively, in highly enantiomerically enriched form. Notably, highly enantioselective allylation, crotylation, and reverse prenylation of isatins employing conventional allylmetal reagents have not been reported (Scheme 27) (For enantioselective catalytic allylation of isatins, see: [310]).

## 4 Conclusion

Organic molecules are defined as compounds of carbon and hydrogen. Hence, the formation of C–C bonds under the conditions catalytic hydrogenation and transfer hydrogenation is a natural endpoint in the evolution of strategies for the synthesis of organic molecules. The use of unsaturates as surrogates to premetallated nucleophiles under hydrogenation conditions represents a departure from the use of stoichiometric organometallic reagents in carbonyl and imine addition that circumvents generation of stoichiometric byproducts. This concept is extended further via C–C bond-forming transfer hydrogenation – processes in which alcohol reactants

**Scheme 28** Examples of two aspirational processes: byproduct-free couplings of  $\alpha$ -olefins to (renewable) alcohols and amines with linear or branch regiocontrol



serve dually as hydrogen donors and carbonyl precursors, enabling carbonyl addition directly from the alcohol oxidation level in the absence of stoichiometric byproducts and premetallated nucleophiles.

As highlighted in this review, iridium complexes are especially versatile catalysts for C–C bond-forming hydrogenation and transfer hydrogenation. And now, based on these studies, many aspirational transformations are placed within reach (Scheme 28). However, to fully realize the potential of hydrogen-mediated C–C coupling, alternate metal catalysts should be developed and applied to processes beyond those described herein. This challenge will ultimately evoke catalytic methods for the sustainable manufacture of pharmaceutical ingredients, bulk chemicals, and fuels from renewable feedstocks.

**Acknowledgments** Acknowledgment is made to the Robert A. Welch Foundation (F-0038), NIH-NIGMS (RO1-GM69445), and the ACS-GCI Pharmaceutical Roundtable for partial support of the research described in this account. Dr Oliver Briel of Umicore is thanked for the generous donation of iridium salts. Dr Wataru Kuriyama and Dr Yasunori Ino of Takasago are thanked for the generous donation of (*R*)- and (*S*)-SEGPPOS.

## References

1. Board on Chemical Sciences and Technology, National Research Council of the National Academies (2005) Sustainability in the chemical industry: grand challenges and research needs. National Academies Press, Washington, DC
2. Nobel Foundation (1966) Nobel lectures, chemistry, 1901–1921. Elsevier, Amsterdam
3. Smil V (2004) Enriching the earth: Fritz Haber, Carl Bosch, and the transformation of world food production. MIT, Cambridge, MA
4. Roelen O (1938) Chemische Verwertungsgesellschaft mbH, Oberhausen. German Patent DE 849,548
5. Roelen O (1944) Chemische Verwertungsgesellschaft mbH. Chem Abstr 38:5501
6. Frohning CD, Kohlpaintner CW (1996) In: Cornils B, Herrmann WA (eds) Applied homogeneous catalysis with organometallic compounds, vol 1. VCH, Weinheim, p 29
7. van Leeuwen PWNM (2004) Homogeneous catalysis, understanding the art. Kluwer, Dordrecht
8. Thommen M (2005) Spec Chem Mag 25(4):26
9. Thayer A (2005) Chem Eng News 83(36):40
10. Jäkel C, Paciello R (2006) Chem Rev 106:2912



11. Ngai M-Y, Kong JR, Krische MJ (2007) *J Org Chem* 72:1063
12. Skucas E, Ngai M-Y, Komanduri V, Krische MJ (2007) *Acc Chem Res* 40:1394
13. Iida H, Krische MJ (2007) *Top Curr Chem* 279:77
14. Shibahara F, Krische MJ (2008) *Chem Lett* 37:1102
15. Bower JF, Kim IS, Patman RL, Krische MJ (2009) *Angew Chem Int Ed* 48:34
16. Han SB, Kim IS, Krische MJ (2009) *Chem Commun* 7278
17. Bower JF, Krische MJ (2009) In: Anastas P (ed) *Handbook of green chemistry*, vol 1. VCH, Weinheim, p 205
18. Molander GA, Hoberg JO (1992) *J Am Chem Soc* 114:3123
19. Kokubo K, Miura M, Nomura M (1995) *Organometallics* 14:4521
20. Moyes RB, Walker DW, Wells PB, Whan DA, Irvine EA (1992) In: Dines TJ, Rochester CH, Thomson J (eds) *Catalysis and surface characterisation (special publication)*, vol 114. Royal Society of Chemistry, London, p 207
21. Bianchini C, Meli A, Peruzzini M, Vizza F, Zanolini F, Frediani P (1989) *Organometallics* 8:2080
22. Guillena G, Ramón DJ, Yus M (2007) *Angew Chem Int Ed* 46:2358
23. Hamid MHSA, Slatford PA, Williams MJ (2007) *Adv Synth Catal* 349:1555
24. Nixon TD, Whittlesey MK, Williams MJ (2009) *Dalton Trans* 753
25. Guillena G, Ramón DJ, Yus M (2010) *Chem Rev* 110:1611
26. Shi L, Tu Y-Q, Wang M, Zhang F-M, Fan C-A, Zhao Y-M, Xia WJ (2005) *J Am Chem Soc* 127:10836
27. Jiang Y-J, Tu Y-Q, Zhang E, Zhang S-Y, Cao K, Shi L (2008) *Adv Synth Catal* 350:552
28. Zhang S-Y, Tu Y-Q, Fan C-A, Zhang F-M, Shi L (2009) *Angew Chem Int Ed* 48:8761
29. Zhang S-Y, Tu Y-Q, Fan C-A, Jiang Y-J, Shi L, Cao K, Zhang E (2008) *Chem Eur J* 14:10201
30. Kamitanaka T, Hikida T, Hayashi S, Kishida N, Matsuda T, Harada T (2007) *Tetrahedron Lett* 48:8460
31. Liu Z-Q, Sun L, Wang J-G, Han J, Zhao YK, Zhou B (2009) *Org Lett* 11:1437
32. Seidel A (ed) (2004) *Kirk-Othmer's encyclopedia of chemical technology*, vol 2, 5th edn. Wiley, Hoboken, p 234
33. Banerjee AK, Poon PS, Laya MS, Vera WJ (2004) *Russ Chem Rev* 73:621
34. Cheikh RB, Chaabouni R, Laurent A, Mison P, Nafti A (1983) *Synthesis* 685
35. Laurent A, Mison P, Nafti A, Cheikh RB, Chaabouni R (1984) *J Chem Res* 354
36. Johannsen M, Jørgensen KA (1998) *Chem Rev* 98:1689
37. Acemoglu L, Williams MJ (2002) In: Negishi E-i, de Meijere A (eds) *Handbook of organopalladium chemistry for organic synthesis*, vol 2. Wiley, New York, p 1689
38. Trost BM, Crawley ML (2003) *Chem Rev* 103:2921
39. Trost BM (2004) *J Org Chem* 69:5813
40. Miyabe H, Takemoto Y (2005) *Synlett* 1641
41. Takeuchi R, Kezuka S (2006) *Synthesis* 3349
42. Helmchen G, Dahnz A, Duebon P, Schelwies M, Weihofen R (2007) *Chem Commun* 675
43. Lu Z, Ma S (2008) *Angew Chem Int Ed* 47:258
44. Oguni N, Omi T (1984) *Tetrahedron Lett* 25:2823
45. Kitamura M, Suga S, Kawai K, Noyori R (1986) *J Am Chem Soc* 108:6071
46. Pu L, Yu H-B (2001) *Chem Rev* 101:757
47. Betancort JM, Garcia C, Walsh PJ (2004) *Synlett* 749
48. Ramón DJ, Yus M (2004) *Angew Chem Int Ed* 43:284
49. Garcia C, Martin VS (2006) *Curr Org Chem* 10:1849
50. Ramón DJ, Yus M (2006) *Chem Rev* 106:2126
51. Hatano M, Ishihara K (2008) *Chem Rec* 8:143
52. Oppolzer W, Radinov R (1992) *Helv Chim Acta* 75:170
53. Oppolzer W, Radinov R (1993) *J Am Chem Soc* 115:1593
54. Soai K, Takahashi KJ (1994) *Chem Soc Perkin Trans 1* 1257

55. Wipf P, Xu W (1994) *Tetrahedron Lett* 35:5197
56. Oppolzer W, Radinov RN, De Brabander J (1995) *Tetrahedron Lett* 36:2607
57. Wipf P, Ribe S (1998) *J Org Chem* 63:6454
58. Oppolzer W, Radinov RN, El-Sayed E (2001) *J Org Chem* 66:4766
59. Dahmen S, Bräse S (2001) *Org Lett* 3:4119
60. Chen YK, Lurain AE, Walsh PJ (2002) *J Am Chem Soc* 124:12225
61. Ji J-X, Qiu L-Q, Yip CW, Chan ASC (2003) *J Org Chem* 68:1589
62. Lurain AE, Walsh PJ (2003) *J Am Chem Soc* 125:10677
63. Ko D-H, Kang S-W, Kim KH, Chung Y, Ha D-C (2004) *Bull Korean Chem Soc* 25:35
64. Sprout CM, Richmond ML, Seto CT (2004) *J Org Chem* 69:6666
65. Jeon S-J, Chen YK, Walsh PJ (2005) *Org Lett* 7:1729
66. Lauterwasser F, Gall J, Hoefener S, Bräse S (2006) *Adv Synth Catal* 348:2068
67. Jeon S-J, Fisher EL, Carroll PJ, Walsh PJ (2006) *J Am Chem Soc* 128:9618
68. Salvi L, Jeon S-J, Fisher EL, Carroll PJ, Walsh PJ (2007) *J Am Chem Soc* 129:16119
69. Wu H-L, Wu P-Y, Uang B-J (2007) *J Org Chem* 72:5935
70. Chai Z, Liu X-Y, Zhang J-K, Zhao G (2007) *Tetrahedron Asymmetry* 18:724
71. Wipf P, Kendall C (2002) *Chem Eur J* 8:1778
72. Wipf P, Nunes RL (2004) *Tetrahedron* 60:1269
73. Ramón DJ, Yus M (1998) *Tetrahedron Lett* 39:1239
74. Ramón DJ, Yus M (1998) *Tetrahedron* 54:5651
75. Dosa PI, Fu GC (1998) *J Am Chem Soc* 120:445
76. Li H, Walsh PJ (2004) *J Am Chem Soc* 126:6538
77. Li H, Walsh PJ (2005) *J Am Chem Soc* 127:8355
78. Jeon S-J, Li H, Garcia C, La Rochelle LK, Walsh PJ (2005) *J Org Chem* 70:448
79. Schmidt F, Rudolph J, Bolm C (2006) *Synthesis* 3625
80. Bloch R (1998) *Chem Rev* 98:1407
81. Kobayashi S, Ishitani H (1999) *Chem Rev* 99:1069
82. Vilaivan T, Bhanthumnavin W, Sritana-Anant Y (2005) *Curr Org Chem* 9:1315
83. Schmidt F, Stemmler RT, Rudolph J, Bolm C (2006) *Chem Soc Rev* 35:454
84. K-i Y, Tomioka K (2008) *Chem Rev* 108:2874
85. Soai K, Hatanaka T, Miyazawa T (1992) *J Chem Soc Chem Commun* 1097
86. Katritsky AR, Harris PA (1992) *Tetrahedron Asymmetry* 3:437
87. Andersson PG, Guijarro D, Tanner D (1996) *Synlett* 727
88. Andersson PG, Guijarro D, Tanner D (1997) *J Org Chem* 62:7364
89. Guijarro D, Pinho P, Andersson PG (1998) *J Org Chem* 63:2530
90. Brandt P, Hedberg C, Lawonn K, Pinho P, Andersson PG (1999) *Chem Eur J* 5:1692
91. Jimeno C, Reddy S, Sola L, Moyano A, Pericas MA, Riera A (2000) *Org Lett* 2:3157
92. Sato I, Kodaka R, Soai K (2001) *J Chem Soc Perkin Trans 1* 22:2912
93. Zhang X, Lin W, Gong L, Mi A, Cui X, Jiang Y, Choi MCK, Chan ASC (2002) *Tetrahedron Lett* 43:1535
94. Sato I, Hosoi K, Kodaka R, Soai K (2002) *Eur J Org Chem* 18:3115
95. Liu H, Zhang H-L, Wang S-J, Mi A-Q, Jiang Y-Z, Gong L-Z (2005) *Tetrahedron Asymmetry* 16:2901
96. Almansa R, Guijarro D, Yus M (2007) *Tetrahedron Asymmetry* 18:2828
97. Fujihara H, Nagai K, Tomioka K (2000) *J Am Chem Soc* 122:12055
98. Nagai K, Fujihara H, Kuriyama M, Yamada K-I, Tomioka K (2002) *Chem Lett* 8
99. Shi M, Wang C-J (2003) *J Org Chem* 68:6229
100. Soeta T, Nagai K, Fujihara H, Kuriyama M, Tomioka K (2003) *J Org Chem* 68:9723
101. Shi M, Zhang W (2003) *Tetrahedron Asymmetry* 14:3407
102. Shi M, Wang C-J (2003) *Adv Synth Catal* 345:971
103. Boezio AA, Charette AB (2003) *J Am Chem Soc* 125:1692
104. Boezio AA, Pytkowicz J, Côté A, Charette AB (2003) *J Am Chem Soc* 125:14260
105. Côté A, Boezio AA, Charette AB (2004) *Proc Nat Acad Sci U S A* 101:5405

106. Côté A, Boezio AA, Charette AB (2004) *Angew Chem Int Ed* 43:6525
107. Wang M-C, Xu C-L, Zou Y-X, Liu H-M, Wang D-K (2005) *Tetrahedron Lett* 46:5413
108. Desrosiers J-N, Côté A, Charette AB (2005) *Tetrahedron* 61:6186
109. Wang M-C, Liu L-T, Hua Y-Z, Zhang J-S, Shi Y-Y, Wang D-K (2005) *Tetrahedron Asymmetry* 16:2531
110. Kim B-S, Kang S-W, Kim KH, Ko D-H, Chung Y, Ha DC (2005) *Bull Korean Chem Soc* 26:1501
111. Lauzon C, Charette AB (2006) *Org Lett* 8:2743
112. Wang M-C, Xu C-L, Cheng F, Ding X (2006) *Tetrahedron* 62:12220
113. Shi M, Lei Z-Y, Xu Q (2006) *Adv Synth Catal* 348:2237
114. Chen J, Li D, Ma H, Cun L, Zhu J, Deng J, Liao J (2008) *Tetrahedron Lett* 49:6921
115. Perron Q, Alexakis A (2008) *Tetrahedron Asymmetry* 19:1871
116. Pizzuti MG, Minnaard AJ, Feringa BL (2008) *J Org Chem* 73:940
117. Nishimura T, Yasuhara Y, Hayashi T (2006) *Org Lett* 8:979
118. Chen J, Li D, Ma H, Cun L, Zhu J, Deng J, Liao J (2008) *Tetrahedron Lett* 49:6921
119. Porter JR, Traverse JF, Hoveyda AH, Snapper ML (2001) *J Am Chem Soc* 123:984
120. Porter JR, Traverse JF, Hoveyda AH, Snapper ML (2001) *J Am Chem Soc* 123:10409
121. Akullian LC, Snapper ML, Hoveyda AH (2003) *Angew Chem Int Ed* 42:4244
122. Akullian LC, Porter JR, Traverse JF, Snapper ML, Hoveyda AH (2005) *Adv Synth Catal* 347:417
123. Basra S, Fennie MW, Kozlowski MC (2006) *Org Lett* 8:2659
124. Fu P, Snapper ML, Hoveyda AH (2008) *J Am Chem Soc* 130:5530
125. Tomioka K (1990) *Synthesis* 541
126. Berrisford DJ (1995) *Angew Chem Int Ed* 34:178
127. Denmark SE, Nicaise OJ-C (1996) *Chem Commun* 999
128. Enders D, Reinhold U (1997) *Tetrahedron Asymmetry* 8:1895
129. Luderer MR, Bailey WF, Luderer MR, Fair JD, Dancer RJ, Sommer MB (2009) *Tetrahedron Asymmetry* 20:981
130. Goldfuss B (2003) *Top Organomet Chem* 5:21
131. Iguchi M, K-i Y, Tomioka K (2003) *Top Organomet Chem* 5:37
132. Goldfuss B (2005) *Synthesis* 14 2271
133. Hayashi T, Ishigedani M (2000) *J Am Chem Soc* 122:976
134. Hayashi T, Ishigedani M (2001) *Tetrahedron* 57:2589
135. Hayashi T, Kawai M, Tokunaga N (2004) *Angew Chem Int Ed* 43:6125
136. Kuriyama M, Soeta T, Hao X, Chen Q, Tomioka K (2004) *J Am Chem Soc* 126:8128
137. Tokunaga N, Otomaru Y, Okamoto K, Ueyama K, Shintani R, Hayashi T (2004) *J Am Chem Soc* 126:13584
138. Weix DJ, Shi Y, Ellman JA (2005) *J Am Chem Soc* 127:1092
139. Otomaru Y, Tokunaga N, Shintani R, Hayashi T (2005) *Org Lett* 7:307
140. Duan H-F, Jia Y-X, Wang L-X, Zhou Q-L (2006) *Org Lett* 8:2567
141. Wang Z-Q, Feng C-G, Xu M-H, Lin G-Q (2007) *J Am Chem Soc* 129:5336
142. Marelli C, Monti C, Gennari C, Piarulli U (2007) *Synlett* 14 2213
143. Nakagawa H, Rech JC, Sindelar RW, Ellman JA (2007) *Org Lett* 9:5155
144. Trincado M, Ellman JA (2008) *Angew Chem Int Ed* 47:5623
145. Kurihara K, Yamamoto Y, Miyaura N (2009) *Adv Synth Catal* 351:260
146. Okamoto K, Hayashi T, Rawal VH (2009) *Chem Commun* 4815
147. Kakuuchi A, Taguchi T, Hanzawa Y (2003) *Tetrahedron Lett* 44:923
148. Wipf P, Kendall C, Stephenson CRJ (2003) *J Am Chem Soc* 125:761
149. Patel SJ, Jamison TF (2004) *Angew Chem Int Ed* 43:3941
150. (2007) *Top Curr Chem* 279 1
151. Ojima I, Tzamarioudaki M, Li Z, Donovan RJ (1996) *Chem Rev* 96:635
152. Montgomery J (2000) *Acc Chem Res* 33:467

153. Montgomery J, Amarashinghe KKD, Chowdhury SK, Oblinger E, Seo J, Savchenko AV (2002) *Pure Appl Chem* 74:129
154. Ikeda S-I (2003) *Angew Chem Int Ed* 42:5120
155. Miller KM, Molinaro C, Jamison TF (2003) *Tetrahedron Asymmetry* 14:3619
156. Montgomery J (2004) *Angew Chem Int Ed* 43:3890
157. Montgomery J, Sormunen GJ (2007) *Top Curr Chem* 279:1
158. Moslin RM, Miller-Moslin K, Jamison TF (2007) *Chem Commun* 4441
159. Ojima I, Tzamarioudaki M, Tsai C-Y (1994) *J Am Chem Soc* 116:3643
160. Crowe WE, Rachita MJ (1995) *J Am Chem Soc* 117:6787
161. Kablaoui NM, Buchwald SL (1995) *J Am Chem Soc* 117:6785
162. Oblinger E, Montgomery J (1997) *J Am Chem Soc* 119:9065
163. Tang X-Q, Montgomery J (1999) *J Am Chem Soc* 121:6098
164. Tang X-Q, Montgomery J (2000) *J Am Chem Soc* 122:6950
165. Mahandru GM, Liu G, Montgomery J (2004) *J Am Chem Soc* 126:3698
166. Knapp-Reed B, Mahandru GM, Montgomery J (2005) *J Am Chem Soc* 127:13156
167. Chaulagain MR, Sormunen GJ, Montgomery J (2007) *J Am Chem Soc* 129:9568
168. Huang W-S, Chan J, Jamison TF (2000) *Org Lett* 2:4221
169. Miller KM, Huang W-S, Jamison TF (2003) *J Am Chem Soc* 125:3442
170. Miller KM, Jamison TF (2005) *Org Lett* 7:3077
171. Liu P, McCaren P, Cheong PH-Y, Jamison TF, Houk KN (2010) *J Am Chem Soc* 132:2050
172. Takai K, Sakamoto S, Isshiki T (2003) *Org Lett* 5:653
173. Ngai M-Y, Barchuk A, Krische MJ (2007) *J Am Chem Soc* 129:280
174. Kong J-R, Ngai M-Y, Krische MJ (2006) *J Am Chem Soc* 128:718
175. Cho C-W, Krische MJ (2006) *Org Lett* 8:3873
176. Hong Y-T, Cho C-W, Skucas E, Krische MJ (2007) *Org Lett* 9:3745
177. Komanduri V, Krische MJ (2006) *J Am Chem Soc* 128:16448
178. Komanduri V, Grant CD, Krische MJ (2008) *J Am Chem Soc* 130:12592
179. Kong J-R, Krische MJ (2006) *J Am Chem Soc* 128:16040
180. Skucas E, Kong J-R, Krische MJ (2007) *J Am Chem Soc* 129:7242
181. Han SB, Kong J-R, Krische MJ (2008) *Org Lett* 10:4133
182. Williams VM, Kong J-R, Ko B-J, Mantri Y, Brodbelt JS, Baik M-H, Krische MJ (2009) *J Am Chem Soc* 131:16054
183. Dewar MJS (1951) *Bull Soc Chim Fr* C71
184. Chatt J, Duncanson LA (1953) *J Chem Soc* 2939
185. Dewar MJS, Ford GP (1979) *J Am Chem Soc* 101:783
186. Li J, Schreckenbach G, Ziegler T (1995) *J Am Chem Soc* 117:486
187. Vaska L, Peone Jr J (1971) *Chem Commun* 418
188. Haynes A, McNish J, Pearson JMJ (1998) *Organomet Chem* 551:339
189. Grotjohn DB, Collins LSB, Wolpert M, Bikzhanova GA, Lo HC, Combs D, Hubbard JL (2001) *J Am Chem Soc* 123:8260
190. Kirchner RM, Ibers JA (1973) *J Am Chem Soc* 95:1095
191. Calabrese JC, Roe DC, Thorn DL, Tulip TH (1984) *Organometallics* 3:1223
192. Rappoli BJ, Churchill MR, Janik TS, Rees WM, Atwood JD (1987) *J Am Chem Soc* 109:5145
193. Rees WM, Churchill MR, Fettinger JC, Atwood JD (1987) *J Organomet Chem* 319:411
194. Marinelli G, Streib WE, Huffman JC, Caulton KG, Gagne MR, Takats J, Dartiguenave M, Chardon C, Jackson SA, Eisenstein O (1990) *Polyhedron* 9:1867
195. Masashi Y, Sakaki S (2002) *J Am Chem Soc* 124:7588
196. Barchuk A, Ngai M-Y, Krische MJ (2007) *J Am Chem Soc* 129:8432
197. Ngai M-Y, Barchuk A, Krische MJ (2007) *J Am Chem Soc* 129:12644
198. K-i Fujita, Yamashita M, Puschmann F, Alvarez-Flacon MM, Incarvito CD, Hartwig JF (2006) *J Am Chem Soc* 128:9044
199. Ramachandran PV (2002) *Aldrichim Acta* 35:23

200. Kennedy JWJ, Hall DG (2003) *Angew Chem Int Ed* 42:4732
201. Denmark SE, Fu J (2003) *Chem Rev* 103:2763
202. Yu C-M, Youn J, Jung H-K (2006) *Bull Korean Chem Soc* 27:463
203. Marek I, Sklute G (2007) *Chem Commun* 1683
204. Hall DG (2007) *Synlett* 1644
205. Mikhailov BM, Bubnov YN (1964) *Izv Akad Nauk SSSR, Ser Khim* 1874
206. Hosomi A, Sakurai H (1976) *Tetrahedron Lett* 17:1295
207. Herold T, Hoffmann RW (1978) *Angew Chem Int Ed Engl* 17:768
208. Hoffmann RW, Herold T (1981) *Chem Ber* 114:375
209. Hayashi T, Konishi M, Kumada M (1982) *J Am Chem Soc* 104:4963
210. Brown HC, Jadhav PK (1983) *J Am Chem Soc* 105:2092
211. Roush WR, Walts AE, Hoong LK (1985) *J Am Chem Soc* 107:8186
212. Reetz M (1988) *Pure Appl Chem* 60:1607
213. Short RP, Masamune S (1989) *J Am Chem Soc* 111:1892
214. Corey EJ, Yu C-M, Kim SS (1989) *J Am Chem Soc* 111:5495
215. Seebach D, Beck AK, Imwinkeiried R, Roggo S, Wonnacott A (1987) *Helv Chim Acta* 70:954
216. Riediker M, Duthaler RO (1998) *Angew Chem Int Ed Engl* 28:494
217. Panek JS, Yang M (1991) *J Am Chem Soc* 113:6594
218. Kinnaird JWA, Ng PY, Kubota K, Wang X, Leighton JL (2002) *J Am Chem Soc* 124:7920
219. Hackman BM, Lombardi PJ, Leighton JL (2004) *Org Lett* 6:4375
220. Burgos CH, Canales E, Matos K, Soderquist JA (2005) *J Am Chem Soc* 127:8044
221. Furuta K, Mouri M, Yamamoto H (1991) *Synlett* 561
222. Costa AL, Piazza MG, Tagliavini E, Trombini C, Umami-Ronchi A (1993) *J Am Chem Soc* 115:7001
223. Keck GE, Tarbet KH, Geraci LS (1993) *J Am Chem Soc* 115:8467
224. Denmark SE, Coe DM, Pratt NE, Griedel BD (1994) *J Org Chem* 59:6161
225. Tabuchi T, Inanaga J, Yamaguchi M (1986) *Tetrahedron Lett* 27:1195
226. Takahara JP, Masuyama Y, Kurusu Y (1992) *J Am Chem Soc* 114:2577
227. Kimura M, Ogawa Y, Shimizu M, Sueishi M, Tanaka S, Tamaru Y (1998) *Tetrahedron Lett* 39:6903
228. Kimura M, Tomizawa T, Horino Y, Tanaka S, Tamaru Y (2000) *Tetrahedron Lett* 41:3627
229. Kimura M, Shimizu M, Shibata K, Tazoe M, Tamaru Y (2003) *Angew Chem Int Ed* 42:3392
230. Zanoni G, Gladiali S, Marchetti A, Piccinini P, Tredici I, Vidari G (2004) *Angew Chem Int Ed* 43:846
231. Kimura M, Shimizu M, Tanaka S, Tamaru Y (2005) *Tetrahedron* 61:3709
232. Howell GP, Minnaard AJ, Feringa BL (2006) *Org Biomol Chem* 4:1278
233. Barczak NT, Grote RE, Jarvo ER (2007) *Organometallics* 26:4863
234. Masuyama Y, Kaneko Y, Kurusu Y (2004) *Tetrahedron Lett* 45:8969
235. Vasylyev M, Alper H (2010) *J Org Chem*. doi:10.1021/jo902706p
236. Tsuji Y, Mukai T, Kondo T, Watanabe Y (1989) *J Organomet Chem* 369:C51
237. Kondo T, Ono H, Satake N, Mitsudo T-a, Watanabe Y (1945) *Organometallics* 1995:14
238. Denmark SE, Nguyen ST (2009) *Org Lett* 11:781
239. Masuyama Y, Chiyo T, Kurusu Y (2005) *Synlett* 14 2251
240. Banerjee M, Roy S (2006) *J Mol Catal A* 246:231
241. Masuyama Y, Marukawa M (2007) *Tetrahedron Lett* 48:5963
242. Masuyama Y (1994) In: Liebeskind LS (ed) *Advances in metal-organic chemistry*, vol 3. JAI, Greenwich, p 255
243. Tamaru Y (2002) In: Negishi E-i, de Meijere A (eds) *Handbook of organopalladium chemistry for organic synthesis*, vol 2. Wiley, New York, p 1917
244. Tamaru Y (1999) In: Tsuji J (ed) *Perspectives in organopalladium chemistry for the XXI century*. Elsevier, Amsterdam, p 215
245. Kondo T, Mitsudo T-A (2002) *Curr Org Chem* 6:1163

246. Tamaru Y (2005) *Eur J Org Chem* 2647
247. Zanoni G, Pontiroli A, Marchetti A, Vidari G (2007) *Eur J Org Chem* 3599
248. Bandini M, Cozzi PG, Umani-Ronchi A (2000) *Polyhedron* 19:537
249. Bandini M, Cozzi PG, Umani-Ronchi A (2001) *Tetrahedron* 57:835
250. Bandini M, Cozzi PG, Umani-Ronchi A (2000) *Angew Chem Int Ed* 39:2327
251. Inoue M, Suzuki T, Nakada M (2003) *J Am Chem Soc* 125:1140
252. Lee J-Y, Miller JJ, Hamilton SS, Sigman MS (2005) *Org Lett* 7:1837
253. McManus HA, Cozzi PG, Guiry PJ (2006) *Adv Synth Catal* 348:551
254. Xia G, Yamamoto H (2006) *J Am Chem Soc* 128:2554
255. Hargaden GC, Müller-Bunz H, Guiry PJ (2007) *Eur J Org Chem* 4235
256. Hargaden GC, O'Sullivan TP, Guiry PJ (2008) *Org Biomol Chem* 6:562
257. Liu X, Henderson JA, Sasaki T, Kishi Y (2009) *J Am Chem Soc* 131:16678
258. Avalos M, Babiano R, Cintas P, Jiménez JL, Palacios JC (1999) *Chem Soc Rev* 28:169
259. Bandini M, Cozzi PG, Umani-Ronchi A (2002) *Chem Commun* 919
260. Hargaden GC, Guiry PJ (2007) *Adv Synth Catal* 349:2407
261. Inoue M, Suzuki T, Kinoshita A, Nakada M (2008) *Chem Rec* 8:169
262. Mikami K, Shimizu M (1992) *Chem Rev* 92:1021
263. Berrisford DJ, Bolm C (1995) *Angew Chem Int Ed* 34:1717
264. Johnson JS, Evans DA (2000) *Acc Chem Res* 33:325
265. Clarke ML, France MB (2008) *Tetrahedron* 64:9003
266. Ho C-Y, Ng S-S, Jamison TF (2006) *J Am Chem Soc* 128:5362
267. Ng S-S, Ho C-Y, Jamison TF (2006) *J Am Chem Soc* 128:11513
268. Ho C-Y, Schleicher KD, Chan C-W, Jamison TF (2009) *Synlett* 2565
269. Skucas E, Bower JF, Krische MJ (2007) *J Am Chem Soc* 129:12678
270. Bower JF, Skucas E, Patman RL, Krische MJ (2007) *J Am Chem Soc* 129:15134
271. Han SB, Kim I-S, Han H, Krische MJ (2009) *J Am Chem Soc* 131:6916
272. Saito T, Yokozawa T, Ishizaki T, Moroi T, Sayo N, Miura T, Kumobayashi H (2001) *Adv Synth Catal* 343:264
273. Bower JF, Patman RL, Krische MJ (2008) *Org Lett* 10:1033
274. Zbieg JR, Fukuzumi T (2010) *Adv Synth Catal* 352: (In Press)
275. Shibahara F, Bower JF, Krische MJ (2008) *J Am Chem Soc* 130:6338
276. Shibahara F, Bower JF, Krische MJ (2008) *J Am Chem Soc* 130:14120
277. Obora Y, Hatanaka S, Ishii Y (2009) *Org Lett* 11:3510
278. Patman RL, Chaulagain MR, Williams VM, Krische MJ (2009) *J Am Chem Soc* 131:2066
279. Kim IS, Ngai M-Y, Krische MJ (2008) *J Am Chem Soc* 130:6340
280. Kim IS, Ngai M-Y, Krische MJ (2008) *J Am Chem Soc* 130:14891
281. Lopez F, Ohmura T, Hartwig JF (2003) *J Am Chem Soc* 125:3426
282. Shu C, Hartwig JF (2004) *Angew Chem Int Ed* 43:4794
283. Ueno S, Hartwig JF (2008) *Angew Chem Int Ed* 47:1928
284. Kondo T, Mukai T, Watanabe Y (1991) *J Org Chem* 56:487
285. Poss CS, Schreiber SL (1994) *Acc Chem Res* 27:9
286. Lu Y, Kim I-S, Hassan A, Del Valle DJ, Krische MJ (2009) *Angew Chem Int Ed* 48:5018
287. Kogure T, Eliel ELJ (1984) *Org Chem* 49:576
288. Midland MM, Gabriel J (1985) *J Org Chem* 50:1144
289. Minami N, Ko SS, Kishi Y (1982) *J Am Chem Soc* 104:1109
290. Ko SY, Lee AWM, Masamune S, Reed LA III, Sharpless KB, Walker FJ (1983) *Science* 220:949
291. Kobayashi S, Ohtsubo A, Mukaiyama T (1991) *Chem Lett* 831
292. Hammadi A, Nuzillard JM, Poulin JC, Kagan HB (1992) *Tetrahedron Asymmetry* 3:1247–1262
293. Doyle MP, Kalinin AV, Ene DG (1996) *J Am Chem Soc* 118:8837
294. Trost BM, Calkins TL, Oertelt C, Zambrano J (1998) *Tetrahedron Lett* 39:1713
295. Balskus EP, Jacobsen EN (2007) *Science* 317:1736

296. Han SB, Kong JR, Krische MJ (2008) *Org Lett* 10:4133
297. Hassan A, Lu Y, Krische MJ (2009) *Org Lett* 11:3112
298. Kim IS, Han SB, Krische MJ (2009) *J Am Chem Soc* 131:2514
299. Han SB, Han H, Krische MJ (2010) *J Am Chem Soc* 132:1760
300. Han SB, Gao X, Krische MJ (2010) *J Am Chem Soc* 132:9153
301. Zhang YJ, Yang JH, Kim SH, Krische MJ (2010) *J Am Chem Soc* 132:4562
302. Masuyama Y, Takahara JP, Kurusu Y (1988) *J Am Chem Soc* 110:4473
303. Masuyama Y, Otake K, Kurusu Y (1988) *Tetrahedron Lett* 29:3563
304. Takahara JP, Masuyama Y, Kurusu Y (1992) *J Am Chem Soc* 114:2577
305. Araki S, Kameda K, Tanaka J, Hirashita T, Yamamura H, Kawai M (2010) *J Org Chem* 66:7919
306. Gagliardo M, Selander N, Mehendale NC, van Koten G, Klein Gebbink RJM, Szabó KJ (2008) *Chem Eur J* 14:4800
307. Bandini M, Cozzi PG, Licciulli S, Umani-Ronchi A (2004) *Synthesis* 409
308. Nakajima M, Saito M, Hashimoto S (2000) *Chem Pharm Bull* 48:306
309. Itoh J, Han SB, Krische MJ (2009) *Angew Chem Int Ed* 48:6313
310. Kitajima M, Mori I, Arai K, Kogure N, Takayama H (2006) *Tetrahedron Lett* 47:3199

# Ir-Catalyzed Functionalization of C–H Bonds

Jongwook Choi and Alan S. Goldman

**Abstract** The ability to selectively functionalize C–H bonds holds enormous potential value in virtually every sphere of organic chemistry, from fuels to pharmaceuticals. Transition metal complexes have shown great promise in this context. Iridium provided the first examples of oxidative addition of C–H bonds; this addition is key to iridium's leading role in alkane dehydrogenation and related reactions. Catalysts based on iridium have also proven highly effective for valuable borylations of C–H bonds and, to a lesser extent, for C–Si coupling. Compared with other platinum group metals, iridium chemistry has not been developed as extensively for the elaboration of C–C bonds from C–H bonds, but significant promise is indicated, particularly for coupling with simple hydrocarbons which lack functionalities that can act as directing groups.

**Keywords** Alkane metathesis · Borylation · C–H bond activation · Dehydrogenation · Hydroarylation · Iridium catalyst · Silylation

## Contents

1	Introduction and Background Stoichiometric Chemistry .....	140
2	Catalytic Dehydrogenation .....	141
2.1	Dehydrogenation of Alkanes .....	141
2.2	Alkane Metathesis .....	145
2.3	Dehydrogenation of Alkyl Groups in Molecules Containing Heteroatoms .....	147
3	Borylation of Arenes .....	148
4	Silylation of Arenes .....	152
5	C–C Bond Formation .....	154
5.1	Hydroarylation and Hydrovinylation of C–C Multiple Bonds .....	154
5.2	Aryl–Aryl Coupling Reactions .....	158
5.3	Coupling with $sp^3$ C–H Bonds .....	159




6	Others .....	160
6.1	$\alpha,\alpha$ -Dehydrogenation and Reactions with Heterocumulenes .....	160
6.2	C–H Bond Activation by Metal Carbenoids or Nitrenoids .....	162
7	Concluding Remarks .....	163
	References .....	163

## 1 Introduction and Background Stoichiometric Chemistry

Given the ubiquitous nature of carbon–hydrogen bonds, the ability to selectively catalyze their transformation is one of the most attractive goals of modern synthetic chemistry [1–9]. The potential applications of such chemistry range in scale from synthesis of the most complex fine chemicals to the development of new routes to transportation fuels.

In the realm of C–H bond transformations applied toward the synthesis of fine chemicals, iridium has not achieved the prominence attained in recent years by the second-row platinum group metals, particularly palladium [10]. A notable exception, however, has been the leading role of iridium in the valuable chemistry of arene borylation [11].


With respect to reactions of the simplest of organic molecules, alkanes, iridium has earned a special place. Indeed, iridium has played a particularly important role in the development and mechanistic understanding of the reactions of organometallic complexes with simple molecules in general. Vaska's complex,  $\text{Ir}(\text{Cl})(\text{CO})(\text{PPh}_3)_2$ , offered the archetype of the oxidative addition reaction with the discovery of addition of  $\text{H}_2$  to give what is formally an Ir(III) complex (in contrast with organic chemical formalisms wherein addition of  $\text{H}_2$  is a reduction) [12]. The same complex offered the archetypal oxidative addition of C–X bonds with the report of addition of methyl iodide [13]. Undoubtedly, it is not a coincidence that the first example of oxidative addition of an alkane C–H bond was to an iridium center: the seminal report by Bergman in which photogenerated  $\text{Cp}^*\text{Ir}(\text{PMe}_3)$  was found to add a wide range of alkane C–H bonds [14].



As in the case of the H–H and C–I addition to Vaska's complex, reaction 1 proceeds via addition to Ir(I) to give a stable formally six-coordinate Ir(III) complex.

The first example of dehydrogenation of alkanes by a transition metal complex was also achieved with iridium. The cationic iridium(III) complex,  $[\text{IrH}_2(\text{acetone})_2(\text{PPh}_3)_2]^+ \text{BF}_4^-$ , with TBE (3,3-dimethyl-1-butene or *t*-butylethene) as

a hydrogen acceptor dehydrogenates cyclopentane or cyclooctane (COA) to give the cyclopentadienyl and cyclooctadiene adducts, respectively (2) [15].



Presumably, TBE first accepts  $H_2$ , thereby generating yet another example of an Ir(I) species that undergoes C–H bond oxidative addition. As in the example of **1**, the resulting Ir product strongly binds the transformed hydrocarbon. Indeed, the ability of Ir to give stable complexes generally reflects a mixed blessing, valuable for the purposes of characterization of complexes and mechanism, but often leading to species too stable to permit catalytic reactivity. There are, however, many exceptions to the tendency of iridium to give species too stable for catalysis, and as we shall see, the first well-characterized examples of catalytic alkane functionalization were dominated by iridium.

In this review, the recent developments in catalytic functionalization of C–H bonds by iridium complexes will be emphasized. For more information on previous work and their details, there are excellent reviews published recently on C–H bond functionalization by transition metal complexes generally [1–9], as well as with an emphasis on iridium [11, 16].

## 2 Catalytic Dehydrogenation

### 2.1 Dehydrogenation of Alkanes


The dehydrogenation of unactivated alkanes to olefins by the removal of two hydrogen atoms is in principle an elegant way to introduce a valuable functionality. This transformation has great potential value considering the importance of olefin intermediates in the commodity and petrochemical industries and the great versatility of olefins as reagents in organic synthesis.

As noted above, the first stoichiometric dehydrogenation of alkanes by soluble homogeneous metal complexes, reported by Crabtree et al. [15], was a reaction of an iridium complex,  $[IrH_2(PPh_3)_2(Me_2CO)_2]^+$ . The closely related bis- $P(p\text{-}FC_6H_4)_3$  complex was subsequently reported to catalytically dehydrogenate COA using TBE as hydrogen acceptor, in the presence of a base that was found to deprotonate the complex [17]. Felkin and coworkers later used the neutral iridium (and ruthenium) polyhydride systems,  $(^iPr_3P)_2IrH_5$ ,  $[(p\text{-}F\text{-}C_6H_4)_3P]_2IrH_5$ , and  $[(p\text{-}F\text{-}C_6H_4)_3P]_3RuH_4$ , for catalytic COA/TBE transfer dehydrogenation [18–20]. Using  $(^iPr_3P)_2IrH_5$ , up to

70 catalytic turnovers were obtained. Soon thereafter,  $[(p\text{-F-C}_6\text{H}_4)_3\text{P}]_2\text{IrH}_2(\kappa^2\text{-O}_2\text{CCF}_3)$  was found to effect dehydrogenation of linear and cyclic alkanes, both thermally (with TBE as acceptor) and photochemically (without hydrogen acceptor) [21, 22]. This study laid the mechanistic groundwork for future alkane dehydrogenation chemistry, in particular highlighting the importance of the key 14-electron  $\text{IrL}_2\text{X}$  unit (in this case, X being the dechelated  $\kappa^1$ -carboxylate ligand).


The systems discussed above suffered from limited turnover numbers ( $\leq 70$ ), presumably due to catalyst degradation; species active enough to react with alkanes could be expected to react, intra- or intermolecularly, with their own ligands. The first molecular catalysts to circumvent this limitation were complexes based on the closest possible analog of  $\text{IrL}_2\text{X}$ , specifically  $\text{Rh}(\text{PMe}_3)_2\text{Cl}$ . Eisenberg had demonstrated that  $\text{M}(\text{PPh}_3)_2\text{Cl}(\text{CO})$  ( $\text{M} = \text{Rh}$  or  $\text{Ir}$ ) catalyzed the photochemical carbonylation of benzene [23, 24]. Tanaka, observing that  $\text{PMe}_3$  was more favorable than  $\text{PPh}_3$  for C–H activation in the case of **1**, found that  $\text{Rh}(\text{PMe}_3)_2\text{Cl}(\text{CO})$  was accordingly more efficient for photochemical carbonylation [25, 26], and this same complex was soon found to be even more effective for photochemical alkane dehydrogenation [27–30]. It was later reported by Goldman and coworkers that species  $\text{Rh}(\text{PMe}_3)_2\text{CIL}$  ( $\text{L} = \text{CO}$  or phosphine) were excellent alkane transfer-dehydrogenation catalysts although (quite surprisingly) only in the presence of dihydrogen [31, 32]. The requirement of  $\text{H}_2$  atmosphere, although mechanistically interesting, was a practical drawback because the system hydrogenated more than one mol acceptor per mol dehydrogenated product.

In 1996, Jensen and Kaska reported that the pincer-ligated iridium complex,  $(^t\text{Bu}^4\text{PCP})\text{IrH}_2$  (**1-H<sub>2</sub>**) ( $^t\text{Bu}^4\text{PCP} = \kappa^3\text{-C}_6\text{H}_3\text{-2,6-}[\text{CH}_2\text{P}(t\text{-Bu})_2]_2$ ), shows excellent catalytic activity for transfer dehydrogenation of cycloalkanes (**3**) and long-term stability at high temperatures (up to  $200^\circ\text{C}$ ) [33, 34]. The high thermal stability was exploited toward the first efficient example of catalytic acceptorless dehydrogenation [35]. The *i*-propyl analog,  $(^i\text{Pr}^4\text{PCP})\text{IrH}_2$  (**2-H<sub>2</sub>**), displayed even higher catalytic activity, achieving close to 1,000 turnovers in the dehydrogenation of cyclodecane without acceptor (**4**) [36]. Most notably perhaps, Jensen and Goldman reported that complexes **1-H<sub>2</sub>** and **2-H<sub>2</sub>** selectively dehydrogenate the terminal position of *n*-alkanes to give  $\alpha$ -olefin, although subsequent isomerization leads to the formation of the thermodynamically more stable internal olefins [37].




The proposed mechanism of *n*-alkane dehydrogenation by  $(t\text{Bu}^4\text{PCP})\text{Ir}$  is shown in Scheme 1 [38]. The reactive  $14e$   $(t\text{Bu}^4\text{PCP})\text{Ir}$  species (**1**) is generated from the hydrogenation of TBE by **1**-H<sub>2</sub> and undergoes oxidative addition of terminal C–H bonds.  $\beta$ -hydride elimination then affords  $\alpha$ -olefin and regenerates **1**-H<sub>2</sub>. In the case of acceptorless dehydrogenation, instead of transferring hydrogen to an acceptor, the active species **1** is generated via thermolytic loss of H<sub>2</sub> from **1**-H<sub>2</sub> [39]. Unfortunately, catalytic activity is inhibited by the buildup of olefin product due to the formation of complexes,  $(t\text{Bu}^4\text{PCP})\text{Ir}(\text{olefin})$ . In the case of TBE, the species of this composition is the vinyl hydride shown in Scheme 1, while linear olefins give  $\pi$ -complexes, and 1-alkenes in particular bind very strongly [40, 41].

The success of derivatives of **1** and **2** as dehydrogenation catalysts has led to the investigation of numerous different pincer ligands for iridium-catalyzed alkane dehydrogenation. The “Anthrphos” pincer iridium complex (**3**-H<sub>2</sub>) was expected to afford even greater thermal stability (Fig. 1), and indeed, the catalyst can tolerate reaction temperatures up to 250°C [42]. The catalytic activity of **3**-H<sub>2</sub>, however, is much less than that of **1**-H<sub>2</sub> under comparable conditions.



**Scheme 1** Proposed mechanism of alkane dehydrogenation with acceptors



**Fig. 1** Various pincer-ligated Ir complexes


The effect of varying substituents at the *para* position of pincer ligands was investigated both computationally and experimentally [43, 44]. DFT calculations reveal that electron-donating substituents such as methoxy (OMe) favor the oxidative addition of R–H (or H<sub>2</sub>) to 14-electron **1** and derivatives; experimentally, the MeO substituent engenders a modest increase in catalytic activity for acceptorless dehydrogenation. The bis-phosphinite complex, (<sup>R4</sup>POCOP)Ir (<sup>R4</sup>POCOP =  $\kappa^3$ -2,6-C<sub>6</sub>H<sub>3</sub>(OPR<sub>2</sub>)<sub>2</sub>), reported by the Brookhart group (**4**, R = <sup>t</sup>Bu) [45–47] and Jensen group (**5**, R = <sup>i</sup>Pr) [48] shows greater activity for the “benchmark” COA/TBE transfer-dehydrogenation than does **1** under similar conditions [45–47]. The  $\nu_{\text{CO}}$  stretching frequency exhibited by the CO adduct of **4a** is greater than that of **1**, which appears to suggest that **4a** is less electron-rich than **1**. DFT calculations, however, show that oxygen substitution for CH<sub>2</sub> increases electron donation to aryl ring in the <sup>tBu4</sup>POCOP ligand, similarly to a methoxy substituent at the *para* position [44]. However, **4a** shows much less reactivity for the transfer-dehydrogenation of linear alkanes than does **1** [41], particularly in the case of 1-alkene hydrogen acceptors.

Brookhart and coworkers found significant differences between **1** and **4** when investigating the mechanism of COA dehydrogenation by precursors of **4** [45–47]. In particular, the rate of TBE hydrogenation by **4b**-H<sub>2</sub> is much faster than by **1**-H<sub>2</sub>. For example, **4b**-H<sub>2</sub> hydrogenates TBE at –70°C with a rate comparable to that by **1**-H<sub>2</sub> at 55°C. Recent DFT calculations indicate that these differences are attributable to the fact that **4** is significantly less crowded at the metal center than **1** [41]. Thus, **1** undergoes oxidative addition of the unhindered vinyl C–H bond of TBE trans to the *t*-butyl group, rather than formation of the  $\pi$ -coordinated complex as seen in the case of **4** [45–47]. Both simple coordination and hydrogenation of TBE are clearly very sterically demanding, much more so than addition of the TBE C–H vinylic bond.

As indicated above, steric factors very significantly impact the catalytic efficiency of the pincer-iridium complexes. In this context, Goldman, Brookhart, and Krogh-Jespersen conducted a combined computational–experimental study of the effect on catalytic activity achieved by substituting *t*-butyl groups with methyl groups on phosphorus [40]. DFT calculation showed that a single methyl for *t*-butyl substitution on **1**, to give (<sup>tBu3Me</sup>PcP)Ir (**6**), has a large favorable effect on the kinetics of the  $\beta$ -hydride elimination step (see Scheme 1), which is calculated to be the rate-determining step in the alkane dehydrogenation cycle. Experimentally, **6** dehydrogenates *n*-alkanes more efficiently than does **1** or even the presumably less crowded (<sup>iPr4</sup>PcP)Ir (**2**); DFT calculations are in accord with these results [40]. After the first methyl for *t*-butyl substitution, further substitutions of this type are not calculated to result in a significant increase of activity because energies of the olefin-bound resting state and the rate-determining TS (generally  $\beta$ -hydride elimination) are lowered by approximately equal amounts. Experimentally, (<sup>tBuMe</sup>PcP<sup>tBuMe</sup>)IrH<sub>4</sub> was synthesized and found to give lower turnovers than **6**; this was attributed to cluster formation, a hypothesis supported by the isolation of dimeric species bearing the (<sup>tBuMe</sup>PcP<sup>tBuMe</sup>) ligand [40].

Dehydrogenation of saturated polyolefins by pincer-iridium complexes has also been reported. Coates and Goldman have reported that (MeO-<sup>iPr4</sup>PcP)IrH<sub>4</sub>


dehydrogenates poly-1-hexene with up to 18% conversion based on C<sub>6</sub>-units (5) [49]. Dehydrogenation selectively takes place at the branches, not the backbone, in accord with the observations that the (<sup>R</sup>PCP)Ir complexes (R = <sup>t</sup>Bu and <sup>i</sup>Pr) have a strong preference for terminal dehydrogenation to give α-olefin, although this is followed by double-bond isomerization.



Recently, Scott, Brookhart, Goldman and coworkers have heterogenized (<sup>t</sup>Bu<sup>4</sup>PCP)Ir and (<sup>t</sup>Bu<sup>4</sup>POCOP)Ir derivatives by supporting them on γ-Al<sub>2</sub>O<sub>3</sub>. This was most successful with derivatives in which the pincer aryl group had a Lewis-basic *para*-substituent, e.g., NH<sub>2</sub>, or a group that could be hydrolyzed and thus react with the alumina surface, e.g., -C(O)(OMe) or -OP(<sup>t</sup>Bu)<sub>2</sub> [50].


## 2.2 Alkane Metathesis

Alkane metathesis (AM; shown for *n*-alkanes in 6), which has also been referred to as “disproportionation” or “redistribution of alkane chains,” has great potential in the context of petroleum refining.



A particularly intriguing potential application of this reaction is the upgrading of the products of Fischer–Tropsch catalysis to increase the eventual yield of desirable *n*-alkane chain lengths (typically C9–C19). FT catalysis, already practiced on a large scale commercially, may prove to be a key route to the future utilization of coal, biomass, or other nonpetroleum carbon sources (including CO<sub>2</sub> reduction powered by solar, wind, or nuclear energy) [51, 52].

Early work by Burnett and Hughes [53] and later by the Basset group [54, 55] demonstrated the feasibility of alkane metathesis by heterogeneous catalysts. However, these heterogeneous systems did not yield products with any selectivity for molecular weight. In 2006, Brookhart, Goldman, and coworkers reported alkane metathesis using a combination of two homogeneous catalysts, pincer-ligated iridium complexes (**1** or **4a**), and a molybdenum-based Schrock olefin metathesis catalyst (**7**) under relatively mild conditions [56]. The proposed pathway is shown in Scheme 2 for the very desirable case in which *n*-alkane H<sub>3</sub>C(CH<sub>2</sub>)<sub>*n*</sub>CH<sub>3</sub> is disproportionated to H<sub>3</sub>C(CH<sub>2</sub>)<sub>2*n*</sub>CH<sub>3</sub> and C<sub>2</sub>H<sub>6</sub>. The pincer iridium complex dehydrogenates the *n*-alkane to give α-olefin. The olefin metathesis catalyst then effects conversion of two C<sub>*n*</sub> olefins to C<sub>2*n*-2</sub> olefin and ethylene. The dihydride complexes



**Scheme 2** Alkane metathesis of linear alkanes


formed in the initial step then hydrogenate these olefins to afford  $\text{C}_{2n-2}$  *n*-alkane and ethane. Thus, the overall role of the iridium complex is to transfer hydrogen from *n*-alkane to the metathesized olefins.

Using **1**-H<sub>2</sub> or **4a**-H<sub>2</sub> as the transfer-dehydrogenation catalysts, 1.25 and 2.05 M total product concentrations, respectively, were obtained from 7.6 M *n*-hexane after 1 day at 125°C. In accord with the proposal illustrated in Scheme 2, exclusively linear *n*-alkanes were formed (in contrast with the Basset systems). But in contrast with the idealized cycle depicted in Scheme 2, the mixtures were not restricted to  $\text{C}_{2n-2}$  alkane and ethane as products.

Interestingly, the nature of the product mixture was very dependent upon the iridium catalyst. **1**-H<sub>2</sub> gave at least moderate selectivity for *n*-decane as the heavy product from *n*-hexane, while **4a**-H<sub>2</sub> gave successively decreasing amounts of linear heptane, octane, nonane, and decane. A priori, the low selectivity obtained using **4a**-H<sub>2</sub> could be explained by faster isomerization of olefin intermediate prior to olefin metathesis and hydrogenation. Recent experimental evidence, however, indicates that **4a** can directly dehydrogenate *n*-alkane to give the *internal* olefin [57], unlike **1** that shows high kinetic selectivity for the terminal position of *n*-alkanes.

It was observed that turnover numbers for AM using **1** or **4a** and **7** are limited by the decomposition of **7**. The use of the more thermally stable, heterogeneous, olefin-metathesis catalyst  $\text{Re}_2\text{O}_7/\text{Al}_2\text{O}_3$  in place of **7** was therefore investigated for this purpose. The resulting system was less active, requiring higher temperature (175°C) to achieve even rates that were comparable, though still slower. However, catalytic activity was maintained for a longer time and higher turnover numbers were ultimately obtained. In more recent work, using  $\text{Re}_2\text{O}_7/\text{Al}_2\text{O}_3$  and pincer-iridium catalysts supported on  $\text{Al}_2\text{O}_3$  through *para*-substituents (e.g., *t*-Bu<sub>2</sub>PO-), even higher turnover numbers (>2,000) were ultimately achieved [58].

Schrock and coworkers have generated and tested more than forty molybdenum and tungsten imido complexes with pincer iridium complex for AM [59]. In general, tungsten complexes perform better than molybdenum complexes, presumably due to



**Scheme 3** Alkane metathesis of cyclic alkanes

greater stability of the former at the high temperatures required for dehydrogenation. The tungsten imido complex,  $W(NAr)(CHR)(OSiPh_3)_2$  ( $Ar = 2,6$ -diisopropylphenyl,  $R = CMe_2Ph$ ) gave the best product yield, about double that reported with the originally used molybdenum catalyst **7**.


Scott and coworkers [60] have also investigated applications of AM to cycloalkanes. They found, for example, that cyclooctane undergoes metathesis by **7** in tandem with **1**, **4a**, or **6**, to give a range of cyclic products, including (surprisingly) cycloheptane, but mostly low cyclooligomers in which the carbon number is a multiple of 8. The proposed mechanism is shown in Scheme 3.

### 2.3 Dehydrogenation of Alkyl Groups in Molecules Containing Heteroatoms

Examples of iridium-catalyzed dehydrogenation of alkyl moieties (H–C–C–H linkages) in heteroatom-containing molecules are relatively scarce although stoichiometric reactions have been reported in several cases [61, 62]. Jensen, Kaska and coworkers reported that **1-H<sub>2</sub>** catalyzes transfer-dehydrogenation of tetrahydrofuran to produce dihydrofuran as well as furan [63]. The catalytic activity was significantly lower than with cycloalkanes under comparable conditions. It was suggested that the oxygen coordination to the metal center inhibits the reaction. Knapp, Goldman and coworkers reported that dehydrogenation of tertiary amines to give enamines (**7**) occurs in high yield and with good selectivity, particularly in the case of *N*-ethyl amines [64]. The resulting terminal enamines are difficult to synthesize by conventional synthetic routes.

Several iridium-catalyzed dehydrogenations of HCXH ( $X = O, N$ ) linkages have been reported by pincer-iridium complexes. While these dehydrogenations are not generally considered as C–H bond activations or functionalizations, it is generally not known if they proceed via initial activation of the X–H or C–H bond [65, 66].






### 3 Borylation of Arenes

Arylboronate esters are widely used synthons in organic transformations such as C–C or C–X bond forming reaction, thanks to their low toxicity, good functional group tolerance, and high stability toward atmospheric oxidation. Traditionally, they have been prepared from aryl halides via conversion to Grignard or lithium reagents followed by reaction with trialkyl borates. The possibility of direct borylation via C–H bond cleavage was first hinted at by Rablen and Hartwig, who computationally predicted the bond dissociation energies of B–H and B–C bonds and suggested that conversion of C–H to C–B bonds is thermodynamically favorable [67, 68]. Shortly thereafter, the first examples of metal-catalyzed borylations of alkane and arene C–H bonds were discovered, initially photochemical [69–72] and subsequently thermochemical [73, 74]. Since then, numerous stoichiometric and catalytic borylation of C–H bonds have been reported, particularly of arenes [11, 75, 76]. While many transition metal complexes including Rh, Ru, Re, Fe, W, and Pd have been studied in this context, iridium complexes have proven the most effective for the more practical thermocatalytic transformations, affording high reactivity as well as regioselectivity.


Iridium-catalyzed formation of B–C bonds from arene C–H bonds was first reported by Smith and coworkers [73]. They demonstrated that the archetypal C–H activation products,  $\text{Cp}^*\text{Ir}(\text{PMe}_3)(\text{H})(\text{R})$ , could mediate B–C bond formation ( $\text{R} = \text{Ph}$ , cyclohexyl) and were able to effect the catalytic borylation of benzene with HBpin (**8**) to produce  $\text{C}_6\text{H}_5\text{Bpin}$  and  $\text{H}_2$  at  $150^\circ\text{C}$  (**8**).

Subsequent reports showed that a wide range of arene substrates could be borylated, with regioselectivity predominantly dictated by steric factors [77]. In the case of monosubstituted arenes, borylation generally occurs predominantly at the *meta* and *para* positions, with a near-statistical ratio while substitution at the *ortho* position is minor (Scheme 4). In the case of 1,3-disubstituted arenes, borylation occurs exclusively at 5-position. The selectivity was reversed in favor of the *ortho* position in the case of diethylbenzamide, presumably due to chelation [77].




**Scheme 4** Regioselectivity in monosubstituted arenes

Smith also found that catalytic activity is dramatically increased by generating the active catalyst in situ from  $(\text{Ind})\text{Ir}(\text{COD})$  ( $\text{Ind} = \eta^5\text{-C}_9\text{H}_7$ ) in the presence of  $\text{PMe}_3$  or chelating phosphines such as 1,2-bis(diphenylphosphino)ethane (dppe) or 1,2-bis(dimethylphosphino)ethane (dmppe) [76]. Moreover, these systems show remarkable functional group tolerance, allowing, for example, borylation of aryl halides, esters and heteroarenes. The selectivity for C–H over C–X ( $\text{X} = \text{halogen}$ ) bonds in particular allowed the group to develop one-pot biaryl syntheses via borylation/Pd-catalyzed Miyaura-Suzuki coupling (9) [76].




One of the most active and well-studied catalytic borylation systems is that generated from iridium(I) precursors such as  $[\text{Ir}(\text{COD})\text{Cl}]_2$  or  $[\text{Ir}(\text{COD})(\text{OMe})_2]$  and bipyridine type ligands such as 2,2'-bipyridine or 4,4'-di-*tert*-butyl-2,2'-bipyridine (dtbpy). In 2002, Ishiyama, Miyaura, and Hartwig et al. reported that the combination of  $[\text{Ir}(\text{COD})\text{Cl}]_2$  and 2,2'-bipyridine catalyzes arene borylation in the presence of excess arene under mild conditions ( $80^\circ\text{C}$ ). When the catalyst is generated from  $[\text{Ir}(\text{COE})_2\text{Cl}]_2$  and dtbpy, the reaction proceeds even at room temperature [78, 79]. The same groups optimized conditions and found that the combination of  $[\text{Ir}(\text{COD})(\text{OMe})_2]$  and dtbpy (10) is a highly effective catalyst in the borylation of arenes so that reactions can be successfully performed with equimolar ratio of arenes and

$B_2Pin_2$  [79] or  $HBpin$  [80] at room temperature in nonpolar solvents. The reactions tolerate a wide range of functional groups including Br, I,  $CO_2Me$ , and CN.




Extensive studies of kinetics and isotope effects by Hartwig and coworkers support the mechanism shown in Scheme 5 for the Ir(I)/dtbpy catalyzed borylation [81]. In particular, these studies indicate that the iridium(III) trisboryl bipyridine complex (**10**) is the species that activates the arene C–H bond; this is in agreement with DFT calculations by Sakaki et al. predicting the key intermediacy of the trisboryl complex and the seven-coordinated Ir(V) species resulting from C–H addition [82]. C–H addition to Ir(III) was also proposed in the (Ind)Ir(COD)/phosphine-catalyzed borylation by Smith et al. [76].

The iridium-catalyzed direct borylation has been applied to a wide variety of heteroaromatics including thiophene, pyrrole, furan, indole, and pyridine, with good yield under relatively mild conditions [83]. Again, the combination of [Ir(COD)(OMe)<sub>2</sub>] and dtbpy yielded a highly efficient system, catalyzing borylation at room temperature in high yield [84]. In the case of five-membered heteroarenes, the reactions are highly selective for borylation at the  $\alpha$ -position of the heteroaromatic ring. However, the regioselectivity can be controlled by imposing sufficient steric bulk at the heteroatom (e.g., **11**). Interestingly, borylation of six-membered




**Scheme 5** Proposed mechanism for borylation by bipyridine ligated iridium complexes

heteroaromatic rings shows different regioselectivity. Borylation of pyridine occurs at the 3- and 4-position with a statistical ratio and the reaction of quinoline selectively gives the 3-borylated product (12). Although the reason for different regioselectivity was not determined, one possibility proposed by the authors is that the reversible coordination at nitrogen (by either iridium or boron) blocks borylation at the 2-position while perhaps activating the 3-position [83].




The  $[\text{Ir}(\text{OMe})(\text{COD})]_2/\text{dtbpy}$  catalytic system borylates indole selectively at the 2-position (Scheme 6). Smith and coworkers reported that borylation of N-unprotected 2-substituted indoles exclusively occurs at 7-position (Scheme 6) [85, 86]. It has been suggested that nitrogen interaction with the iridium center or possibly the empty p-orbital of boron in a boryl ligand induces the observed regioselectivity. Borylation of other heteroarenes have been reported using the same or similar Ir(I) and bipyridine combination [85, 87–90].

In the borylation of nonheteroaromatic rings, sterics are generally the most important factor influencing regioselectivity so that boronate is generally introduced at the sterically less hindered sites (e.g., *meta* or *para* in monosubstituted



**Scheme 6** Borylation of indoles


arenes). However, it was shown that selective borylation of an *ortho* position is possible with the proper directing group [91]. Using hydrosilyl groups, benzylic silanes, phenols, and anilines have been successfully boronated with high regioselectivity for the *ortho* position (13). Although the mechanism has not yet been fully investigated, evidence supports the formation of an Ir–Si bond in the course of the reaction. Recently, Sawamura and coworkers reported a silica-supported iridium monophosphine catalyst that efficiently catalyzes selective *ortho* borylation of arenes possessing directing groups (e.g., methyl benzoate or chlorobenzene) [92]. The high *ortho* selectivity was attributed to the fact that the catalytically active iridium species possesses only a single phosphine ligand; presumably, this allows the directing group to coordinate while still maintaining the degree of coordinative unsaturation required for catalytic activity (14).



## 4 Silylation of Arenes


In 1982, Curtis reported one of the earliest examples of catalytic functionalization that appeared to proceed via activation of a simple hydrocarbon C–H bond. Pentamethyldisiloxane was observed to undergo dehydrogenative coupling with benzene in the presence of Vaska's complex,  $\text{Ir}(\text{CO})(\text{Cl})(\text{PPh}_3)_2$ , to yield phenyl-pentamethyldisiloxane along with other redistribution products [93]. The reaction required long reaction time at relatively high temperature and gave a variety of redistribution products along with the Ph–Si coupling product in low yield (~20% after 49 days at 100°C); nevertheless, precedent for catalytic formation of aryl–silyl coupling was established. Two decades later, Ishiyama and Miyaura et al. developed an efficient iridium-catalyzed aromatic silylation system [94].  $[\text{Ir}(\text{COD})(\text{OMe})_2]$  and dtbpy, which also comprises an efficient system for aromatic borylation, catalyzes aromatic silylation with fluorinated disilanes,  $(t\text{-BuF}_2\text{Si})_2$  (11), to give the corresponding aryl difluorosilanes (15). Steric factors appear to influence the regioselectivity of the coupling, similar to borylation of arenes, but the silylations in some cases afford lower regioselectivity. For example, *o*-xylene

gives exclusively 4-silylated products, but silylation of 3-chlorotoluene (**12**) gave a mixture of products (**13** and **14**) (16). The nature of the substituents on silicon is critical;  $({}^t\text{BuCl}_2\text{Si})_2$  or  $(\text{Me}_3\text{Si})_2$  do not undergo the analogous reaction. Subsequently, the same group found significant improvement in regioselectivity by using 2,9-diisopropyl-1,10-phenanthroline (**16**) as the chelating ligand, and 1,2-di-*sec*-butyl-1,1,2,2-tetrafluorodisilane (**15**) as the silyl coupling substrate (**17**) [95].




The silylation of heteroarenes was also reported [96]. Silylation of thiophene or furan with **11** occurs selectively at the  $\alpha$ -position in the presence of  $[\text{Ir}(\text{COD})(\text{OMe})_2]/2$ -*tert*-butyl-1,10-phenanthroline (tbphen, **17**). Silylation of pyrrole or indole under the same conditions was unsuccessful presumably due to the presence of the acidic N–H bond. Accordingly, N-substituted pyrrole and indole undergo silylation with **11**, to selectively give 3-substituted products (Scheme 7).

Falck has recently reported dehydrogenative silylation of heteroarenes with triethylsilane (**18**) [97]. Coupling with the Si–H bond of triethylsilane, rather than the disilane Si–Si bond, in conjunction with the use of norbornene that presumably acts as a hydrogen acceptor, gives good yields with indoles, thiophenes, and furans, under relatively mild condition ( $80^\circ\text{C}$ ). Unlike the reaction shown in Scheme 7, silylation of indole did not require protection of the N–H group.




**Scheme 7** Regioselectivity in silylation of heteroarenes



## 5 C–C Bond Formation


### 5.1 Hydroarylation and Hydrovinylation of C–C Multiple Bonds

Periana and coworkers reported in 2000 that the binuclear iridium (III) complex  $[\text{Ir}(\mu\text{-acac-O,O,C}^3)(\text{acac-O,O})(\text{acac-C}^3)]_2$  (**18**) catalyzes olefin hydroarylation with benzene and unactivated alkenes [98]. In the reaction of 1-alkenes, weak selectivity for anti-Markovnikov addition to yield *n*-alkyl arenes is observed, in sharp contrast with Friedel–Crafts arene alkylations, which essentially only afford branched products. For example, iridium-catalyzed alkylation of benzene with propylene produces *n*-propylbenzene (**20**) and cumene (**21**) in 69 and 31% yield, respectively, whereas the  $\text{AlCl}_3$ -catalyzed reaction gives entirely cumene (**21**) (19). Unlike Friedel–Crafts reactions, in which ethylene reacts more slowly than substituted olefins due to lesser stability of the respective carbonium ion, ethylene is more reactive than propene for the iridium-catalyzed hydroarylation. The reaction is sensitive to steric effects so that alkylation of toluene with ethylene gives only *meta* and *para* substituted products in an approximately statistical ratio (63:37). No *ortho* product was observed. These observations constitute compelling evidence that **18**-catalyzed hydroarylation does not involve free carbonium ions, and are consistent with a C–H activation mechanism. Monomeric iridium complexes,  $\text{Ir}(\text{acac-O,O})_2(\text{R})(\text{L})$  (**19**) where R is Ph or  $(\text{acac-C}^3)$  and L is pyridine or  $\text{H}_2\text{O}$ , were also reported to catalyze the reactions [99]. The use of dimer **18** or four different monomers **R-19-L** (including R = Ph or  $\text{acac-C}^3$  and L = pyridine or  $\text{H}_2\text{O}$ ) all gives comparable turnover frequencies and, most importantly, identical ratios of linear to branched product for reaction 19, thus strongly indicating that these species all give rise to the same catalytically active intermediate [99]. Kinetic studies indicate that the active catalyst is monomeric [99–102].




When **acac-C<sup>3</sup>-19-Py** is used as the precatalyst for reaction 19, the iridium containing species recovered (65% yield) is **Ph-19-Py**. Displacement of pyridine by olefin can presumably generate **22**. Based on extensive experimental and theoretical study by Periana and Goddard, the mechanism proposed for the hydroarylation is that shown in Scheme 8. *Cis*-bis(acac-O,O) Ir complex (**22**) undergoes olefin insertion into the Ir–aryl bond, followed by arene coordination to generate **23a** and **23b**. Aromatic C–H bond activation then takes place via “oxidative hydrogen migration”, a step which may be viewed as proceeding via a transition state – but not an intermediate – that is essentially Ir(V) in character [100–103]. This step, followed by olefin coordination, yields the coupled organic product and regenerates **22**. DFT calculations show that regioselectivity in olefin arylation is determined by the relative energetics of the transition states for formation of **23a** and **23b**.

In 2005, the same authors reported that the iridium(III) complex with tropolonato ligands (**25**) shows faster C–H bond activation rates than the analogs with acetato ligands (**20**) [104]. For example, **25-Me** catalyzes H/D exchange with a mixture of C<sub>6</sub>H<sub>6</sub>/ toluene-*d*<sub>8</sub> 50 times faster than the acetato analog. However, hydroarylation catalyzed by **25-Ph** is slightly less effective than with the acetato analog [105].




Periana and Goddard et al. also found that the iridium(III) vinyl bis-acetylacetonate complex (**Vinyl-Ir-Py**, **26**) catalyzes dimerization of olefins (e.g., **21**) [106, 107]. Analogously to the hydroarylation of Scheme 8, the mechanism is proposed to






**Scheme 8** Proposed mechanism for hydroarylation


involve insertion of olefin into an Ir–vinyl bond followed by vinyl C–H bond activation (Scheme 9).




Several arylations involving reactive alkenes such as norbornene or allenes have been reported. Togni and coworkers have shown that norbornene is selectively added to the *ortho* positions of phenols to produce a mixture of **30** and **31** in 69% and 13% yield, respectively, after 72 hours at 100°C (22) [108, 109]. 1,1-dimethylallene also reacts with aromatic carboxamides (**33**) to produce prenylation products (**34**) in the presence of cationic iridium complexes (**23**) [110]. In both cases, initial *ortho* C–H bond activation in arenes directed by coordinating groups followed by olefin insertion has been proposed.



**Scheme 9** Proposed mechanism of hydrovinylation




Insertion of alkynes into aromatic C–H bonds has been achieved by iridium complexes. Shibata and coworkers found that the cationic complex  $[\text{Ir}(\text{COD})_2]\text{BF}_4$  catalyzes the hydroarylation of internal alkynes with aryl ketones in the presence of BINAP (**24**) [111]. The reaction selectively produces *ortho*-substituted alkenated-aryl products. Styrene and norbornene were also found to undergo hydroarylation under similar condition.  $[\text{Cp}^*\text{IrCl}_2]_2$  catalyzes aromatization of benzoic acid with two equivalents of internal alkyne to form naphthalene derivatives via decarboxylation in the presence of  $\text{Ag}_2\text{CO}_3$  as an oxidant (**25**) [112].




## 5.2 Aryl–Aryl Coupling Reactions

Relatively few examples have been reported of iridium-catalyzed arylation of arenes via C–H bond cleavage. The scope of the reactions has been, to date, limited to iodoarenes as coupling partners. The Yamaguchi group reported Cp\*Ir-catalyzed benzene coupling with iodoarenes in the presence of excess of KO<sup>t</sup>Bu (26) [113]. Various types of methyl or methoxy substituted iodoarenes undergo arylation with benzene in moderate yield. The reaction of anisole or toluene with iodobenzene gives poor regioselectivity. These limitations notwithstanding, this was a notable example of arylation of an aryl C–H bond that did not require any “directing group” adjacent to the C–H bond substrate. The mechanism is proposed to involve a phenyl radical intermediate based on the following (1) The regioselectivity shown in 27 is similar to that observed in radical substitution reactions. (2) In the presence of radical scavenger, the authors observed a significant decrease in reaction yield [113].




The report by Yamaguchi et al. revealed the first example of iridium-catalyzed direct arylation of aromatic C–H bonds. Subsequently, the Itami group reported that [Ir(COD)(py)PCy<sub>3</sub>]PF<sub>6</sub>, Crabtree’s catalyst [114], mediates direct arylation of a broad range of heteroarenes including thiophene and furan in good yield (28) [115]. The choice of additive turns out to be crucial in that only Ag<sub>2</sub>CO<sub>3</sub> promotes the reaction among those investigated including AgOTf, AgOAc, Na<sub>2</sub>CO<sub>3</sub>, K<sub>2</sub>CO<sub>3</sub>, and KO<sup>t</sup>Bu. The system shows good functional group tolerance; bromo, nitro, and ester group substituted iodoarenes were used as coupling partners.




### 5.3 Coupling with $sp^3$ C–H Bonds

Examples of catalytic formation of C–C bonds from  $sp^3$  C–H bonds are even more scarce than from  $sp^2$  C–H bonds and, in general, are limited to C–H bonds adjacent to heteroatoms. A remarkable iridium-catalyzed example was reported by the group of Lin [116] the intermolecular oxidative coupling of methyl ethers with TBE to form olefin complexes in the presence of  $(P^iPr_3)_2IrH_5$  (**29**). In their proposed mechanism, the reactive  $14e^-$  species **38** undergoes oxidative addition of the methyl C–H bond in methyl ethers followed by olefin insertion to generate the intermediate **39**.  $\beta$ -hydride elimination affords **35**, which can isomerize to products **36** and **37** (Scheme 10). The reaction proceeds under mild condition ( $50^\circ C$ ) but suffers from poor selectivity as well as low yield (TON of 12 after 24 h).




Sames et al. have reported the intramolecular cyclization of alkene–amide substrates catalyzed by  $[\text{Ir}(\text{COE})_2\text{Cl}]_2$  and the *N*-heterocyclic carbene ligand, *N,N'*-bis-(2,6-diisopropylphenyl)-imidazolyl, via olefin insertion following oxidative addition of an  $sp^3$  C–H bond (Scheme 11) [117].

Recently, Shibita et al. reported catalysis of alkyne insertion into an arylamide  $sp^3$  C–H bond to give allylamides (**42**) by a cationic iridium complex [118]. An interesting aspect of this work is the unusually selective cleavage of an  $sp^3$  C–H bond over  $sp^2$  aromatic C–H bonds so that the alkenyl arylamide (**43**) is only a very minor product (**30**). The carbonyl group is required for the reaction as no coupling




**Scheme 10** Proposed mechanism for C–C bond coupling via  $sp^3$  C–H bond cleavage



**Scheme 11** Cyclization of alkene–amide via  $\text{sp}^3$  C–H Bond cleavage

was observed with *N,N*-dimethylbenzylamine under the same conditions. This was tentatively proposed to be due to carbonyl coordination directing C–H addition of the amido  $\text{sp}^3$  C–H bond (although the same functionality is well known to result in functionalization of *ortho*-aryl-C–H bonds [119]). Alkyne insertion presumably follows C–H addition to give the allylamides.



In an unusual example of C–C coupling, the benzylation of a bromofullerene, C<sub>60</sub>Me<sub>3</sub>Br, with toluene, has been reported to be catalyzed by [Ir(COE)<sub>2</sub>Cl]<sub>2</sub> in the presence of triethylamine [120].


## 6 Others

### 6.1 $\alpha,\alpha$ -Dehydrogenation and Reactions with Heterocumulenes


Initiating a remarkable series of papers in 2008–2009, Grubbs and Whited reported the formation of an Ir(I) carbene complex, **44**, from the reaction of methyl *t*-butyl ether (MTBE) with Ozerov's (PNP)IrH<sub>2</sub> (PNP = [N(2-*i*PrPr<sub>2</sub>-4-Me-C<sub>6</sub>H<sub>3</sub>)<sub>2</sub>]<sup>−</sup>) and NBE [121–127]. C–H addition to (PNP)Ir presumably occurs followed by

$\alpha$ -H-elimination and loss of  $\text{H}_2$ , which is scavenged by a second mol of NBE; the overall result can be considered an  $\alpha,\alpha$ -dehydrogenation. Carbene complex **44**, although formally a Fischer carbene, does not display typical electrophilic activity at the carbene carbon. Rather, the characteristic reactivity elucidated of **44** was with heterocumulenes, particularly  $\text{CO}_2$  to give *t*-Bu formate (**45**) and  $(\text{PNP})\text{Ir}(\text{CO})$  (**47**) (Scheme 12). Analogous reactions were found with carbonyl sulfide and phenyl isocyanate to give **47** and the thioester and formimidate **46**. Attempts to incorporate these reactions into catalytic cycles were not successful, no doubt due to the very strong Ir–CO bond [128] in **47**.

In order to avoid formation of **47**, the use of diazo reagents ( $\text{E}=\text{N}=\text{N}$ ) was investigated in place of the analogous carbonyls and indeed the organic products **45** was produced from  $\text{N}_2\text{O}$ , along with the dinitrogen complex **48**. However, even dinitrogen proved too strongly bound to allow incorporation into a thermal catalytic cycle. Dinitrogen could be extruded photochemically, however, allowing the authors to achieve the photocatalytic formation of *N*-adamantyl *t*-butylformimidate shown in Scheme 13 [121–127].



**Scheme 12** Reactions  $(\text{PNP})\text{Ir}=\text{CH}(\text{O}^t\text{Bu})$  with heterocumulenes




**Scheme 13** Catalytic cycle for *t*-butylformimidate synthesis

## 6.2 C–H Bond Activation by Metal Carbenoids or Nitrenoids

Functionalization of C–H bonds by metal carbenoid or nitrenoid insertions represents a promising alternative to the more “traditional” approach of direct activation by a metal center. Carbenoids and nitrenoids show unusual regio- and stereoselectivity in insertions into C–H bonds, and unlike insertions of metal centers, these are intrinsically “functionalizations” rather than “activations,” which must be followed by functionalization (although in either case, loss of the functionalized group, to regenerate the active metal complex, is still required for catalysis) [129]. The use of dimeric Rh(II) complexes in this area has been extensively developed [129].

Recent studies show that iridium complexes are also capable of catalyzing C–C or C–N bond coupling via carbenoid or nitrenoid insertion into C–H bonds. Katsuki et al. reported that iridium(III)–salen complexes catalyze asymmetric C–H insertion of carbenes at the  $\alpha$ -position of THF and the allylic position of 1,4-cyclohexadiene [130]. In both cases, the reactions are highly enantioselective and in the case of THF, highly diastereoselective (Scheme 14). Importantly, while previous related work was limited to the use of  $\alpha$ -aryl- $\alpha$ -diazoacetates, this study gave comparably good results with simple  $\alpha$ -methyl- $\alpha$ -diazoacetate (R=Me, Scheme 14); this may be due to inhibition of  $\beta$ -H-elimination as a result of the salen ligand precluding access to a coordination site cis to the site of the carbene.

Intramolecular C–N bond coupling in arylazides via C–H bond activation catalyzed by [Ir(COD)(OMe)<sub>2</sub>] was recently reported [131]. The intermediate iridium nitrenoid complex (**50**) formed after the extrusion of dinitrogen is proposed as a reactive species, which can cleave the benzylic C–H bond to yield indoline derivatives (**31**).



**Scheme 14** Stereoselective C–H functionalization by Ir(III)–salen

## 7 Concluding Remarks

Extensive research over the past two decades on the catalytic functionalization of C–H bonds by complexes of iridium, as with other transition metals, has begun to pay significant dividends. Nevertheless, only a small fraction of the great potential offered by such chemistry has been realized, and there seems little doubt that progress will continue at a steady pace at least.

Iridium has a tendency to form very stable – and catalytically inactive – Ir(III) complexes. The metal, however, has proven particularly effective for some hydrocarbon functionalizations, the first of which to be discovered was alkane dehydrogenation. Key to this reactivity is (a) the ability of three-coordinate Ir(I) to undergo facile oxidative addition of alkyl–H bonds, an addition that is generally quite unfavorable thermodynamically, and (b) the fact that the products of such additions are coordinatively unsaturated and therefore able to undergo  $\beta$ -hydrogen elimination. The other well-developed C–H bond functionalization in which iridium plays a leading role is borylation. In contrast with dehydrogenation, the key oxidative couple for borylation, and perhaps silylation, arylation, and other couplings as well, appears to be Ir(III)/Ir(V). These reactions involve bonds much more prone to oxidative addition than alkyl–H, including aryl–H, B–H, B–B, and Si–H (For some relevant examples of Ir(V) complexes and discussions of factors stabilizing them, see [132–138]). This may explain why the Ir(III)/Ir(V) couple is accessible with these substrates, although even with such substrates the Ir(V) oxidation state is certainly not so stable that catalytic activity is precluded. None of this however is understood definitively, and any generalizations are still highly speculative. Thus, although iridium provides some of the most classic examples in mechanistic organometallic chemistry, there seems to be great room for growth in its applications in C–H bond functionalization, which will probably be coupled with commensurate growth in our mechanistic understanding of this chemistry.

## References

1. Crabtree RH (2001) *Dalton Trans* 17 2437–2450
2. Labinger JA, Bercaw JE (2002) *Nature* 417:507–514
3. Ritleng V, Sirlin C, Pfeffer M (2002) *Chem Rev* 102:1731–1769
4. Goldman AS, Goldberg KI (2004) In: Goldberg KI, Goldman AS (eds) *Activation and functionalization of C–H bonds*, ACS symposium series, vol 885, pp 1–43
5. Colby DA, Bergman RG, Ellman JA (2010) *Chem Rev* 110:624–655
6. Balcells D, Clot E, Eisenstein O (2010) *Chem Rev* 110:749–823
7. Dobereiner GE, Crabtree RH (2010) *Chem Rev* 110:681–703
8. Bellina F, Rossi R (2010) *Chem Rev* 110:1082–1146
9. Gunay A, Theopold KH (2010) *Chem Rev* 110:1060–1081
10. Lyons TW, Sanford MS (2010) *Chem Rev* 110:1147–1169
11. Mkhalid IAI, Barnard JH, Marder TB, Murphy JM, Hartwig JF (2010) *Chem Rev* 110:890–931



12. Vaska L, DiLuzio JW (1962) *J Am Chem Soc* 84:679–680
13. Vaska L (1968) *Accounts Chem Res* 1:335–344
14. Janowicz AH, Bergman RG (1982) *J Am Chem Soc* 104:352–354
15. Crabtree RH, Mihelcic JM, Quirk JM (1979) *J Am Chem Soc* 101:7738–7739
16. Oro LA, Claver C (eds) (2009) *Iridium complexes in organic synthesis*. Wiley-VCH, Weinheim
17. Burk MJ, Crabtree RH, Parnell CP, Uriarte RJ (1984) *Organometallics* 3:816–817
18. Felkin H, Fillebeen-Khan T, Gault Y, Holmes-Smith R, Zakrzewski J (1984) *Tetrahedron Lett* 25:1279–1282
19. Baudry D, Ephritikhine M, Felkin H, Zakrzewski J (1984) *Tetrahedron Lett* 25:1283–1286
20. Felkin H, Fillebeen-Khan T, Holmes-Smith R, Lin Y (1985) *Tetrahedron Lett* 26:1999–2000
21. Burk MJ, Crabtree RH, McGrath DV (1985) *J Chem Soc Chem Commun* 1829–1830
22. Burk MJ, Crabtree RH (1987) *J Am Chem Soc* 109:8025–8032
23. Kunin AJ, Eisenberg R (1986) *J Am Chem Soc* 108:535–536
24. Kunin AJ, Eisenberg R (1988) *Organometallics* 7:2124–2129
25. Sakakura T, Tanaka M (1987) *J Chem Soc Chem Commun* 758–759
26. Sakakura T, Tanaka M (1987) *Chem Lett* 249–252
27. Sakakura T, Sodeyama T, Tanaka M (1989) *New J Chem* 13:737–745
28. Sakakura T, Sodeyama T, Tokunaga Y, Tanaka M (1988) *Chem Lett* 263–264
29. Nomura K, Saito Y (1988) *J Chem Soc Chem Commun* 161
30. Maguire JA, Boese WT, Goldman AS (1989) *J Am Chem Soc* 111:7088–7093
31. Maguire JA, Goldman AS (1991) *J Am Chem Soc* 113:6706–6708
32. Maguire JA, Petrillo A, Goldman AS (1992) *J Am Chem Soc* 114:9492–9498
33. Gupta M, Hagen C, Flesher RJ, Kaska WC, Jensen CM (1996) *Chem Commun* 2083–2084
34. Gupta M, Hagen C, Kaska WC, Cramer RE, Jensen CM (1997) *J Am Chem Soc* 119:840–841
35. Xu W, Rosini GP, Gupta M, Jensen CM, Kaska WC, Krogh-Jespersen K, Goldman AS (1997) *Chem Commun* 2273–2274
36. Liu F, Goldman AS (1999) *Chem Commun* 655–656
37. Liu F, Pak EB, Singh B, Jensen CM, Goldman ASJ (1999) *J Am Chem Soc* 121:4086–4087
38. Renkema KB, Kissin YV, Goldman AS (2003) *J Am Chem Soc* 125:7770–7771
39. Krogh-Jespersen K, Czerw M, Kanzelberger M, Goldman AS (2001) *J Chem Inf Comput Sci* 41:56–63
40. Kundu S, Choliy Y, Zhuo G, Ahuja R, Emge TJ, Warmuth R, Brookhart M, Krogh-Jespersen K, Goldman AS (2009) *Organometallics* 28:5432–5444
41. Biswas S, Ahuja R, Ray A, Choliy Y, Krogh-Jespersen K, Brookhart M, Goldman AS (2008) Abstracts of papers, 236th ACS national meeting, Philadelphia, PA, 17–21 August 2008, INOR-125
42. Haenel MW, Oevers S, Angermund K, Kaska WC, Fan H-J, Hall MB (2001) *Angew Chem Intl Ed* 40:3596–3600
43. Krogh-Jespersen K, Czerw M, Zhu K, Singh B, Kanzelberger M, Darji N, Achord PD, Renkema KB, Goldman AS (2002) *J Am Chem Soc* 124:10797–10809
44. Zhu K, Achord PD, Zhang X, Krogh-Jespersen K, Goldman AS (2004) *J Am Chem Soc* 126:13044–13053
45. Göttker-Schnetmann I, White P, Brookhart M (2004) *J Am Chem Soc* 126:1804–1811
46. Göttker-Schnetmann I, White PS, Brookhart M (2004) *Organometallics* 23:1766–1776
47. Göttker-Schnetmann I, Brookhart M (2004) *J Am Chem Soc* 126:9330–9338
48. Morales-Morales D, Redon R, Yung C, Jensen CM (2004) *Inorg Chim Acta* 357:2953–2956
49. Ray A, Zhu K, Kissin YV, Cherian AE, Coates GW, Goldman AS (2005) *Chem Commun* 3388–3390
50. Huang Z, Brookhart M, Goldman AS, Kundu S, Ray A, Scott SL, Vicente BC (2009) *Adv Synth Catal* 351:188–206

51. Dry ME (2002) *Catalysis Today* 71:227–241
52. Schulz H (1999) *Appl Catal A* 186:3–12
53. Burnett RL, Hughes TR (1973) *J Catal* 31:55–64
54. Vidal V, Theolier A, Thivolle-Cazat J, Basset J-M (1997) *Science* 276:99–102
55. Basset JM, Coperet C, Lefort L, Maunders BM, Maury O, Le Roux E, Saggio G, Soignier S, Soulivong D, Sunley GJ, Taoufik M, Thivolle-Cazat J (2005) *J Am Chem Soc* 127:8604–8605
56. Goldman AS, Roy AH, Huang Z, Ahuja R, Schinski W, Brookhart M (2006) *Science* 312:257–261
57. Biswas S, Choliy Y, Krogh-Jespersen K, Brookhart M, Goldman AS (2009) Abstract of papers, 238th ACS national meeting, Washington, DC, 16–20 August 2009, INOR-479
58. Huang Z, Rolfe E, Carson EC, Brookhart M, Goldman AS, El-Khalafy SH, MacArthur AHR (2010) *Adv Synth Catal* 352:125–135
59. Bailey BC, Schrock RR, Kundu S, Goldman AS, Huang Z, Brookhart M (2009) *Organometallics* 28:355–360
60. Ahuja R, Kundu S, Goldman AS, Brookhart M, Vicente BC, Scott SL (2008) *Chem Commun* 253–255
61. Luecke HF, Arndtsen BA, Burger P, Bergman RG (1996) *J Am Chem Soc* 118:2517–2518
62. Whited MT, Zhu Y, Timpa SD, Chen C-H, Foxman BM, Ozerov OV, Grubbs RH (2009) *Organometallics* 28:4560–4570
63. Gupta M, Kaska WC, Jensen CM (1997) *Chem Commun* 461–462
64. Zhang X, Fried A, Knapp S, Goldman AS (2003) *Chem Commun* 2060–2061
65. Gu X-Q, Chen W, Morales-Morales D, Jensen CM (2002) *J Mol Cat A* 189:119–124
66. Bernskoetter WH, Brookhart M (2008) *Organometallics* 27:2036–2045
67. Rablen PR, Hartwig JF, Nolan SP (1994) *J Am Chem Soc* 116:4121–4122
68. Rablen PR, Hartwig JF (1996) *J Am Chem Soc* 118:4648–4653
69. Waltz KM, He X, Muhoro C, Hartwig JF (1995) *J Am Chem Soc* 117:11357–11358
70. Waltz KM, Hartwig JF (1997) *Science* 277:211–213
71. Chen H, Hartwig JF (1999) *Angew Chem Intl Ed* 38:3391–3393
72. Waltz KM, Hartwig JF (2000) *J Am Chem Soc* 122:11358–11369
73. Iverson CN, Smith MR III (1999) *J Am Chem Soc* 121:7696–7697
74. Chen H, Schlecht S, Semple TC, Hartwig JF (2000) *Science* 287:1995–1997
75. Ishiyama T, Miyaura N (2006) *Pure Appl Chem* 78:1369–1375
76. Cho J-Y, Tse MK, Holmes D, Maleczka RE Jr, Smith MR III (2002) *Science* 295:305–308
77. Cho J-Y, Iverson CN, Smith MR III (2000) *J Am Chem Soc* 122:12868–12869
78. Ishiyama T, Takagi J, Ishida K, Miyaura N, Anastasi NR, Hartwig JF (2002) *J Am Chem Soc* 124:390–391
79. Ishiyama T, Takagi J, Hartwig JF, Miyaura N (2002) *Angew Chem Intl Ed* 41:3056–3058
80. Ishiyama T, Nobuta Y, Hartwig JF, Miyaura N (2003) *Chem Commun* 2924–2925
81. Bollner TM, Murphy JM, Hapke M, Ishiyama T, Miyaura N, Hartwig JF (2005) *J Am Chem Soc* 127:14263–14278
82. Tamura H, Yamazaki H, Sato H, Sakaki S (2003) *J Am Chem Soc* 125:16114–16126
83. Takagi J, Sato K, Hartwig JF, Ishiyama T, Miyaura N (2002) *Tetrahedron Lett* 43:5649–5651
84. Ishiyama T, Takagi J, Yonekawa Y, Hartwig JF, Miyaura N (2003) *Adv Synth Catal* 345:1103–1106
85. Paul S, Chotana GA, Holmes D, Reichle RC, Maleczka RE Jr, Smith MR III (2006) *J Am Chem Soc* 128:15552–15553
86. Lo WF, Kaiser HM, Spannenberg A, Beller M, Tse MK (2006) *Tetrahedron Lett* 48:371–375
87. Mkhalid IAI, Conventry DN, Albesa-Jove D, Batsanov AS, Howard JAK, Perutz RN, Marder TB (2006) *Angew Chem Intl Ed* 45:489–491
88. Mertins K, Zapf A, Beller M (2004) *J Mol Cat A* 207:21–25
89. Chotana GA, Kallepalli VA, Maleczka RE Jr, Smith MR III (2008) *Tetrahedron* 64:6103–6114

90. Kallepalli VA, Shi F, Paul S, Onyeozili EN, Maleczka RE, Smith MR (2009) *J Org Chem* 74:9199–9201
91. Boebel TA, Hartwig JF (2008) *J Am Chem Soc* 130:7534–7535
92. Kawamorita S, Ohmiya H, Hara K, Fukuoka A, Sawamura M (2009) *J Am Chem Soc* 131:5058–5059
93. Gustavson WA, Epstein PS, Curtis MD (1982) *Organometallics* 1:884–885
94. Ishiyama T, Sato K, Nishio Y, Miyaura N (2003) *Angew Chem Intl Ed* 42:5346–5348
95. Saiki T, Nishio Y, Ishiyama T, Miyaura N (2006) *Organometallics* 25:6068–6073
96. Ishiyama T, Sato K, Nishio Y, Saiki T, Miyaura N (2005) *Chem Commun* 5065–5067
97. Lu B, Falck JR (2008) *Angew Chem Intl Ed* 47:7508–7510
98. Matsumoto T, Taube DJ, Periana RA, Taube H, Yoshida H (2000) *J Am Chem Soc* 122:7414–7415
99. Periana RA, Liu XY, Bhalla G (2002) *Chem Commun* 3000–3001
100. Oxgaard J, Muller RP, Goddard WA, Periana RA (2004) *J Am Chem Soc* 126:352–363
101. Oxgaard J, Periana RA, Goddard WA III (2004) *J Am Chem Soc* 126:11658–11665
102. Bhalla G, Liu XY, Wong-Foy A, Jones CJ, Periana RA (2004) In: *Activation and functionalization of C-H bonds*, ACS symposium series, vol 885, pp 105–115
103. Ghosh R, Zhang X, Achord P, Emge TJ, Krogh-Jespersen K, Goldman AS (2007) *J Am Chem Soc* 129:853–866
104. Bhalla G, Periana RA (2005) *Angew Chem Intl Ed* 44:1540–1543
105. Bhalla G, Oxgaard J, Goddard WA III, Periana RA (2005) *Organometallics* 24:3229–3232
106. Bhalla G, Oxgaard J, Goddard WA II, Periana RA (2005) *Organometallics* 24:5499–5502
107. Oxgaard J, Bhalla G, Periana RA, Goddard WA III (2006) *Organometallics* 25:1618–1625
108. Dorta R, Tognib A (2003) *Chem Commun* 760–761
109. Aufdenblatten R, Diezi S, Togni A (2000) *Monatsh Chem* 131:1345–1350
110. Zhang YJ, Skucas E, Krische MJ (2009) *Org Lett* 11:4248–4250
111. Tsuchikama K, Kasagawa M, Hashimoto Y-K, Endo K, Shibata T (2008) *J Organomet Chem* 693:3939–3942
112. Ueura K, Satoh T, Miura M (2007) *J Org Chem* 72:5362–5367
113. Fujita K-i, Nonogawa M, Yamaguchi R (2004) *Chem Commun* 1926–1927
114. Crabtree RH, Felkin H, Morris GE (1977) *J Organomet Chem* 141:205–215
115. Join B, Yamamoto T, Itami K (2009) *Angew Chem Intl Ed* 48:3644–3647, S3644/3641-S3644/3679
116. Lin Y, Ma D, Lu X (1987) *Tetrahedron Lett* 28:3249–3252
117. DeBoef B, Pastine SJ, Sames D (2004) *J Am Chem Soc* 126:6556–6557
118. Tsuchikama K, Kasagawa M, Endo K, Shibata T (2009) *Org Lett* 11:1821–1823
119. Kakiuchi F, Murai S (2002) *Acc Chem Res* 35:826–834
120. Matsuo Y, Iwashita A, Nakamura E (2006) *Chem Lett* 35:858–859
121. Whited MT, Grubbs RH (2008) *Organometallics* 27:5737–5740
122. Whited MT, Grubbs RH (2008) *J Am Chem Soc* 130:5874–5875
123. Romero PE, Whited MT, Grubbs RH (2008) *Organometallics* 27:3422–3429
124. Whited MT, Grubbs RH (2008) *J Am Chem Soc* 130:16476–16477
125. Whited MT, Grubbs RH (2009) *Organometallics* 28:161–166
126. Whited MT, Zhu Y, Timpa SD, Chen C-H, Foxman BM, Ozerov OV, Grubbs RH (2009) *Organometallics* 28:4560–4570
127. Whited MT, Grubbs RH (2009) *Acc Chem Res* 42:1607–1616
128. Rosini GP, Liu F, Krogh-Jespersen K, Goldman AS, Li C, Nolan SP (1998) *J Am Chem Soc* 120:9256–9266
129. Davies HML, Manning JR (2008) *Nature* 451:417–424
130. Suematsu H, Katsuki T (2009) *J Am Chem Soc* 131:14218–14219
131. Sun K, Sachwani R, Richert KJ, Driver TG (2009) *Org Lett* 11:3598–3601
132. Klei SR, Tilley TD, Bergman RG (2000) *J Am Chem Soc* 122:1816–1817
133. Kawamura K, Hartwig JF (2001) *J Am Chem Soc* 123:8422–8423

134. Mohammad HAY, Grimm JC, Eichele K, Mack H-G, Speiser B, Novak F, Quintanilla MG, Kaska WC, Mayer HA (2002) *Organometallics* 21:5775–5784
135. Krogh-Jespersen K, Czerw M, Summa N, Renkema KB, Achord PD, Goldman AS (2002) *J Am Chem Soc* 124:11404–11416
136. Webster CE, Hall MB (2003) *Coord Chem Rev* 238–239:315–331
137. Lam WH, Lam KC, Lin Z, Shimada S, Perutz RN, Marder TB (2004) *Dalton Trans* 1556–1562
138. Bernskoetter WH, Lobkovsky E, Chirik PJ (2005) *Organometallics* 24:4367–4373



# Iridium-Catalyzed Allylic Substitution

John F. Hartwig and Mark J. Pouy

**Abstract** Iridium-catalyzed asymmetric allylic substitution has become a valuable method to prepare products from the addition of nucleophiles at the more substituted carbon of an allyl unit. The most active and selective catalysts contain a phosphoramidite ligand possessing at least one aryethyl substituent on the nitrogen atom of the ligand. In these systems, the active catalyst is generated by a base-induced cyclometalation at the methyl group of this substituent to generate an iridium metalacycle bound by the COD ligand of the  $[\text{Ir}(\text{COD})\text{Cl}]_2$  precursor and one additional labile dative ligand. Such complexes catalyze the reactions of linear allylic esters with alkylamines, arylamines, phenols, alcohols, imides, carbamates, ammonia, enolates and enolate equivalents, as well as typical stabilized carbon nucleophiles generated from malonates and cyanoesters. Iridium catalysts for enantioselective allylic substitution have also been generated from phosphorus ligands with substituents bound by heteroatoms, and an account of the studies of such systems, along with a description of the development of iridium catalysts is included.

**Keywords** Allylic substitution · Asymmetric catalysis · Heterocycles · Iridium · Phosphoramidite

## Contents

1	Introduction .....	170
2	Background .....	171
2.1	An Overview of Allylic Substitution .....	171
2.2	Nucleophilic Attack on $\eta^3$ -Allyl Metal Complexes .....	171
2.3	Early Metal-Catalyzed Allylic Substitution .....	172
2.4	The State of Allylic Substitution in 1997 .....	173
3	Early Iridium-Catalyzed Allylic Substitutions .....	174
3.1	Allylic Substitution Catalyzed by $[\text{Ir}(\text{COD})\text{Cl}]_2$ and Triphenylphosphite .....	174

---

J.F. Hartwig (✉) and M.J. Pouy

Department of Chemistry, University of Illinois, 600 South Mathews Avenue, Urbana,  
IL 61801, USA

e-mail: jhartwig@illinois.edu


3.2	Asymmetric Iridium-Catalyzed Allylic Substitution: A Survey of Ligands	177
3.3	Iridium Catalysts Containing Phosphoramidite Ligands	181
4	Advances in Mechanistic Understanding and Catalyst Design: Cyclometalation and Modification of the Phosphoramidite Ligand	184
4.1	Discovery of the Active Metalacyclic Catalyst	184
4.2	Effect of the Phosphoramidite Ligand	185
4.3	Scope of Allylic Substitution Catalyzed by Metalacyclic Iridium-Phosphoramidite Complexes	187
5	New Mechanistic Insights: The Isolation of Catalytic Intermediates	195
5.1	General Mechanism of Metal-Catalyzed Allylic Substitution	195
5.2	Isolation and Study of Intermediates in Allylic Substitution Reactions Catalyzed by Metalacyclic Iridium-Phosphoramidite Complexes	196
5.3	Reaction Development Based on Mechanistic Insights	198
6	Challenges in Iridium-Catalyzed Allylic Substitution	201
6.1	Allylic Substitution of Allylic Alcohols	201
6.2	Diastereoselectivity of Iridium-Catalyzed Allylic Substitution Reactions	202
6.3	Allylic Substitution of Branched Allylic Esters	203
6.4	Allylic Substitution with Highly Substituted Allylic Esters	204
6.5	Stability of Iridium-Phosphoramidite Catalysts to Air and Water	205
7	Conclusion	205
	References	206

## 1 Introduction

A wide range of carbon, nitrogen, and oxygen nucleophiles react with allylic esters in the presence of iridium catalysts to form branched allylic substitution products. The bulk of the recent literature on iridium-catalyzed allylic substitution has focused on catalysts derived from  $[\text{Ir}(\text{COD})\text{Cl}]_2$  and phosphoramidite ligands. These complexes catalyze the formation of enantiomerically enriched allylic amines, allylic ethers, and  $\beta$ -branched  $\gamma$ - $\delta$  unsaturated carbonyl compounds. The latest generation and most commonly used of these catalysts (Scheme 1) consists of a cyclometalated iridium-phosphoramidite core chelated by 1,5-cyclooctadiene. A fifth coordination site is occupied in catalyst precursors by an additional  $\kappa^1$ -phosphoramidite or ethylene. The phosphoramidite that is used to generate the metalacyclic core typically contains one BINOLate and one bis-arylethylamino group on phosphorus.

Iridium-catalyzed allylic substitution has undergone rapid development since it was first reported [1]. What began as a reaction of malonate nucleophiles catalyzed by complexes containing achiral ligands to form racemic product has become a valuable method for the enantioselective formation of carbon–carbon, –nitrogen, and –oxygen bonds. Mechanistic investigations have slowly increased our understanding of the inner workings of the catalytic cycle, and experimentally grounded proposals for the origins of enantioselectivity can now be made.

This chapter describes the development of iridium-catalyzed, enantioselective allylic substitution. It is organized to focus on how modifications to the catalyst, combined with mechanistic insights, have provided the foundation for a steady



**Scheme 1** Asymmetric allylic substitution catalyzed by metalacyclic iridium-phosphoramidite complexes

increase in substrate scope, yield, and selectivity. Ultimately, this review describes how these advances have led to the most recently reported generation of metalacyclic iridium-phosphoramidite catalysts.

## 2 Background


### 2.1 An Overview of Allylic Substitution

Metal-catalyzed allylic substitution (Fig. 1) leads to the displacement of an ester, halide, or other leaving group by a nucleophile in an allylic position. Substitution occurs in a formal  $\text{S}_{\text{N}}2$  or  $\text{S}_{\text{N}}2'$  fashion, depending on the identity of the two reactants and the catalyst. However, the mechanism is not similar to uncatalyzed  $\text{S}_{\text{N}}2$  or  $\text{S}_{\text{N}}2'$  reactions of purely organic systems. Instead, the metal catalyst initially displaces the leaving group to form an  $\eta^3$ -allyl complex, which is subsequently attacked by the nucleophile at one terminus of the allyl ligand over the other to form the product. Nucleophilic attack on the substituted terminus of an allyl ligand can produce a chiral product. Thus, the use of an appropriate chiral, non-racemic ligand can result in the formation of this product in enantiomerically enriched form.


### 2.2 Nucleophilic Attack on $\eta^3$ -Allyl Metal Complexes

Iridium-catalyzed allylic substitution was first investigated after many years of development of allylic substitution reactions catalyzed by a variety of complexes of other metals, particularly those containing palladium. While iridium-catalyzed





**Fig. 1** Metal-catalyzed allylic substitution



**Fig. 2** Stoichiometric nucleophilic attack on the terminus of an allyl ligand

allylic substitution dates back to 1997, the addition of nucleophiles to  $\pi$ -allyl metal complexes has been known since 1965, when Tsuji demonstrated that the carbon nucleophiles ethyl malonate, ethyl acetoacetate, and 1-morpholino-1-cyclohexene react with the terminus of the  $\pi$ -allyl ligand of allylpalladium chloride to form mixtures of mono- and diallylation products (Fig. 2) [2].

## 2.3 Early Metal-Catalyzed Allylic Substitution

### 2.3.1 First Metal-Catalyzed Allylic Substitution

After the initial demonstration of stoichiometric nucleophilic attack on  $\pi$ -allyl ligands, catalytic allylic substitution reactions were pursued. In 1970, groups from Union Carbide [3, 4], Shell Oil [5], and Toray Industries [6] published or patented examples of catalytic allylic substitution. All three groups reported allylic amination with palladium catalysts. The Toray Industries report also demonstrated the exchange of aryl ether and ester leaving groups, and the patent from Shell Oil includes catalysts based on rhodium and platinum.




Fig. 3 Enantiotopic termini or faces of  $\eta^3$ -allyl ligands


### 2.3.2 The Selectivity of Allylic Substitution

Many of the enantioselective allylic substitution reactions are based on the two reaction manifolds in Fig. 3. In the first manifold, enantioselection can result from nucleophilic attack at one of two enantiotopic termini of a symmetrical  $\pi$ -allyl ligand. In the second manifold, enantioselection can result from attack at one of the two enantiotopic faces of an allyl ligand that is substituted at only one of the two termini. Thus, initial selective addition of the metal fragment to one face of the allyl or unselective addition to both faces followed by isomerization to form one stereoisomer selectively will lead to the formation of one enantiomer of product over the other when nucleophilic attack occurs at the more substituted terminus of the allyl ligand. Palladium-catalyzed reactions that generate  $\pi$ -allyl intermediates containing one unsubstituted terminus typically lead to reaction at the unsubstituted terminus, forming to products lacking a stereocenter at the electrophilic carbon. Thus, the formation of enantioenriched branched products from  $\pi$ -allyl intermediates containing one unsubstituted terminus required the development of a new type of catalyst.

## 2.4 The State of Allylic Substitution in 1997

The first iridium catalysts for allylic substitution were published in 1997. Takeuchi showed that the combination of  $[\text{Ir}(\text{COD})\text{Cl}]_2$  and triphenylphosphite catalyzes the addition of malonate nucleophiles to the substituted terminus of  $\eta^3$ -allyliridium intermediates that are generated from allylic acetates. This selectivity for attack at the more substituted terminus gives rise to the branched allylic alkylation products (Fig. 4), rather than the linear products that had been formed by palladium-catalyzed allylic substitution reactions at that time [7]. The initial scope of iridium-catalyzed allylic substitution was also restricted to stabilized enolate nucleophiles, but it was quickly expanded to a wide range of other nucleophiles.

**Fig. 4** Branched and linear products formed from  $\eta^3$ -allyl intermediates that have one unsubstituted terminus



### 2.4.1 Contemporaneous Metal-Catalyzed Allylic Substitution

Most allylic substitution reactions catalyzed by other metals are selective for the formation of branched products. Although this had been demonstrated for a large portion of the d-block before Takeuchi's work with iridium, most of the progress in this area was restricted to stabilized enolate nucleophiles.

Prior to the initial disclosure of iridium-catalyzed allylic substitution, catalysts based on complexes of molybdenum [8–18], tungsten [19–24], iron [25–28], ruthenium [11, 29, 30], cobalt [27], rhodium [11, 31], nickel [11, 32–37], and platinum [38] had all been shown to favor the formation of branched products from substrates that generate  $\pi$ -allyl intermediates that are unsubstituted at one terminus. In fact the widely recognized selectivity of palladium catalysts for the formation of linear products is actually an anomaly among transition metals. Some reactions catalyzed by rhodium and iron catalysts exhibit regiochemical “memory effects,” meaning that the nucleophile adds to the same terminus of the allyl group originally occupied by the leaving group. Branched starting materials led to branched products, and linear starting materials led to linear products [39, 40]. Catalysts based on ruthenium, cobalt, and nickel complexes were reported to form branched products selectively, but this regioselectivity was not typically high. Only catalysts based on molybdenum and tungsten formed branched products from allylic substitution with high selectivity. However, these early reactions typically required catalyst loadings of 10 mol% or more, and their use with nitrogen or oxygen nucleophiles had not yet been reported.

Thus, a catalyst for allylic substitution that is highly active and selective for the formation of products from addition to the more substituted terminus from a variety of nucleophiles and allylic esters was unknown. Iridium catalysts have now been developed that begin to fill this void.

## 3 Early Iridium-Catalyzed Allylic Substitutions

### 3.1 *Allylic Substitution Catalyzed by [Ir(COD)Cl]<sub>2</sub> and Triphenylphosphite*

The first examples of iridium-catalyzed allylic substitution [1] occurred between stabilized carbon nucleophiles and both alkyl- and aryl-substituted allylic alcohol derivatives with exceptional selectivity for the branched substitution product.

**Table 1** Allylic substitution catalyzed by  $[\text{Ir}(\text{COD})\text{Cl}]_2$  and  $\text{P}(\text{O}Ph)_3$ 

Entry	b or I/R <sup>1</sup> /R <sup>2</sup>	LG	T (°C)	Time (h)	Yield (%)	b:I or b:(E)-I:(Z)-I
1	I/Ph/H	OAc	rt	3	98	99:1
2	b/Ph/H	OAc	rt	3	91	99:1
3	I/H/C <sub>3</sub> H <sub>7</sub>	OAc	65	5	85	7:3:90
4	I/Ph/H	OH	65	1	98	99:1
5	I/Me/Me	OAc	65	24	n.r. <sup>a</sup>	
6	I/Me/Me	OPO(OEt) <sub>2</sub>	rt	4	81	96:4
7	b/Me/Me	OAc	rt	2	80	100:0
8	I/C <sub>3</sub> H <sub>7</sub> /H	OAc	rt	3	89	96:4
9 <sup>b</sup>	I/C <sub>3</sub> H <sub>7</sub> /H	OAc	65	9	90	95:5
10 <sup>c</sup>	I/C <sub>3</sub> H <sub>7</sub> /H	OAc	65	16	6	24:48:28
11	I/Ph/H	OCO <sub>2</sub> Me	50	3	95	95:5

<sup>a</sup>No reaction<sup>b</sup>Conducted with  $\text{Ir}(\text{COD})_2\text{BF}_4$ <sup>c</sup>Conducted with  $\text{PPh}_3$  (4 mol%)

In this and subsequent studies [41–44], Takeuchi and coworkers described the scope and selectivity of catalysts derived from  $[\text{Ir}(\text{COD})\text{Cl}]_2$  and achiral phosphorus ligands. Many of the trends that they uncovered have also been observed with more recently developed catalyst systems derived from chiral phosphoramidite ligands (Table 1).

Stabilized enolate and amine nucleophiles add to both branched and (*E*)-linear allylic alcohol derivatives to form products in high yield and over 95% selectivity for the branched product when catalyzed by a combination of  $[\text{Ir}(\text{COD})\text{Cl}]_2$  and  $\text{P}(\text{O}Ph)_3$ . In contrast, substitution of (*Z*)-allylic alcohol derivatives results in formation of (*Z*)-linear products, especially for alkyl-substituted allylic esters. Based on the net retention of configuration observed for allylic substitution of a cyclic allylic carbonate, such substitutions were proposed to occur through a “double-inversion mechanism” in which both oxidative addition of the allylic ester and reaction of the allyl intermediate with the nucleophilic reagent occur with inversion of configuration. Later work by Helmchen and coworkers supported a double inversion mechanism by showing that hindered allylic acetates that are unable to undergo an inversion of configuration were unreactive toward allylic substitution in the presence of iridium catalysts, whereas

analogous reactions of unhindered cyclopentenyl acetates proceeded to full conversion [45]. Finally, as discussed in Sect. 5, recent studies on discrete, isolated allyliridium intermediates by Hartwig and coworkers support a mechanism for nucleophilic attack on allyliridium intermediates that proceeds by inversion of configuration [46].

Although most of these early reactions were conducted with allylic acetates, reactions of allylic carbonates, trifluoroacetates, and phosphates also occur. Sodium diethylmalonate also reacts with allylic alcohols in the presence of the iridium-triphenylphosphite catalyst. However, the alcohol itself does not act as a leaving group. Instead, transesterification occurs with one equivalent of malonate nucleophile to form a more labile ester leaving group.

Addition to linear 1,1-disubstituted allylic acetates is slower than addition to monosubstituted allylic esters. Additions to allylic trifluoroacetates or phosphates are faster than additions to allylic carbonates or acetates, and reactions of branched allylic esters are faster than additions to linear allylic esters. Aryl-, vinyl, alkynyl, and alkyl-substituted allylic esters readily undergo allylic substitution. Amines and stabilized enolates both react with these electrophiles in the presence of the catalyst generated from an iridium precursor and triphenylphosphite.

### 3.1.1 Iridium Precursor and Ligands

Branched allylic substitution products were obtained with the highest yields and selectivities when the reactions were conducted with the combination of  $[\text{Ir}(\text{COD})\text{Cl}]_2$  and  $\text{P}(\text{OPh})_3$  as catalyst. Other iridium precursors were tested, but they all required higher reaction temperatures, and most of these catalyst systems produced allylic substitution products in lower yield and branched-to-linear selectivity. The use of a  $\pi$ -accepting ligand such as  $\text{P}(\text{OPh})_3$  proved to be necessary in most cases to obtain high branched selectivity. Reactions conducted with catalysts containing less  $\pi$ -accepting ligands occurred with lower selectivity for the branched isomer.

Catalysts lacking phosphorus ligands have also been used as catalysts for allylic substitutions.  $[\text{Ir}(\text{COD})\text{Cl}]_2$  itself, which contains a  $\pi$ -accepting diolefin ligand, catalyzes the alkylation of allylic acetates, but the formation of branched products was only favored when the substitution reaction was performed with branched allylic esters. Takemoto and coworkers later reported the etherification of branched allylic acetates and carbonates with oximes catalyzed by  $[\text{Ir}(\text{COD})\text{Cl}]_2$  without added ligand [47]. Finally, as discussed in Sect. 6, Carreira reported kinetic resolutions of branched allylic carbonates from reactions of phenol catalyzed by the combination of  $[\text{Ir}(\text{COE})_2\text{Cl}]_2$  and a chiral diene ligand [48].

One notable deviation from this trend of increased branched selectivity with increased  $\pi$ -accepting character has been reported. Nomura and coworkers reported allylic substitution reactions catalyzed by  $[\text{Ir}(\text{COD})\text{Cl}]_2$  and triphenylphosphine to form polymers linked by branched 1,1'-(1,4-phenylene)diprop-2-enyl units [49]. Despite this exception, most iridium-catalyzed allylic substitution


reactions have been conducted with complexes derived from  $[\text{Ir}(\text{COD})\text{Cl}]_2$  and an electron-accepting ligand.

### 3.2 Asymmetric Iridium-Catalyzed Allylic Substitution: A Survey of Ligands

The formation of chiral products from nucleophilic addition to the more substituted terminus of monosubstituted allyl groups in the presence of iridium catalysts containing achiral  $\pi$ -accepting phosphorus ligands led to efforts to conduct these reactions with related enantioenriched ligands that control the absolute stereochemistry. Catalysts generated from phosphoramidite ligands have been used most commonly for this process, and complexes of these ligands catalyze allylic substitution reactions with broad substrate scope, high activity, and high regio- and enantioselectivity. However, catalysts containing several other families of ligands have also been studied. Reactions catalyzed by complexes of the other families of ligands will be reviewed first.

#### 3.2.1 Reactions Catalyzed by Iridium Complexes of PHOX Ligands

The first enantioselective, iridium-catalyzed allylic substitution was reported by Helmchen and coworkers soon after the initial report by Takeuchi. Helmchen studied catalysts generated from phosphinooxazoline (PHOX) ligands and  $[\text{Ir}(\text{COD})\text{Cl}]_2$  for the reactions of sodium dimethylmalonate with cinnamyl acetates (Scheme 2) [50]. The alkylation products were isolated in nearly quantitative yield and were formed with ratios of branched-to-linear products up to 99:1 and with enantioselectivities up to 95% ee. In this and subsequent studies with PHOX ligands [51, 52], Helmchen et al. demonstrated that the highest yields and selectivities were obtained with a PHOX ligand containing electron-withdrawing substituents and



**Scheme 2** Allylic alkylation catalyzed by  $[\text{Ir}(\text{COD})\text{Cl}]_2$  and a PHOX ligand


with additives such as LiF. Mark, This is not shown in Scheme 2. Although substitution of only two allylic carbonates was reported, this work showed that iridium complexes could catalyze allylic substitution reactions with high enantioselectivity. However, intramolecular allylic amination catalyzed by complexes of the PHOX ligand [53] occurred with low enantioselectivity, and the regioselectivity was lower for the reactions of alkyl-substituted allylic esters than for the reactions of aromatic allylic esters. Thus, further studies on iridium-catalyzed allylic substitution with complexes of PHOX ligands have not been reported.

### 3.2.2 Reactions Catalyzed by Iridium Complexes of Chiral Phosphite Ligands


Iridium complexes containing chiral phosphite ligands also catalyze allylic alkylation. Fuji and coworkers reported reactions of dimethylmalonate with cinnamyl methyl carbonate catalyzed by complexes of a BINOL-derived phosphite (Scheme 3) [54, 55]. With *n*BuLi as base and ZnCl<sub>2</sub> as additive, the alkylation product was isolated in 99% yield with 93% branched selectivity and 96% ee. Iridium complexes of another BINOL-derived phosphite containing a thioether substituent were shown to catalyze the alkylation of cinnamyl phosphates with a protected glycine nucleophile (Scheme 4) [56, 57]. This reaction is particularly noteworthy because it occurs with high diastereoselectivity. Such high diastereoselectivity has not been achieved when using other iridium catalysts. Furthermore, varying the solvent and base led to the formation of one diastereomer over the other, with the major diastereomer formed in excellent enantiomeric excess in both cases. Less than 1% of linear products were detected in these reactions.

### 3.2.3 Reactions Catalyzed by Iridium Complexes of Pybox Ligands


Complexes derived from the oxazolinyipyridine ligand Ph-Pybox and [Ir(COD)Cl]<sub>2</sub> catalyze allylic substitution reactions of several oxygen and nitrogen nucleophiles



**Scheme 3** Allylic alkylation catalyzed by [Ir(COD)Cl]<sub>2</sub> and a chiral phosphite ligand



**Scheme 4** Diastereoselective allylic alkylation with protected glycine



**Scheme 5** Allylic substitution catalyzed by [Ir(COD)Cl<sub>2</sub>] and Ph-Pybox

that have not been used in systems catalyzed by metalacyclic iridium phosphoramidite complexes (Scheme 5). Reactions of amines [58], hydroxylamines [58, 59], oximes [58, 60], and guanidines [61] with cinnamyl phosphates occur in good to excellent yield, branched selectivity, and enantiomeric excess. *O*-substituted hydroxylamines react at nitrogen. However, hydroxylamines react preferentially




at oxygen when the hydroxylamine possesses an electron-withdrawing group on nitrogen.


### 3.2.4 Reactions Catalyzed by Iridium Complexes of DIAPHOX and Phosphorodiamidite Ligands

The phosphoramidite ligands that are the focus of the remainder of this chapter have prompted the investigation of ligands containing related structures. Iridium complexes of aspartic acid-derived P-chirogenic diaminophosphine oxides (DIAPHOXs) catalyze the amination [62] and alkylation [63] of allylic carbonates (Scheme 6). With BSA as base and catalytic amounts of  $\text{NaPF}_6$  as additive, branched amination and alkylation products were obtained from cinnamyl carbonates in excellent yields and enantioselectivities. However, the yields and enantioselectivities were lower for the reactions of alkyl-substituted allylic carbonates. Added  $\text{LiOAc}$  increased the enantioselectivities of allylic alkylation products.


Catalysts generated from  $[\text{Ir}(\text{COD})\text{Cl}]_2$  and phosphorodiamidite ligands containing a pyrrolo[1,2-*c*] diazaphosphol-1-one backbone catalyze the addition of phenol to methyl cinnamyl carbonate (Scheme 7) [64]. The arrangement of the steric bulk around the ligand determines the configuration of the product; catalysts containing the two ligands in which the *N*-aryl and *O*-aryl groups are reversed form the opposite enantiomers. DFT calculations were performed to determine the origin of this reversal in selectivity, and these studies suggested that reversal of the aryl groups changes the face of the allyl ligand that is preferentially coordinated to the metal center.



**Scheme 6** Allylic substitution catalyzed by  $[\text{Ir}(\text{COD})\text{Cl}]_2$  and DIAPHOX



**Scheme 7** Allylic etherification catalyzed by [Ir(COD)Cl]<sub>2</sub> and a phosphorodiamidite ligand



**Scheme 8** Allylic substitution catalyzed by [Ir(COD)Cl]<sub>2</sub> and monophos

### 3.3 Iridium Catalysts Containing Phosphoramidite Ligands

#### 3.3.1 Reactions Catalyzed by Complexes of Monophos

Although Helmchen et al. showed that asymmetric iridium-catalyzed allylic substitution could be achieved, the scope of the reactions catalyzed by iridium complexes of the PHOX ligands was limited. Thus, they evaluated reactions catalyzed by complexes generated from [Ir(COD)Cl]<sub>2</sub> and the dimethylamine-derived phosphoramidite monophos (Scheme 8) [45, 51]. Although selectivity for the branched isomer from addition of malonate nucleophiles to allylic acetates was excellent, the highest enantiomeric excess obtained was 86%. This enantiomeric excess was obtained from a reaction of racemic branched allylic acetate. The enantiomeric excess was lower when linear allylic acetates were used. This system catalyzed addition of the lithium salts of *N*-benzyl sulfonamides to allylic acetates, but the product of the reaction between this reagent and an alkyl-substituted linear allylic acetate was formed with an enantiomeric excess of 13%.

#### 3.3.2 Reactions Catalyzed by Complexes of Bis-Phenethylamine-Derived Phosphoramidite, L1

##### Amination with L1

Although reactions catalyzed by [Ir(COD)Cl]<sub>2</sub> and the dimethylamine-based monophos occurred with only modest enantioselectivity, a wide variety of reactions

catalyzed by iridium complexes of other phosphoramidite ligands have occurred with excellent enantioselectivity. Ohmura and Hartwig reported that the combination of bisphenethylamine-derived phosphoramidite **L1** (Scheme 1) and [Ir(COD)Cl]<sub>2</sub> catalyze the amination of a series of allylic carbonates with primary and secondary amines in excellent yield, branched selectivity, and enantioselectivity (Table 2) [65]. **L1** was originally developed by Feringa and coworkers for use in copper-catalyzed conjugate additions of alkylzinc reagents to  $\alpha$ - $\beta$ -unsaturated ketones [67].


The scope of allylic electrophiles that react with amines was shown to encompass electron-neutral and electron-rich cinnamyl methyl carbonates, as well as furan-2-yl and alkyl-substituted allylic methyl carbonates. An *ortho*-substituted cinnamyl carbonate was found to react with lower enantioselectivity, a trend that has been observed in later studies of reactions with other nucleophiles. The electron-poor *p*-nitrocinnamyl carbonate also reacted, but with reduced enantioselectivity. Allylic amination of dienyl carbonates also occur in some cases with high selectivity for formation of the product with the amino group at the  $\gamma$ -position over the  $\varepsilon$ -position of the pentadienyl unit [66]. Arylamines did not react with allylic carbonates under these conditions. However, they have been shown to react in the presence of the metalacyclic iridium-phosphoramidite catalysts that are discussed in Sect. 4.

### Iridium-Catalyzed Allylic Etherification with Catalysts Derived from **L1**

Hartwig et al. demonstrated that the same combination of iridium precursor and phosphoramidite **L1** also catalyzes allylic etherifications (Scheme 9) [68]. Lithium and sodium aryloxides were shown to react with cinnamyl and hex-2-enyl carbonates to form the branched allylic ethers in high yield, with high branched-to-linear

**Table 2** Allylic amination catalyzed by [Ir(COD)Cl]<sub>2</sub> and **L1**

Entry	R	Nu	Yield (%)	b:l	ee (%)	t(h)	Reference
1	Ph	BnNH <sub>2</sub>	84	98:1	95	10	[65]
2	Ph	n-hexylNH <sub>2</sub>	88	98:2	96	9	[65]
3	Ph	pyrrolidine	75	98:2	97	2	[65]
4	4-(MeO)C <sub>6</sub> H <sub>4</sub>	BnNH <sub>2</sub>	88	99:1	96	9	[65]
5	4-(NO <sub>2</sub> )C <sub>6</sub> H <sub>4</sub>	BnNH <sub>2</sub>	67	83:13	86	12	[65]
6	2-(MeO)C <sub>6</sub> H <sub>4</sub>	BnNH <sub>2</sub>	77	95:4	76	16	[65]
7	C <sub>3</sub> H <sub>7</sub>	BnNH <sub>2</sub>	66	88:8	95	10	[65]
8	PMBO	BnNH <sub>2</sub>	71	96:4	96	14	[66]



R <sup>1</sup>	R <sup>2</sup>	ArOM	Yield (%)	b:l	ee (%)	t (h)
Ph	Me	PhOLi	86	96/4	96	20
Ph	Me	4-(MeO)C <sub>6</sub> H <sub>4</sub> OLi	88	98:2	97	8
Ph	Et	4-BrC <sub>6</sub> H <sub>4</sub> ONa	91	96/4	90	8
Ph	Et	4-(NO <sub>2</sub> )C <sub>6</sub> H <sub>4</sub> ONa	n.r. <sup>a</sup>			
2-(MeO)C <sub>6</sub> H <sub>4</sub>	Me	PhOLi	79	98/2	75	41
nPr	Me	PhOLi	93	92/8	92	14


<sup>a</sup>No reaction

### Scheme 9 Allylic etherification catalyzed by [Ir(COD)Cl]<sub>2</sub> and L1


ratios, and enantiomeric excesses of up to 97%. The use of alkali metal aryloxides was necessary to obtain useful reaction rates; reactions of phenols in combination with amine bases occurred in lower yield and enantioselectivity. In some cases, transesterification of the carbonate with the phenoxide competed with the allylic substitution process. In these cases, reactions of ethyl allylic carbonates, rather than methyl allylic carbonates, occurred in high yield. As was observed for iridium-catalyzed allylic amination, the reaction of an *ortho*-substituted cinnamyl carbonate occurred with lower enantioselectivity than reactions of *meta*- and *para*-substituted cinnamyl carbonates. In contrast, the reactivity of *ortho*-substituted aryloxides was comparable to that of less hindered aryloxides. However, reactions of electron-deficient aryloxides occurred with lower branched-to-linear ratios and lower enantioselectivity. The most electron-poor aryloxides derived from 4-(NO<sub>2</sub>)C<sub>6</sub>H<sub>4</sub>OH did not react. The lithium phenoxides were added directly to reaction mixtures, or they were generated in situ from the corresponding phenol and *n*BuLi or Cy<sub>2</sub>NLi.

### Iridium-Catalyzed Asymmetric Allylic Alkylation with Catalysts Derived from L1

Reactions of allylic electrophiles with stabilized carbon nucleophiles were shown by Helmchen and coworkers to occur in the presence of iridium-phosphoramidite catalysts containing L1 (Scheme 10) [66, 69], but alkylations of linear allylic acetates with salts of dimethylmalonate occurred with variable yield, branched-to-linear selectivity, and enantioselectivity. Although selectivities were improved by the addition of lithium chloride, enantioselectivities still ranged from 82–94%, and branched selectivities from 55–91%. Reactions catalyzed by complexes of phosphoramidite ligands derived from primary amines resulted in the formation of alkylation products with higher branched-to-linear ratios but lower enantioselectivities. These selectivities were improved by the development of metalacyclic iridium catalysts discussed in the next section and “salt-free” reaction conditions described later in this chapter.



**Scheme 10** Iridium-catalyzed allylic alkylation catalyzed by  $[\text{Ir}(\text{COD})\text{Cl}]_2$  and **L1**



**Scheme 11** Formation of the active metalacyclic iridium phosphoramidite catalyst

## 4 Advances in Mechanistic Understanding and Catalyst Design: Cyclometalation and Modification of the Phosphoramidite Ligand

### 4.1 Discovery of the Active Metalacyclic Catalyst

Although the combination of  $[\text{Ir}(\text{COD})\text{Cl}]_2$  and **L1** was shown to catalyze the alkylation, amination, and etherification of allylic esters to form the branched substitution product in high yield and enantioselectivity, the identity of the active catalyst in these reactions had not been identified. The combination of  $[\text{Ir}(\text{COD})\text{Cl}]_2$  and **L1** forms the square-planar  $[\text{Ir}(\text{COD})(\text{Cl})\text{L1}]$  (**4**) (Scheme 11) [45]. However, this complex does not react with allylic carbonates to form an appreciable amount of an allyl complex, and the absence of this reactivity suggested that the mechanism or identity of the active catalyst was more complex than that from simple addition of the allylic ester to the square-planar complex containing a  $\kappa^1$ -phosphoramidite ligand.

Instead of reacting with the allylic ester, the square-planar complex **4** reacts with pyrrolidine to form a new species, **1a**, which contains a cyclometalated

*P,C*-bound  $\kappa^2$ -phosphoramidite in addition to a  $\kappa^1$ -phosphoramidite (Scheme 11). This cyclometalated complex was identified by NMR spectroscopy and X-ray crystallography [70]. A cyclometalated species had been proposed to form in reactions catalyzed by  $[\text{Ir}(\text{COD})\text{Cl}]_2$  and  $\text{P}(\text{OPh})_3$ . However, no complexes were isolated from this mixture or shown to be competent to catalyze allylic substitution [45].

Mechanistic studies showed that metalacycle **1a** is competent to be a catalyst in asymmetric allylic substitution reactions. The reaction of benzylamine with methyl cinnamyl carbonate catalyzed by a mixture of **L1** and  $[\text{Ir}(\text{COD})\text{Cl}]_2$  occurs with an induction period and forms product in 84% yield and 95% ee, whereas the same reaction catalyzed by a mixture of metalacycle **1a** and  $[\text{Ir}(\text{COD})\text{Cl}]_2$  occurs without an induction period in just 2 hours to form the substitution product in 81% yield and 97% ee. The latter reaction was conducted with added  $[\text{Ir}(\text{COD})\text{Cl}]_2$  to trap the  $\kappa^1$ -bound **L1** after dissociation. This ligand must dissociate to provide a site for oxidative addition of the allylic carbonate.

The synthesis of pure metalacycle **1a** from  $[\text{Ir}(\text{COD})\text{Cl}]_2$  and **L1** requires only amine base and heat, followed by precipitation and removal of amine hydrochloride. However, this complex was typically generated in situ during early studies by the treatment of a combination of  $[\text{Ir}(\text{COD})\text{Cl}]_2$  and **L1** with an amine base, such as *n*PrNH<sub>2</sub>, 1,5,7-triazabicyclo[4.4.0]dec-5-ene (TBD), or 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU). If a 1:1 ratio of iridium to ligand is used, a mixture of **1a** and  $[\text{Ir}(\text{COD})\text{Cl}]_2$  is produced. Helmchen et al. have reported that catalyst activation in the presence of tetrahydrothiophene (THT) prevents coordination of the  $\kappa^1$ -phosphoramidite [71].

## 4.2 Effect of the Phosphoramidite Ligand

Concurrent with studies on cyclometalation, studies on the effects of the structure of phosphoramidite ligand had been conducted. Several groups studied the effect of the structure of ligand on the rate and selectivity of these iridium-catalyzed allylic substitutions. **L1** contains three separate chiral components – the two phenethyl moieties on the amine as well as the axially chiral BINOL backbone. These portions of the catalyst structure can control reaction rates by affecting the rate of cyclometalation, by inhibiting catalyst decomposition, or by forming a complex that reacts faster in the turnover-limiting step(s) of the catalytic cycle.

### 4.2.1 Modification of the Amino Groups on the Phosphoramidite

Catalysts containing **L1** were much more reactive and selective than those containing monophos. The conclusion that the cyclometalated core of complex **1a** is present in the active catalyst explains why the complexes of **L1** are more reactive

than those of monophos. Complexes containing monophos cannot form a species containing an analogous stable, five-membered metalacycle.

Modifications were made to the aryylethyl substituent of the phosphoramidite to identify catalysts that would exhibit even higher reactivity and selectivity than **L1**. Some studies of reactions catalyzed by complexes of different phosphoramidites were performed with preactivated catalyst, but others were performed without activation of the catalyst. Thus, it is not straightforward to determine which ligands form the most active catalysts. Nonetheless, certain modifications to the bisphenethylamine portion of phosphoramidite **L1** improve the rates, yields, and selectivities of allylic substitution reactions catalyzed by the modified ligands and [Ir(COD)Cl]<sub>2</sub>. Phosphoramidite ligands **L2** or **L3** derived from bisnaphthethylamine or 1-(2-methoxyphenyl)ethylamine, the latter of which is not yet commercially available from typical suppliers (Table 3), generate catalysts that are more reactive or more selective for reactions of certain substrate combinations. Complexes derived from **L2** and **L3** cyclometalate faster than complexes derived from **L1**, leading to a shorter induction period when the catalyst is not formed prior to addition of the reagents [75]. Likewise, the metalacyclic complex **2a** derived from **L2** is more resilient than **1a** to deactivation by acidic pronucleophiles [76].

Additional studies were conducted to determine how further modifications to the amine portion of the phosphoramidite ligand affect iridium-catalyzed allylic substitution. One aryylethyl moiety is necessary for the formation of metalacyclic active catalyst, but it was unclear how changes to the structure of the second substituent on nitrogen would affect reactivity and selectivity. A stereocenter on this second

**Table 3** Comparison of phosphoramidite ligands

R' = cyclododecyl

Entry	R	L	Time (h)	Yield (%)	ee (%)	b:l	Activated prior to adding reagents?	Reference
1	C <sub>3</sub> H <sub>7</sub>	<b>L1</b>	3	58	95	94:6	Yes	[72]
2	C <sub>3</sub> H <sub>7</sub>	<b>L3</b>	0.5	67	98	94:6	Yes	[72]
3	Ph	<b>L1</b>	2	81	97	98:2	Yes	[70]
4	Ph	<b>L1</b>	12	84	95	98:2	No	[70]
5	Ph	<b>L2</b>	16	88	97	98:2	No	[73]
6	Ph	<b>L4</b>	3	93	95	99:1	Yes	[74]
7	Ph	<b>L5</b>	3	86	94	98:2	Yes	[74]

substituent was shown to be unnecessary to form metalacyclic iridium catalyst that reacted with high enantioselectivities. Phosphoramidites synthesized from  $C_1$ -symmetric amines in which one phenethyl group has been replaced by a bulky cycloalkyl group generate catalysts that display reactivity similar to that of the catalysts containing related phosphoramidites synthesized from  $C_2$ -symmetric amines (Table 3, entry 6) [74, 77, 78].

#### 4.2.2 Modification of the Diol Backbone of the Phosphoramidite

The configuration of the chiral BINOLate backbone of the phosphoramidite ligand affects the rates and enantioselectivities of allylic substitution reactions. Hartwig and coworkers found that allylic substitution conducted with a catalyst derived from the simplified ligand ( $S_a,R$ )-**L4** occurred more slowly than that conducted with a catalyst derived from ( $R_a,R$ )-**L4** [74]. Complexes of the “mismatched” ( $S_a,R$ )-**L4** undergo cyclometalation slowly. The products formed from reactions catalyzed by complexes of ( $S_a,R$ )-**L4** and ( $R_a,R$ )-**L4** have the opposite absolute configuration.

The resolved BINOLate backbone of the phosphoramidite ligand is not required for high catalyst activity and selectivity. Catalysts generated from ligands in which the BINOLate backbone has been replaced with an achiral biphenolate catalyze allylic substitution reactions in high yield and selectivity despite rapid room temperature atropisomerism of the biphenolate [74, 77]. Iridium-catalyzed allylic substitution occurs with a catalyst generated from the simplified phosphoramidite containing just one chiral center (one phenethyl group on nitrogen) and no resolved, axially chiral components (Table 3, entry 7).

### 4.3 Scope of Allylic Substitution Catalyzed by Metalacyclic Iridium-Phosphoramidite Complexes

Metalacyclic iridium-phosphoramidite complexes generated prior to addition of the reagents react with a substantially broader scope than those generated in situ. Some nucleophiles, such as arylamines, are not sufficiently basic to induce cyclometalation and do not react with allylic carbonates in the presence of  $[\text{Ir}(\text{COD})\text{Cl}]_2$  and **L1**. In reactions of some other nucleophiles, catalyst activation by cyclometalation is slow, and a long induction period results. Thus, it is typically advantageous to conduct reactions with the “pre-activated” metalacyclic species generated prior to addition of the reagents. Coupled with a set of modified phosphoramidite ligands that lead to improved reactivity and selectivity, these new catalyst systems have been applied quickly to reactions of a variety of nucleophiles, some of which have been commonly used for allylic substitution with catalysts containing other metals and others that have rarely been used with catalysts of any metal.




### 4.3.1 Iridium-Catalyzed Allylic Substitution with Carbon Nucleophiles

#### Iridium-Catalyzed Allylic Substitution with Stabilized Enolates


Malonate nucleophiles react with allylic carbonates in the presence of **L1** and  $[\text{Ir}(\text{COD})\text{Cl}]_2$ . However, the yields and selectivities of these reactions were lower than those obtained from analogous reactions with nitrogen and oxygen nucleophiles. The use of a preactivated metalacycle increased the selectivities of products formed from iridium-catalyzed allylic alkylation reactions. By forming the active metalacycle from a mixture of **L1** and  $[\text{Ir}(\text{COD})\text{Cl}]_2$  and base, Helmchen et al. demonstrated that products formed from the reaction of sodium dimethylmalonate with linear allylic carbonates were formed with enantioselectivities above 96% (Scheme 12) [71]. Additional improvements were realized by applying “salt-free” conditions [79]. Rather than deprotonating the malonate nucleophile prior to addition of the other reaction components, the neutral malonate was added; the alkoxide generated from decarboxylation of the carbonate leaving group deprotonates the malonate. By combining “salt-free” conditions with catalysts derived from the more active ligand **L2**, high yields of the branched alkylation product were obtained in excellent enantioselectivity (Scheme 12).

#### Allylic Substitution with Unstabilized Enolates

Unstabilized enolates react with allylic carbonates in the presence of metalacyclic iridium-phosphoramidite catalysts. Although ketones and aldehydes have not yet been used directly as pronucleophiles with this catalyst system, silyl enol ethers [80] and enamines [81] react with linear allylic carbonates to form, after workup,  $\beta$ -branched,  $\gamma$ - $\delta$  unsaturated ketones (Scheme 13). Both methods form products in high yield, branched selectivity, and enantioselectivity for a range of cinnamyl and alkyl-substituted allylic carbonates. However, the silyl enol ethers derived from aliphatic ketones reacted in lower yields than enamines derived from the same ketones.



**Scheme 12** Allylic alkylation with activated catalyst before and after applying “salt-free” conditions

**Scheme 13** Allylic substitution with unstabilized enolate nucleophiles

In addition, unstabilized enolate nucleophiles have been generated by decarboxylation of  $\beta$ -keto-carboxylates. In this case, no additives are required to activate the nucleophile, but the highest yields and selectivities were obtained in the presence of two equivalents of DBU [82]. Although reactions of allylic carbonates containing aromatic, heteroaromatic, and aliphatic substituents occurred, only reactions to form aryl ketone products were published.

### Allylic Substitution with Nitroalkanes


Nitroalkanes react with allylic carbonates in the presence of iridium catalysts [83]. However, nitroalkanes are prochiral, and products are formed with poor diastereoselectivity. Nitroalkanes and other prochiral nucleophiles are discussed further in Sect. 6.

### Allylic Substitution with “Hard” Carbon Nucleophiles

Hard carbon nucleophiles that do not contain stabilizing electron-withdrawing substituents have been shown to react with allylic carbonates only recently in the

presence of iridium catalysts. Alexakis et al. reported conditions for the reactions of arylzinc reagents with allylic carbonates in the presence of iridium phosphoramidite catalysts [84, 85]. The arylzinc reagents were generated from the corresponding aryl Grignard reagent and  $\text{ZnBr}_2$ . Although regioselectivities were poor, typically around 1:1, the branched-to-linear ratio of this reaction was higher than that of other allylic arylation reactions, which exclusively form linear products. Although the active metalacyclic catalyst in these reactions was not formed prior to addition of the reagents, the high reactivity of the combination of  $[\text{Ir}(\text{COD})\text{Cl}]_2$  and ligand **L2** led to acceptable rates; the organozinc reagent might serve as the base to form the active metalacyclic catalyst (Scheme 14).

Nitrogen heterocycles undergo C-allylation in the presence of metalacyclic iridium-phosphoramidite catalysts. Recently, Yu and coworkers reported the C-allylation of indoles with the iridium catalyst derived from **L1** [86] and from phosphoramidites containing 2-methylindoline- and 2-methyl-1,2,3,4-tetrahydroquinoline as the amino group (Table 4) [87]. No N-allylation was reported. However,



**Scheme 14** Addition of aryl zinc reagents to allylic carbonates

**Table 4** C-allylation of indoles

Entry	R <sup>1</sup>	R <sup>2</sup>	L	Yield (%)	ee (%)
1	Ph	H	<b>L1</b>	49	85
2	4-(MeO)C <sub>6</sub> H <sub>4</sub>	5-MeO	<b>L1</b>	85	89
3	4-(CF <sub>3</sub> )C <sub>6</sub> H <sub>4</sub>	H	<b>L1</b>	63	87
4	nBu	H	<b>L1</b>	55	85
5	4-(CF <sub>3</sub> )C <sub>6</sub> H <sub>4</sub>	H	<b>L6</b>	41	65

*N*-allylation of indoles and other azoles in the presence of iridium catalysts has been reported recently and is discussed in Sect. 5.3.


### 4.3.2 Allylic Substitution with Nitrogen Nucleophiles

#### Allylic Substitution with Amines and Anilines


As previously discussed, activation of the iridium-phosphoramidite catalyst before addition of the reagents allows less basic nitrogen nucleophiles to be used in iridium-catalyzed allylic substitution reactions [70, 88]. Arylamines, which do not react with allylic carbonates in the presence of the combination of **L1** and [Ir(COD)Cl]<sub>2</sub> as catalyst, form allylic amination products in excellent yields and selectivities when catalyzed by complex **1a** generated in situ (Scheme 15). The scope of the reactions of aromatic amines is broad. Electron-rich and electron-neutral aromatic amines react with allylic carbonates to form allylic amines in high yields and excellent regio- and enantioselectivities as do hindered *ortho*-substituted aromatic amines. Electron-poor aromatic amines require higher catalyst loadings, and the products from reactions of these substrates are formed with lower yields and selectivities.

#### Ammonia, Imides, Carbamates, and Sulfonamides

A diverse group of secondary and tertiary amines are readily synthesized from the reaction of primary and secondary amines with allylic carbonates in the presence of preformed iridium metalacycles, but the direct synthesis of primary amines via iridium-catalyzed allylic amination requires the use of ammonia as a nucleophile. The asymmetric allylation of ammonia had not been reported until very recently, and it is not a common reagent in other metal-catalyzed reactions. Nonetheless, Hartwig and coworkers developed the reactions of ammonia with allylic carbonates in the presence of **1a** generated in situ [89]. Reactions conducted in the initial work led exclusively to the products from diallylation (Scheme 16). Further advances in



**Scheme 15** Amination of allylic carbonates with arylamines enabled by preactivated catalyst



**Scheme 16** Allylic substitution with ammonia to form diallylamines

**Table 5** Allylic substitution with ammonia equivalents



Entry	R	Nu	L	Yield (%)	b:l	ee (%)	Time (h)	Reference
1	Ph	<i>o</i> -NsNH <sub>2</sub>	<b>L2</b>	92	90:10	96	3	[91]
2	Ph	phthalimide	<b>L2</b>	95	96:4	98	2.5	[91]
3	Ph	HN(Boc)(CHO)	<b>L2</b>	96	98:2	99	0.7	[91]
4	Ph	NaN(Boc) <sub>2</sub>	<b>L2</b>	80	97:3	99	0.7	[91]
5 <sup>a</sup>	Ph	HN(Boc) <sub>2</sub>	<b>L2</b>	90	99:1	99	1	[94]
6	4-CF <sub>3</sub> (C <sub>6</sub> H <sub>4</sub> )	LiN(Boc) <sub>2</sub>	<b>L7</b>	76	85:15	92	4	[89]
7	4-MeO(C <sub>6</sub> H <sub>4</sub> )	KTFAc	<b>L7</b>	93	nd	98	12	[89]

<sup>a</sup>Conducted at 50°C

catalyst development led to the selective formation of the monoallylation product, and these reactions are discussed in Sect. 5.3 after introduction of these catalysts that were refined further.

Protected primary allylic amines were generated from allylic carbonates and ammonia equivalents. Iridium-catalyzed allylic substitution has now been reported with sulfonamides [90, 91], imides [89, 91–93], and trifluoroacetamide [89] to form branched, protected, primary allylic amines (Table 5). When tested, yields and selectivities were highest from reactions catalyzed by complexes derived from **L2**. Reactions of potassium trifluoroacetamide and lithium di-*tert*-butyliminodicyclohexylidene were conducted with catalysts derived from the simplified ligand **L7**. Reactions of nosylamide and trifluoroacetamide form singly-protected amine products. The other ammonia equivalents lead to the formation of doubly protected allylic amine products, but one protecting group can be removed selectively, except when the product is derived from phthalimide.


Allylic substitution with free carbamate nucleophiles was not accomplished until advanced catalyst precursors were developed. However, products from substitution with carbamate nucleophiles were generated by a decarboxylative allylation. In this process, the imidodicarbonate was shown to undergo decarboxylation to form the carbamate nucleophile, and reaction of the resulting carbamate with the  $\pi$ -allyliridium intermediate formed branched allylic carbamate products (Scheme 17) [92, 95].

**Scheme 17** Intramolecular decarboxylative allylic amidation**Scheme 18** Allylic substitution reactions between copper alkoxides and allylic carbonates


### 4.3.3 Allylic Substitution with Oxygen Nucleophiles

Iridium catalysts generated from modified phosphoramidite ligands prior to addition of the reagents were applied to the allylation of oxygen nucleophiles. As discussed earlier, iridium catalysts derived from  $[\text{Ir}(\text{COD})\text{Cl}]_2$  and **L1** readily catalyze the addition of phenoxides to linear allylic carbonates to form branched allyl aryl ethers with high enantioselectivity. More challenging has been the development of asymmetric allylic substitutions with alkoxide nucleophiles because of their high basicity. The addition of copper salts moderates the basicity of the alkoxides by forming softer copper alkoxides. Thus, Shu and Hartwig developed conditions for asymmetric allylic etherification with copper alkoxides in the presence of iridium-phosphoramidite catalysts. Analogous reactions of alkoxide nucleophiles catalyzed by other transition metals have not been reported to occur with high yield and enantioselectivity [96]. Copper alkoxides, both with and without branching  $\alpha$  to the oxygen, were shown to add to electron-rich, electron-neutral, and electron-poor cinnamyl carbonates, as well as alkyl-substituted allylic carbonates. Branched products formed in high yields and enantioselectivities (Scheme 18). Reactions conducted with ligand **L3** occurred in higher yields, branched-to-linear ratios and enantioselectivities than those conducted with ligand **L1**.

Ueno and Hartwig subsequently improved upon this method by developing conditions for allylic alkoxylation that do not require copper additives [97]. In reactions conducted with **1a** generated in situ as catalyst and  $\text{K}_3\text{PO}_4$  as base,



**Scheme 19** Allylic substitution reactions between aliphatic alcohols and allylic carbonates



**Scheme 20** Reactions of potassium triethylsiloxide with allylic carbonates to form branched allylic alcohols [98]

aliphatic alcohols reacted with cinnamyl and alkyl-substituted allylic acetates to form allylic ethers in high yield and selectivity (Scheme 19). An achiral vinyl ether was formed in these reactions from isomerization of the product, but the addition of 1-phenyl-1-propyne minimized this isomerization, presumably by poisoning an isomerization catalyst that forms during the reaction. This method can also be used to synthesize protected allylic alcohols from cinnamyl acetate and TBSOH, which is related to an earlier method developed by Carreira et al. that uses potassium triethylsiloxide (TESOK) as a nucleophile for the formation of branched allylic alcohols [98]. TESOK adds to *tert*-butyl allylic carbonates in the presence of catalytic amounts of **1a** generated in situ (Scheme 20).

#### 4.3.4 Intramolecular Allylic Substitution Catalyzed by Metalacyclic Iridium Phosphoramidite Complexes


Several types of intramolecular allylic substitution reactions of carbon, nitrogen, and oxygen nucleophiles catalyzed by metalacyclic iridium phosphoramidite complexes have been reported. Intramolecular allylic substitution is much faster than the competing intermolecular process when conducted in the presence of iridium catalysts. Thus, conditions involving high dilution are not required. Intramolecular

allylic substitution has been applied to form products that are precursors to natural products via allylic alkylation [79, 99], amination [53, 78], and etherification [78].

## 5 New Mechanistic Insights: The Isolation of Catalytic Intermediates

### 5.1 General Mechanism of Metal-Catalyzed Allylic Substitution

Studies of the mechanism of palladium-catalyzed allylic substitution have been aided by the isolation and characterization of allylpalladium complexes that are catalytic intermediates. Related mechanistic studies of iridium-catalyzed allylic substitution were, until recently, hampered by the inability to observe catalytically active allyliridium species. Based on the mechanism of palladium-catalyzed allylic substitution, iridium-catalyzed allylic substitution was assumed to occur by the series of steps shown in Scheme 21, but information on the relative rates of individual steps, the structures of intermediates, and the origins of enantioselectivity was limited. Soon after the discovery of the iridium-phosphoramidite system that catalyzes allylic substitution with high enantioselectivity, the active catalyst was shown to consist of a metalacyclic iridium fragment that is derived from the phosphoramidite ligand and that contains a coordinated cyclooctadiene. However, this fragment had only been isolated as a species coordinated to an additional ligand, such as a **L1** (**1a**) or ethylene (**1b**). This five-coordinate, 18-electron species presumably lies off of the actual catalytic cycle and requires dissociation of the phosphoramidite or ethylene ligand to form a 16-electron intermediate that lies on the catalytic cycle. No other species shown to lie on the catalytic cycle had been



**Scheme 21** General mechanism for metal-catalyzed allylic substitution




observed or isolated. Attempts to generate a  $\pi$ -allyliridium(III) intermediate from the species containing **L1** or ethylene as ligand led to complexes that neither catalyzed allylic substitution nor produced the expected branched allylation products upon exposure to a stoichiometric quantity of nucleophile [52].

## 5.2 Isolation and Study of Intermediates in Allylic Substitution Reactions Catalyzed by Metalacyclic Iridium-Phosphoramidite Complexes

### 5.2.1 Identification and Isolation of the Catalyst Resting State

Marković and Hartwig isolated and characterized the first intermediate in iridium-catalyzed allylic substitution [100]. They isolated the metalacyclic iridium-phosphoramidite fragment containing COD and the olefinic portion of *N*-(1-phenylallyl)aniline, the product of the allylic substitution reaction between cinnamyl carbonate and aniline (**5** in Scheme 22). This complex containing the product of allylic substitution was first detected by  $^{31}\text{P}$  NMR spectroscopy during catalytic reactions. It was then isolated, prepared independently, and shown to be chemically and kinetically competent to be an intermediate in allylic substitutions.

In addition to isolating complex **5**, Marković and Hartwig performed kinetic studies on the amination of methyl cinnamyl carbonate with aniline. The proposed mechanism involves reversible dissociation of product, reversible, endothermic oxidative addition of the allylic carbonate to form a  $\pi$ -allyliridium species, and irreversible nucleophilic attack on the  $\pi$ -allyliridium intermediate, as depicted in



**Scheme 22** Catalytic cycle for the reaction of aniline with methyl cinnamyl carbonate catalyzed by a metalacyclic iridium-phosphoramidite complex

Scheme 22. The rate equation for this mechanism is described in (1). The authors determined that the reaction is first-order in allylic carbonate, aniline and catalyst, and inverse first-order in allylamine product. These results are consistent with the proposed mechanism. Thus, iridium-catalyzed allylic substitution is inhibited by product. In addition, the formation of the allyliridium intermediate is disfavored.


$$\text{rate} = \frac{d[\text{P}]}{dt} = \frac{K_1 K_2 k_3 [\text{S}][\text{A}][\text{Ir}]_{\text{tot}}}{[\text{P}]} \quad (1)$$

### 5.2.2 Isolation of $\pi$ -Allyliridium Intermediates


Additional mechanistic insights were gained when Hartwig and coworkers isolated and characterized the first  $\pi$ -allyl complexes that are chemically and kinetically competent to be intermediates in iridium-catalyzed allylic substitution [46]. These complexes were prepared independently from allylic electrophiles that are more reactive than allylic carbonates. The isolation and structural characterization of these species provided a detailed view into the origins of enantioselectivity.

The allyliridium species were formed from allylic chlorides, rather than allylic carbonates. Treatment of complex metalacyclic iridium complex **1b** with allylic methyl carbonates led to no apparent reaction and no accumulation of a new iridium species. The lack of observable product from this reaction is consistent with prior kinetic data indicating that the formation of an allyliridium intermediate from a resting state containing ethylene or allylic amine as ligand was thermodynamically unfavorable. Although the reaction of allylic carbonates with metalacyclic iridium phosphoramidite precursors is thermodynamically unfavorable, the reaction of allylic chlorides with **1b** formed a new species, as determined by  $^{31}\text{P}$  NMR spectroscopy. This new species was trapped by addition of  $\text{AgOTf}$ , isolated as single crystals, and shown to be the intermediate allyliridium species (Scheme 23). The parent  $\pi$ -allyliridium complex **6a** and the analogous crotyl complex **6b** were both isolated and characterized by X-ray diffraction.

Allyliridium complexes **6a** and **6b** reacted with a series of nucleophiles to form products in yields that are similar to those of allylic substitution reactions catalyzed




**Scheme 23** Synthesis of  $\eta^3$ -allyliridium intermediates



**Scheme 24** The reaction of allyl complexes with benzylamine

**Fig. 5** ORTEP Diagram of Crotyl Complex **6b** [46]



by **1b**. Moreover, nucleophilic attack on the crotyl complex **6b** also resulted in the formation of products with branched-to-linear ratios and enantioselectivities that are similar to those of catalytic reactions with methallyl methyl carbonate (Scheme 24).

Information on the origins of enantioselection by the iridium catalyst can be gleaned from the structure of the crotyl complex **6b** (Fig. 5). The phosphoramidite and COD ligands on iridium create a steric environment that constrains the orientation of the allyl ligand. First, the allyl unit binds in an orientation in which the methyl substituent on the allyl unit lies in solution away from the other ligands. Second, coordination to one face of the allyl ligand is favored over coordination to the other because it minimizes interaction between the COD ligand and the central carbon of the allyl group (C38). Comparison of the configuration of the isolated allyl complex with that of the isolated organic products indicates that nucleophilic attack occurs anti to the metal center [46].

### 5.3 Reaction Development Based on Mechanistic Insights


These recent mechanistic studies have provided the foundation for the most recent work that has expanded the scope of iridium-catalyzed allylic substitution. The synthesis and characterization of the ethylene-bound complex **1b** resulted directly

from studies on the synthesis of catalytic intermediates. This stable adduct of the catalytically active metalacyclic fragment is a convenient, single-component catalyst precursor that does not require an activator to generate the active catalyst. With **1b** this activation occurs by dissociation of the coordinated ethylene.


In contrast, reactions catalyzed by **1a** were typically conducted with added  $[\text{Ir}(\text{COD})\text{Cl}]_2$  to trap the  $\kappa^1$ -phosphoramidite ligand after dissociation, and thereby, to leave the unsaturated active catalyst. Under these conditions, as much as half of the iridium in the system is present in an inactive acyclic species. In contrast, when ethylene adduct **1b** is used as the catalyst, all of the iridium belongs to the active metalacyclic species. Hartwig and coworkers have recently taken advantage of the increased availability of the active catalyst generated from **1b** to develop new allylic substitution reactions. These new processes include the reactions of carbamates, nitrogen heterocycles, and ammonia.

Previously, iridium-catalyzed reactions to form *N*-allyl carbamates required the use of allylic imidodicarbonates as reagents that undergo decarboxylative allylation, and the scope of these reactions encompassed only allylic benzyloxycarbamates. Direct allylic substitution with commercially available carbamate nucleophiles would allow the formation of a variety of protected primary amine products without the need to synthesize imidodicarbonates. Hartwig and coworkers have shown that the ethylene complex **2b**, which is the 2-anisyl variant of **1b**, catalyzes the reaction between *tert*-butyl allylic carbonates and  $\text{BocNH}_2$ , fluorenyloxycarbamate ( $\text{FmocNH}_2$ ), benzyloxycarbamate ( $\text{CbzNH}_2$ ), 2,2,2-trichloroethoxycarbamate ( $\text{TrocNH}_2$ ), 2-trimethylsilylethoxycarbamate ( $\text{TeocNH}_2$ ), and 2-oxazolidinone (Scheme 25) [101]. Yields and selectivities were high, and the substrate scope encompasses a range of aryl- and alkyl-substituted allylic carbonates. Reactions of  $\text{BocNH}_2$  occurred without the use of exogenous base.


Complex **2b** also catalyzes the addition of imidazoles, benzimidazoles, and several purines to methyl allylic carbonates (Scheme 26) [76]. These reactions occurred with good yields and excellent enantioselectivities. *N*-substituted heterocycles are found in a number of compounds that exhibit a range of biological activity. The syntheses of several pharmacologically active compounds were conducted with yields that were improved over those published previously.



**Scheme 25** Allylic substitution with carbamate nucleophiles




**Scheme 26** Allylic substitution with *N*-heterocycles



**Scheme 27** Allylic substitution with ammonia catalyzed by **1a** and **1b**

The use of ethylene adduct **1b** is particularly important when the species added to activate catalyst **1a** is incompatible with one of the reaction components. Iridium-catalyzed monoallylation of ammonia requires high concentrations of ammonia, but these conditions are not compatible with the additive [Ir(COD)Cl]<sub>2</sub> because this complex reacts with ammonia [102]. Thus, a reaction between ammonia and ethyl cinnamyl carbonate catalyzed by ethylene adduct **1b** produces the monoallylation product in higher yield than the same reaction catalyzed by **1a** and [Ir(COD)Cl]<sub>2</sub> (Scheme 27). Ammonia reacts with a range of allylic carbonates in the presence of **1b** to form branched primary allylic amines in good yield and high enantioselectivity (Scheme 28). Quenching these reactions with acyl chlorides or anhydrides leads to a one-pot synthesis of branched allylic amides that are not yet directly accessible by metal-catalyzed allylation of amides.

The high concentration of ammonia that is required for selective formation of primary allylic amines suggests that ammonia is much less reactive toward nucleophilic attack on an allyl ligand than the primary allylic amine products. The reactivity of different nucleophiles toward allyl ligands was assessed by treating the allyliridium complex **6a** with ammonia and 1-phenylprop-2-en-1-amine. No product from allylation of ammonia was detected from this reaction. In addition, the reaction of an allylic carbonate with the combination of ammonia and 1-phenylprop-2-en-1-amine in the presence of **1b** did not produce detectable amounts of the product from allylation of ammonia. These results confirm that the primary amine is much more



**Scheme 28** Selective monoallylation of ammonia catalyzed by **1b**


reactive toward the allyliridium species than is ammonia. This greater reactivity of the allylamine explains why iridium-catalyzed allylic substitutions conducted with low concentrations of ammonia selectively form diallylation products over monoallylation products. However, the stability of catalyst **1b** toward ammonia allows the reaction to be conducted with high concentrations of ammonia and to form the primary allylic amine selectively.

## 6 Challenges in Iridium-Catalyzed Allylic Substitution

The scope of reactions catalyzed by metalacyclic iridium-phosphoramidite complexes is remarkably broad, but reactions with some substrates, such as allylic alcohols, prochiral nucleophiles, branched allylic esters, and highly substituted allylic esters, that would form synthetically valuable products or would lead to simpler synthesis of reactants occur with low yields and selectivities. In addition, iridium-catalyzed allylic substitution reactions are sensitive to air and water and must be conducted with dry solvents under an inert atmosphere. Several advances have helped to overcome some, but not all of these challenges.

### 6.1 Allylic Substitution of Allylic Alcohols

Most iridium-catalyzed allylic substitutions have been performed with allylic esters, which are typically synthesized from allylic alcohols. Reactions of allylic alcohols as electrophiles would alleviate the need to prepare the esters from the alcohol. In a few cases, however, iridium catalyzed allylic substitutions have been conducted with allylic alcohols as the electrophile. As discussed earlier in this




**Scheme 29** Allylic substitution with linear allylic alcohols activated by Lewis acids

chapter, Takeuchi found that allylic alkylation reactions between malonate nucleophiles and allylic alcohols occur via transesterification between the allylic alcohol and one equivalent of the nucleophile, thereby producing a more reactive leaving group. Thus, allylic substitution occurs in these cases with the allylic alcohol via the formation of an allylic ester.


More recently, Hartwig and coworkers reported iridium-catalyzed, asymmetric aminations of allylic alcohols in the presence of Lewis acid activators [103]. The addition of molecular sieves and  $\text{Nb(OEt)}_5$  or catalytic amounts of  $\text{BPh}_3$  activated the allylic alcohol sufficiently to allow allylic amination reactions to occur in high yield, branched-to-linear selectivity, and enantioselectivity (Scheme 29). Without the activators, only trace amounts of product were observed.

## 6.2 Diastereoselectivity of Iridium-Catalyzed Allylic Substitution Reactions

Control of diastereoselectivity in allylic substitution reactions remains a significant challenge. Nucleophilic attack by a prochiral nucleophile can produce linear allylic substitution products containing one stereogenic center or branched products containing two stereogenic centers. The diastereoselectivity of reactions of prochiral nucleophiles catalyzed by iridium complexes has been low. For example, nitroalkanes [83], ethyl nitroacetates [83], and malonic amides [104] undergo allylic substitution catalyzed by the metalacyclic iridium complexes in high yield, branched-to-linear selectivity, and enantioselectivity, but the alkylation products form as 1:1 mixtures of diastereomers (Scheme 30). However, the reactions of allylic esters with glycine derivatives reported by Kanayama and described in Sect. 3 occurred with good diastereoselectivity in the presence of iridium-phosphite catalysts (Sect. 3) [56, 57].



**Scheme 30** Allylic alkylation with ethyl nitroacetate



**Scheme 31** Sequential Pd-catalyzed isomerization and Ir-catalyzed allylic substitution


### 6.3 Allylic Substitution of Branched Allylic Esters

Allylic substitution reactions catalyzed by metalacyclic iridium-phosphoramidite complexes form branched products from linear allylic esters with high regioselectivity. However, reactions with racemic, branched allylic esters would be particularly valuable because they are readily accessible from a wide array of aldehydes and vinylmagnesium halides. However, iridium-catalyzed allylic substitution reactions of branched allylic esters have so far occurred with low enantioselectivities [45, 75].


Hartwig and coworkers reported an approach to address this limitation involving tandem catalytic reactions. In this tandem process, sequential palladium-catalyzed isomerization of the branched isomer to the linear isomer, followed by iridium-catalyzed allylic substitution leads to the branched product with high enantiomeric excess [105]. More specifically, treatment of branched allylic esters with catalytic amounts of the combination of Pd(dba)<sub>2</sub> and PPh<sub>3</sub> led to rapid isomerization of the branched allylic ester to the linear isomer, and the linear isomer underwent allylic substitution after addition of the iridium catalyst and nucleophile (Scheme 31).

Carreira et al. reported the kinetic resolution of branched allylic carbonates catalyzed by an iridium complex derived from a chiral [2.2.2]-bicyclooctadiene [48]. Reactions of allylic carbonates with phenol were run to 50% conversion of the carbonate, leaving unreacted allylic carbonate in high enantiomeric excess (Scheme 32). The phenyl ether products were also isolated in mid-to-high enantiomeric excess.





**Scheme 32** Kinetic resolution of branched allylic carbonates




**Scheme 33** The reaction of sulfamic acid with a branched allylic alcohol


More recently, Carreira and coworkers reported the reactions of sulfamic acid with branched allylic alcohols to form branched primary allylic amines [106]. One enantioselective reaction was disclosed, and the product was isolated in 70% ee when using a 5*H*-dibenzo[*b*, *f*]azepine-derived phosphoramidite with  $[\text{Ir}(\text{COE})_2\text{Cl}]_2$  (Scheme 33). This reaction occurs under acidic conditions, which is incompatible with the metalacyclic catalysts.

#### 6.4 Allylic Substitution with Highly Substituted Allylic Esters

No examples have been reported of enantioselective, iridium-catalyzed allylic substitutions of linear allylic esters to generate 1,1-disubstituted or 2-substituted  $\pi$ -allyl intermediates. Takeuchi published reactions in which the proposed allyliridium intermediates are 1,1- or 1,3-disubstituted, but these substrates have not been shown to undergo reactions catalyzed by chiral iridium complexes. No reactions of 1,2-disubstituted substrates have been published (Scheme 34).



**Scheme 34** The reaction of highly substituted allylic carbonates have not been catalyzed by metalacyclic iridium-phosphoramidite complexes



**Scheme 35** Air-stable allylic substitution catalyzed by  $[\text{Ir}(\text{dbcot})\text{Cl}]_2$  and **L2**

## 6.5 Stability of Iridium-Phosphoramidite Catalysts to Air and Water

Except for one recent example, all iridium-catalyzed allylic substitution reactions have been performed under an inert atmosphere with dry solvent and reagents. The iridium metalacycle is sensitive to protonation, which opens the metalacycle and results in the formation of a less-active complex containing a  $\kappa^1$ -phosphoramidite ligand. A paper by Helmchen et al. addressed this issue [107]. Nearly all iridium catalysts used for allylic substitution consist of an iridium fragment chelated by COD. In the presence of a catalyst containing dibenzo[*a,e*]cyclooctatetraene (dbcot) in place of COD, allylic substitution reactions occur in air with results that are comparable to those of reactions performed under an inert atmosphere (Scheme 35).

## 7 Conclusion

Takeuchi's initial report on iridium-catalyzed allylic substitution demonstrated that the allylation of carbon nucleophiles forms branched, racemic substitution products. Chiral catalysts were quickly developed that led to methods to prepare the

products of allylic substitution at the more substituted position of mono-substituted allylic carbonates with high enantioselectivities. Since these initial reports, the scope of iridium-catalyzed asymmetric allylic substitution has grown to include reactions of a wide range of carbon, nitrogen, and oxygen nucleophiles. Substitution products are now typically formed in excellent enantiomeric excess. Only recently have studies begun to reveal the mechanism of iridium-catalyzed allylic substitution, but these mechanistic insights are now aiding in the development of new catalyst precursors that further improve reaction scope and catalyst activity.

## References

1. Takeuchi R, Kashio M (1997) *Angew Chem Int Ed Engl* 36:263
2. Tsuji J, Takahash H, Morikawa M (1965) *Tetrahedron Lett* 4387
3. Walker WE, Manyik RM, Atkins KE, Farmer ML (1970) *Tetrahedron Lett* 3817
4. Atkins KE, Walker WE, Manyik RM (1970) *Tetrahedron Lett* 3821
5. Shryne TS, Smutny EJ, Stevenson DP (1970) US Patent 3,493,617
6. Hata G, Takahash K, Miyake A (1970) *J Chem Soc D* 1392
7. Trost BM, Van Vranken DL (1996) *Chem Rev* 96:395
8. Trost BM, Lautens MJ (1982) *Am Chem Soc* 104:5543
9. Trost BM, Lautens MJ (1983) *Am Chem Soc* 105:3343
10. Trost BM, Lautens M (1983) *Organometallics* 2:1687
11. Minami I, Shimizu I, Tsuji J (1985) *J Organomet Chem* 296:269
12. Trost BM, Lautens MJ (1987) *Am Chem Soc* 109:1469
13. Trost BM, Lautens M (1987) *Tetrahedron* 43:4817
14. Faller JW, Lambert C, Mazzieri MR (1990) *J Organomet Chem* 383:161
15. Trost BM, Merlic CA (1990) *J Org Chem* 55:1127
16. Trost BM, Merlic CA (1990) *J Am Chem Soc* 112:9590
17. Dvořák D, Starý I, Kočovský P (1995) *J Am Chem Soc* 117:6130
18. Dvořáková H, Dvořák D, Šrogl J, Kočovský P (1995) *Tetrahedron Lett* 36:6351
19. Trost BM, Hung M-H (1983) *J Am Chem Soc* 105:7757
20. Trost BM, Hung MHJ (1984) *J Am Chem Soc* 106:6837
21. Trost BM, Tometzki GB, Hung MH (1987) *J Am Chem Soc* 109:2176
22. Lehmann J, Lloyd-Jones GC (1995) *Tetrahedron* 51:8863
23. Frisell H, Åkermark B (1995) *Organometallics* 14:561
24. Lloyd-Jones GC, Pfaltz A (1995) *Angew Chem Int Ed Engl* 34:462
25. Xu YY, Bo Z (1987) *J Org Chem* 52:974
26. Silverman GS, Strickland S, Nicholas KM (1986) *Organometallics* 5:2117
27. Roustan JL, Merour JY, Houlihan F (1979) *Tetrahedron Lett* 3721
28. Roustan JLA, Houlihan F (1979) *Can J Chem Rev Can Chim* 57:2790
29. Zhang SW, Mitsudo T, Kondo T, Watanabe Y (1993) *J Organomet Chem* 450:197
30. Kondo T, Ono H, Satake N, Mitsudo T, Watanabe Y (1995) *Organometallics* 14:1945
31. Tsuji J, Minami I, Shimizu I (1984) *Tetrahedron Lett* 25:5157
32. Bricout H, Carpentier J-F, Mortreux A (1996) *Tetrahedron Lett* 37:6105
33. Mizojiri R, Kobayashi Y (1995) *J Chem Soc Perkin Trans* 1:2073
34. Trost BM, Spagnol MD (2083) *J Chem Soc Perkin Trans* 1:1995
35. Alvarez E, Cuvigny T, Julia M (1988) *J Organomet Chem* 339:199
36. Cuvigny T, Julia M (1986) *J Organomet Chem* 317:383
37. Bricout H, Carpentier J-F, Mortreux A (1995) *J Chem Soc Chem Commun* 1863
38. Brown JM, Macintyre JE (1985) *J Chem Soc Perkin Trans* 2:961

39. Evans PA, Robinson JE, Nelson JD (1999) *J Am Chem Soc* 121:6761
40. Evans PA, Robinson JE (1999) *Org Lett* 1:1929
41. Takeuchi R, Kashio M (1998) *J Am Chem Soc* 120:8647
42. Takeuchi R, Shiga N (1999) *Org Lett* 1:265
43. Takeuchi R, Tanabe K (2000) *Angew Chem Int Ed Engl* 39:1975
44. Takeuchi R, Ue N, Tanabe K, Yamashita K, Shiga N (2001) *J Am Chem Soc* 123:9525
45. Bartels B, García-Yebra C, Rominger F, Helmchen G (2002) *Eur J Inorg Chem* 2569
46. Madrahimov ST, Markovic D, Hartwig JF (2009) *J Am Chem Soc* 131:7228
47. Miyabe H, Matsumura A, Yoshida K, Yamauchi M, Takemoto Y (2004) *Synlett* 2123
48. Fischer C, Defieber C, Suzuki T, Carreira EM (2004) *J Am Chem Soc* 126:1628
49. Nomura N, Komiyama S, Kasugai H, Saba M (2008) *J Am Chem Soc* 130:812
50. Janssen JP, Helmchen G (1997) *Tetrahedron Lett* 38:8025
51. Bartels B, Helmchen G (1999) *Chem Commun* 741
52. García-Yebra C, Janssen JP, Rominger F, Helmchen G (2004) *Organometallics* 23:5459
53. Welter C, Koch O, Lipowsky G, Helmchen G (2004) *Chem Commun* 896
54. Fuji K, Kinoshita N, Tanaka K, Kawabata T (1999) *Chem Commun* 2289
55. Kinoshita N, Marx KH, Tanaka K, Tsubaki K, Kawabata T, Yoshikai N, Nakamura E, Fuji K (2004) *J Org Chem* 69:7960
56. Kanayama T, Yoshida K, Miyabe H, Kimachi T, Takemoto Y (2003) *J Org Chem* 68:6197
57. Kanayama T, Yoshida K, Miyabe H, Takemoto Y (2003) *Angew Chem Int Ed Engl* 42:2054
58. Miyabe H, Matsumura A, Moriyama K, Takemoto Y (2004) *Org Lett* 6:4631
59. Miyabe H, Yoshida K, Yamauchi M, Takemoto Y (2005) *J Org Chem* 70:2148
60. Miyabe H, Matsumura A, Yoshida K, Takemoto Y (2009) *Tetrahedron* 65:4464
61. Miyabe H, Yoshida K, Reddy VK, Takemoto Y (2009) *J Org Chem* 74:305
62. Nemoto T, Sakamoto T, Matsumoto T, Hamada Y (2006) *Tetrahedron Lett* 47:8737
63. Nemoto T, Sakamoto T, Fukuyama T, Hamada Y (2007) *Tetrahedron Lett* 48:4977
64. Kimura M, Uozumi Y (2007) *J Org Chem* 72:707
65. Ohmura T, Hartwig JF (2002) *J Am Chem Soc* 124:15164
66. Lipowsky G, Helmchen G (2004) *Chem Commun* 116
67. Feringa BL, Pineschi M, Arnold LA, Imbos R, de Vries AHM (1997) *Angew Chem Int Ed Engl* 36:2620
68. López F, Ohmura T, Hartwig JF (2003) *J Am Chem Soc* 125:3426
69. Bartels B, García-Yebra C, Helmchen G (2003) *Eur J Org Chem* 1097
70. Kiener CA, Shu CT, Incarvito C, Hartwig JF (2003) *J Am Chem Soc* 125:14272
71. Lipowsky G, Miller N, Helmchen G (2004) *Angew Chem Int Ed Engl* 43:4595
72. Leitner A, Shu CT, Hartwig JF (2005) *Org Lett* 7:1093
73. Tissot-Croset K, Polet D, Alexakis A (2004) *Angew Chem Int Ed Engl* 43:2426
74. Leitner A, Shekhar S, Pouy MJ, Hartwig JF (2005) *J Am Chem Soc* 127:15506
75. Polet D, Alexakis A, Tissot-Croset K, Corminboeuf C, Ditrich K (2006) *Chem Eur J* 12:3596
76. Stanley LM, Hartwig JF (2009) *J Am Chem Soc* 131:8971
77. Leitner A, Shu CT, Hartwig JF (2004) *Proc Natl Acad Sci USA* 101:5830
78. Welter C, Dahnz A, Brunner B, Streiff S, Dübon P, Helmchen G (2005) *Org Lett* 7:1239
79. Gnamm C, Forster S, Miller N, Brödner K, Helmchen G (2007) *Synlett* 790
80. Graening T, Hartwig JF (2005) *J Am Chem Soc* 127:17192
81. Weix DJ, Hartwig JF (2007) *J Am Chem Soc* 129:7720
82. He H, Zheng X-J, Li Y, Dai L-X, You S-L (2007) *Org Lett* 9:4339
83. Dahnz A, Helmchen G (2006) *Synlett* 697
84. Alexakis A, El Hajjaji S, Polet D, Rathgeb X (2007) *Org Lett* 9:3393
85. Polet D, Rathgeb X, Falcicola CA, Langlois JB, El Hajjaji S, Alexakis A (2009) *Chem Eur J* 15:1205
86. Liu WB, He H, Dai LX, You S-L (2008) *Org Lett* 10:1815
87. Liu WB, He H, Dai LX, You SL (2009) *Synthesis* 2076
88. Shu CT, Leitner A, Hartwig JF (2004) *Angew Chem Int Ed Engl* 43:4797

89. Pouy MJ, Leitner A, Weix DJ, Ueno S, Hartwig JF (2007) *Org Lett* 9:3949
90. Weihofen R, Dahnz A, Tverskoy O, Helmchen GN (2005) *Chem Commun* 3541
91. Weihofen R, Tverskoy E, Helmchen G (2006) *Angew Chem Int Ed Engl* 45:5546
92. Singh OV, Han H (2007) *J Am Chem Soc* 129:774
93. Spiess S, Berthold C, Weihofen R, Helmchen G (2007) *Org Biomol Chem* 5:2357
94. Singh OV, Han H (2007) *Tetrahedron Lett* 48:7094
95. Singh OV, Han HS (2007) *Org Lett* 9:4801
96. Shu CT, Hartwig JE (2004) *Angew Chem Int Ed Engl* 43:4794
97. Ueno S, Hartwig JF (2008) *Angew Chem Int Ed Engl* 47:1928
98. Lyothier I, Defieber C, Carreira EM (2006) *Angew Chem Int Ed Engl* 45:6204
99. Streiff S, Welter C, Schelwies M, Lipowsky G, Miller N, Helmchen G (2005) *Chem Commun* 2957
100. Marković D, Hartwig JF (2007) *J Am Chem Soc* 129:11680
101. Weix DJ, Marković D, Ueda M, Hartwig JF (2009) *Org Lett* 11:2944
102. Pouy MJ, Stanley LM, Hartwig JF (2009) *J Am Chem Soc* 131:11312
103. Yamashita Y, Gopalarathnam A, Hartwig JF (2007) *J Am Chem Soc* 129:7508
104. Schelwies M, Dübon P, Helmchen G (2006) *Angew Chem Int Ed Engl* 45:2466
105. Shekhar S, Trantow B, Leitner A, Hartwig JF (2006) *J Am Chem Soc* 128:11770
106. Defieber C, Ariger MA, Moriel P, Carreira EM (2007) *Angew Chem Int Ed Engl* 46:3139
107. Spiess S, Welter C, Franck G, Taquet J-P, Helmchen G (2008) *Angew Chem Int Ed Engl* 47:7652

# Iridium-Catalyzed 1,3-Dipolar Cycloadditions

Daniel Carmona and Luis A. Oro

**Abstract** 1,3-Dipolar cycloaddition reactions (DCR) are atom-economic processes that permit the construction of heterocycles. Their enantioselective versions allow for the creation of up to four adjacent chiral centers in a concerted fashion. In particular, well-defined half-sandwich iridium (III) catalysts have been applied to the DCR between enals or methacrylonitrile with nitrones. Excellent yield and stereoselectivities have been achieved. Support for mechanistic proposals stems from the isolation and characterization of the true catalysts.

**Keywords** 1,3-Dipolar cycloadditions · Enals · Iridium catalysts · Methacrylonitrile · Nitrones

## Contents


1	Introduction .....	210
1.1	The Frontier Molecular Orbital Theory .....	211
2	DCR of Nitrones with Alkenes .....	212
2.1	DCR of Nitrones with Enals .....	212
3	DCR of Nitrones with Enals Catalyzed by Iridium Complexes .....	216
3.1	DCR of Nitrones with Methacrolein .....	217
3.2	DCR of Nitrones with Other $\alpha,\beta$ -Unsaturated Aldehydes .....	223
4	DCR of Nitrones with Methacrylonitrile Catalyzed by Iridium Complexes .....	225
4.1	Recycling Experiments .....	226
5	Conclusions .....	228
	References .....	228

## 1 Introduction


Cycloaddition reactions are a fundamental class of processes in synthetic chemistry. Within this class, the 1,3-dipolar cycloaddition reaction (DCR) has found extensive use as an efficient method for the synthesis of different heterocyclic compounds. These type of reactions involve the addition of a 1,3-dipole to a multiple  $\pi$ -bond system (dipolarophile) leading to five-membered heterocycles (Scheme 1) [1].

Huisgen in the 1960s undertook a systematic study that led to the generalization and classification of DCR [2] and to the proposal of a concerted synchronous mechanism (Scheme 2) [3]. However, Firestone considered the DCR to proceed via a stepwise diradical mechanism. Stereochemistry would be retained for short-lived diradical intermediates but may be destroyed for longer-lived diradicals (Scheme 3) [4]. Currently, the most widely accepted mechanism is a concerted process with both new bonds being formed at the same time but not necessarily to the same extent. Additionally, very recently, Houk et al. have found an unambiguous correlation between the calculated activation barriers and the distortion energies, the latter being “the energy required to distort the 1,3-dipole and the dipolarophile from their equilibrium geometries into the transition-state geometries without allowing any interaction between them” [5]. These findings provide a new way of understanding the DCR that cannot be neglected in accurate computational studies [6–8].


**Scheme 1** 1,3-Dipolar cycloaddition reaction



**Scheme 2** Concerted mechanism for DCR



**Scheme 3** Stepwise diradical mechanism for DCR



**Fig. 1** Sustmann's classification of DCR

### 1.1 The Frontier Molecular Orbital Theory

The most successful qualitative model for DCR is based on frontier molecular orbital (FMO) theory. According to this theory, reaction takes place by maximizing overlap between the HOMO and LUMO of the reagents. Sustman applied FMO theory to the reactivity of concerted DCR and classified them into three types depending on the interactions established between the FMO of the dipole and dipolarophile (Fig. 1) [9–12]. Type I reactions involve dipoles with high-lying HOMO and dipolarophiles with low-lying LUMO. The most favorable FMO interaction will imply transference of charge from the nucleophilic dipole to the electrophilic dipolarophile. In this case, the reaction rate may be increased, for example, introducing electron-withdrawing groups on the dipolarophile. By the contrary, type III reactions involve dipoles with low-lying LUMO and dipolarophiles with high-lying HOMO. The dominant FMO interaction will take place between the HOMO of the dipolarophile and the LUMO of the dipole. Reactions of this type can be accelerated by using dipolarophiles with electron-donating groups. Finally, type II reactions involve dipoles and dipolarophiles with similar energy gaps between the HOMO and LUMO. These reactions are called ambiphilic, and, in particular, reactions of nitrones with alkenes belong to this type. Introduction of electron-donating or electron-withdrawing substituents on the dipolarophile or the dipole can alter the relative FMO energies resulting in a change to type I or III reaction.

Interestingly, the presence of metallic Lewis acids in the reaction media may substantially affect reactions. When coordinating to the dipolarophile, the Lewis acid lowers the energy of its frontier orbitals. This lowering reduces the energy gap between the LUMO of the dipolarophile and the HOMO of the dipole (Fig. 2) leading to an acceleration of the cycloaddition. Similarly, coordination to the dipole reduces the energy difference between the HOMO of the dipolarophile and the LUMO of the dipole, this reduction resulting in an increment of the reaction rate. Notably, the efficiency of this Lewis acid interaction relies not only on the fastening of the reaction but also on the capability of metallic complexes to control the selectivities of the cycloaddition. Thus, a variety of Lewis acids based on metals have been employed as catalysts in stereoselective DCR often providing excellent results [13–22].







Fig. 2 Changes in FMO by coordination to a Lewis acid

Scheme 4 DCR between nitrones and alkenes





## 2 DCR of Nitrones with Alkenes

Over the last years, one of the most studied DCR has been the asymmetric version of the cycloaddition of nitrones with alkenes. This reaction leads to the construction of up to three contiguous asymmetric carbon centers (Scheme 4). The resulting five-membered isoxazolidine derivatives may be converted into amino alcohols, alkaloids, or  $\beta$ -lactams. Several chiral metal complexes have been used as catalysts for this process [13–15, 18–22]. However, the employment of iridium derivatives is very scarce.

### 2.1 DCR of Nitrones with Enals

Due to the presence of an electron-withdrawing group on the dipolarophile, these processes are classified as type I reactions. The process involves the transference of charge from the dipole to the dipolarophile. When catalyzed by metallic compounds, coordination of the dipolarophile is highly desired. Usually, coordination of a nitrone to the Lewis acid is more feasible than coordination of a carbonyl compound. For this reason, alkenes that enable a bidentate coordination to the Lewis acid, such as 3-alkenoyl-oxazolidinones (Scheme 5), have been frequently employed as a model system to study the metal-catalyzed 1,3-dipolar cycloaddition


**Scheme 5** 3-Alkenoyl-oxazolidinones**Scheme 6** DCR catalyzed by Binop-F complexes

of nitrones [14, 15, 18, 20]. In contrast, examples of one point binding catalysts for the activation of electron-deficient monofunctionalized alkenes, such as enals, are scarce [18, 20]. Only recently has it been possible to find a few examples in the literature.


In 2002, Kündig et al. [23, 24] developed catalytic DCR between diaryl nitrones and  $\alpha,\beta$ -unsaturated aldehydes in the presence of Binop-F iron and ruthenium complexes as chiral Lewis-acid catalysts (Scheme 6). The corresponding cycloadducts were obtained in good yields with complete *endo* selectivity and up to 94% ee. The isoxazolidine products were obtained as a mixture of regioisomers in molar ratios varying from 96:4 to 15:85. Experimental and computational data show that the regioselectivity correlates directly with the electronic properties of the nitron.

Yamada and coworkers [25–27] have employed chiral  $\beta$ -ketoiminato cobalt (III) complexes as catalysts for the same type of reaction. The cycloadducts were obtained with complete *endo* selectivity and in moderate to high ee (Scheme 7).

Complexes of nickel(II) or magnesium(II) with the chiral ligand DBFOX (Scheme 8) catalyze the DCR of nitrones with  $\alpha$ -alkyl- and acryloleins rendering preferentially the 5-carbaldehyde cycloadducts. However, the reactions with  $\alpha$ -bromoacrolein catalyzed by the zinc(II) complex of the same ligand afford isoxazoline-4-carbaldehydes. The corresponding cobalt(II) complex is also active for the cycloaddition between cyclopenten-1-carbaldehyde and diphenylnitron.




**Scheme 7** Cycloadditions catalyzed by  $\beta$ -ketoiminato complexes of Co(III)



**Scheme 8** DCR catalyzed by DBFOX complexes

The adducts are characterized as alcohols after reduction with  $\text{NaBH}_4$ . Enantioselectivities up to 99.5 ee were achieved at room temperature [28, 29].

The chiral dinuclear titanium complexes depicted in Scheme 9 have been utilized as catalysts for the DCR between *N*-benzyl and *N*-diphenylmethyl nitrones and



**Scheme 9** Cycloaddition catalyzed by bis-titanium chiral catalysts


acrolein or methacrolein. Interestingly, the 3,4-*endo* adduct was the major product. Several *N*-benzyl and *N*-diphenylmethyl nitrones with aromatic rings with different electronic properties, alkyl groups such as *tert*-butyl or cyclohexyl, and 1-cyclopentenyl or 1-methyl-1-propenyl have been investigated. In general, good yields and enantioselectivities up to 99% ee with a catalytic loading of 10 mol% were achieved [30–32].

The enantioselective catalytic 1,3-dipolar cycloaddition of linear or cyclic nitrones to enals was accomplished using the half-sandwich rhodium(III) complex ( $S_{\text{Rh}}, R_{\text{C}}$ )- $[(\eta^5\text{-C}_5\text{Me}_5)\text{Rh}\{(R)\text{-Prophos}\}(\text{H}_2\text{O})](\text{SbF}_6)_2$  as catalyst precursor [33, 34]. At  $-25^\circ\text{C}$ , quantitative conversions to the cycloadducts, with up to 95% ee, were achieved (Scheme 10). The intermediate with the dipolarophile coordinated to the rhodium has been isolated and completely characterized, including the X-ray determination of its molecular structure [33, 34].

The same rhodium precursor, ( $S_{\text{Rh}}, R_{\text{C}}$ )- $[(\eta^5\text{-C}_5\text{Me}_5)\text{Rh}\{(R)\text{-Prophos}\}(\text{H}_2\text{O})](\text{SbF}_6)_2$ , promotes the reaction between the nitrones *N*-benzylideneaniline *N*-oxide or 3,4-dihydroisoquinoline *N*-oxide with other enals different from methacrolein (Scheme 10). The cycloadducts were prepared with excellent regioselectivity, perfect *endo* selectivity, and enantiomeric excesses up to 94% [35].

Cationic chiral  $\text{Rh}_2^{5+}$  carboxamidates show catalytic activity for the DCR between diaryl nitrones and methacrolein (Scheme 11). Cycloadducts were obtained as 3,4-*endo*/3,5-*endo* mixtures with molar ratios ranging from 13/87 to 90/10. The enantioselectivity of the 3,4 isomers is higher than that of the 3,5 adducts and increases when the steric bulk of the ligand ester group increases [36].

Ruthenium complexes of formula  $[(\eta^6\text{-arene})\text{Ru}(\text{LL}^*)(\text{H}_2\text{O})](\text{SbF}_6)_2$  (arene =  $\text{C}_6\text{H}_6$ , *p*- $\text{MeC}_6\text{H}_4$ <sup>*i*</sup>Pr,  $\text{C}_6\text{Me}_6$ ; LL\* = bidentate chelate chiral ligand with PN, PP, or




**Scheme 10** DCR catalyzed by chiral half-sandwich complexes of rhodium

NN donor atoms) efficiently catalyze the DCR of methacrolein with a series of nitrones (Scheme 12). Typically, quantitative conversions are achieved at  $-25^{\circ}\text{C}$ , in a few hours. Perfect diastereoselectivity for the *endo* adducts is observed, but only low to moderate enantioselectivity is obtained [37].

### 3 DCR of Nitrones with Enals Catalyzed by Iridium Complexes


Methacrolein is the enal most studied in DCR catalyzed by iridium complexes.



**Scheme 11** DCR between diaryl nitrones and methacrolein catalyzed by cationic  $\text{Rh}_2^{5+}$  carboxamidates

### 3.1 DCR of Nitrones with Methacrolein


The enantioselective DCR of nitrones **I–V** to methacrolein catalyzed by  $(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}\{(R)\text{-Prophos}\}$  (Prophos = 1,2-bis(diphenylphosphino)propane) containing complexes (Scheme 13) was published in 2005 [34]. As a catalyst precursor, the aqua-complex  $(S_{\text{Ir}}, R_{\text{C}})\text{-}[(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}\{(R)\text{-Prophos}\}(\text{H}_2\text{O})](\text{SbF}_6)_2$  (**1**), prepared by treating the tris(solvento) complex [38] in acetone with an equimolar amount of  $(R)\text{-Prophos}$  in the presence of traces of water (Scheme 14), was employed. Complex **1** is prepared in a completely diastereoselective manner: only one of the two possible epimers at metal was spectroscopically detected, in solution from  $-90^\circ\text{C}$  to RT, and an X-ray determination showed that it was the *S* at metal isomer. Methacrolein displaces coordinated water from **1** affording, also diastereoselectively, the enal complex  $(S_{\text{Ir}}, R_{\text{C}})\text{-}[(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}\{(R)\text{-Prophos}\}(\text{methacrolein})](\text{SbF}_6)_2$  (**2**) (Scheme 15) [34]. The presence of an electron-withdrawing group on the dipolarophile (CHO) together with the coordination to the metal makes the DCR process a Sustmann's type I reaction and, accordingly, electronic density will be transferred from the nitronium to the methacrolein. In such a process, to achieve good yield, coordination of the dipole (nitronium) has to be avoided. Nitrones may coordinate metals through their oxygen atom and, in fact, while acyclic nitrones **I** and **II** do not displace methacrolein from complex **2**, the cyclic ones **III–V** readily substitute the coordinated dipolarophile even at a low temperature. Thus, for example,  $^{31}\text{P}$  NMR spectra of dichloromethane solutions of the methacrolein complex  $(S_{\text{Ir}}, R_{\text{C}})\text{-}[(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}\{(R)\text{-Prophos}\}(\text{methacrolein})](\text{SbF}_6)_2$  (**2**) remained unchanged for hours, in the  $-90^\circ\text{C}$  to RT range of temperatures, after the addition of 20 equivalents of nitronium **I**. However, immediately after addition, at  $-60^\circ\text{C}$ , of the same amount of the cyclic nitronium **V**, the  $^{31}\text{P}$  NMR spectrum showed the formation of the nitronium complex  $[(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}\{(R)\text{-Prophos}\}(\text{V})]^{2+}$ . In fact, the related nitronium complex  $[(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}\{(R)\text{-Prophos}\}(\text{III})](\text{SbF}_6)_2$  has been




**Scheme 12** DCR catalyzed arene-ruthenium complexes

isolated and characterized by addition of the corresponding cyclic nitron to the water complex  $(S_{Ir}, R_C)-[(\eta^5-C_5Me_5)Ir\{(R)\text{-Prophos}\}(H_2O)](SbF_6)_2$ , in the presence of molecular sieves, according to Scheme 16. When the nitron complex was prepared and isolated at  $-25^\circ\text{C}$ , the two epimers at metal, in ca. 50/50 molar ratio, were obtained. In dichloromethane, complete and irreversible epimerization occurs to the thermodynamically preferred  $S_{Ir}, R_C$  isomer. The latter was the sole isolated isomer when the preparation was carried out at room temperature.


In summary, when the iridium complex **2** is used as catalyst for the DCR reaction between cyclic nitrones and methacrolein, coordination of the nitron to the metal is likely, to the detriment of methacrolein coordination. As we will see later, to avoid



**Scheme 13** Major adducts for the DCR between nitrones I–V and methacrolein




**Scheme 14** Preparation of the catalyst Precursor (*S*<sub>Ir</sub>,*R*<sub>C</sub>)-1



**Scheme 15** Preparation of the methacrolein complex(*S*<sub>Ir</sub>,*R*<sub>C</sub>)-2





**Scheme 16** Preparation of nitrene complexes

**Table 1** Enantioselective DCR of methacrolein with nitrones I–V

Entry	Nitrene	t (h)	Yield (%)	3,4- <i>endo</i>	3,5- <i>endo</i>	ee (%)
1	<b>I</b>	10	100	82	18	95/85.5
2	<b>II</b>	24	78	2	98	–/93
3	<b>III</b>	15	100	–	100	86
4	<b>IV</b>	15	100	–	>99	92
5	<b>V</b>	15	100	–	100	93
6	<b>III</b>	16	75	–	>99	76

this negative effect, concentration of cyclic nitrones has to be maintained low during catalysis.

### 3.1.1 Catalytic Reactions

Complex **2** efficiently catalyzes the cycloaddition reaction of methacrolein with the nitrones **I–V**. Table 1 lists some results obtained. The reactions were performed in  $\text{CH}_2\text{Cl}_2$  in the presence of 4 Å molecular sieves, with 5 mol% of catalyst loading and a 1/140/20 catalyst/methacrolein/nitrene molar ratio. Typically, quantitative yields are obtained after a few hours at  $-25^\circ\text{C}$ . The acyclic nitrene **II** generates the less active system but, even so, 78% conversion was achieved after 24 h at  $-10^\circ\text{C}$  (entry 2). Enantiomeric excesses greater than 90% were achieved in most cases. A greater excess of methacrolein improves both rate and enantioselectivity (compare entry 6 with a catalyst/methacrolein/nitrene molar ratio 1/28/20 with entry 3 with a 1/140/20 molar ratio). To avoid undesired nitrene coordination, addition of the cyclic nitrones **III–V** was accomplished over 10 h.

Temperature variation slightly affected both product distribution and enantioselectivity, the latter smoothly increases as temperature decreases (Table 2). Thus, for example, the ee of the 3,4-*endo* adduct of the DCR between methacrolein and nitrene **I** gradually increases from 85 to 96% when temperature decreases from  $+5$  to  $-35^\circ\text{C}$  (entries 1–5). A similar trend was observed for the 3,5-adduct as well as for the DCR involving other nitrones (Table 2).

Ab initio calculations carried out by Tanaka and Kanemasa conclude that, under Lewis acid-catalyzed conditions, the formation of *endo*-cycloadducts is preferred and that the attack of nucleophilic nitrene oxygen should become more favored to occur at the  $\beta$ -position of the enal rendering 3,4-cycloadducts [39]. In the iridium

**Table 2** Effect of the temperature

Entry	Nitrone	T (°C)	t (h)	Yield (%)	3,4- <i>endo</i>	3,5- <i>endo</i>	ee (%)
1	<b>I</b>	+5	6	100	58	42	85/79
2	<b>I</b>	-5	7	100	71	29	91/81
3	<b>I</b>	-15	8	100	78	22	93.5/82
4	<b>I</b>	-25	10	100	82	18	95/85.5
5	<b>I</b>	-35	24	100	86.5	13.5	96/87
6	<b>II</b>	0	15	78	5	95	-/88
7	<b>II</b>	-10	24	78	2	98	-/93
8	<b>III</b>	-25	15	100	–	100	86
9	<b>III</b>	-35	24	55	–	100	91
10	<b>V</b>	-25	15	100	–	>99	93
11	<b>V</b>	-35	24	53	–	100	95


system that we are referring to, reactions occur with perfect *endo* selectivity. With respect to the regioselectivity, while the uncatalyzed reaction of nitrone **I** with methacrolein gives the 3,5-*endo* adduct, the major product of the catalyzed reaction is the 3,4-*endo* cycloadduct, in good agreement with Tanaka and Kanemasa calculations. On the other hand, cyclic nitrones **III–V**, which are more sterically demanding in their *endo*-approach to coordinated methacrolein, render 3,5-cycloadducts in all cases [23, 40].

### 3.1.2 The Catalytic Cycle

The catalytic cycle that has been proposed [34] is depicted in Scheme 17. At -25°C, the resting state of the catalyst is the [Ir]-methacrolein complex **2** that is the true catalyst. Its reaction with the nitrone is the key step of the productive cycle **A**: it is the rate- and enantioselectivity-determining step, at this temperature. For cyclic nitrones, path **B** becomes operative diminishing the concentration of the active species and, therefore, decreasing the rate. Lowering nitrone concentration (slow addition) hinders this undesired side reaction and, although it also slows down the rate of formation of the [Ir]-adduct intermediate, it favors the catalytic process overall. It has been experimentally shown that the catalytic rate increases when methacrolein concentration increases. Scheme 17 explains this effect because increasing this concentration favors the adduct elimination in path **A** and it recuperates the inactive metallic concentration, present as [Ir]-nitrone in path **B**, by shifting the [Ir]-nitrone/[Ir]-methacrolein equilibrium toward the enal complex, which restarts the cycle.

### 3.1.3 Origin of the Enantioselectivity

The ( $\eta^5$ -C<sub>5</sub>Me<sub>5</sub>)Ir{(R)-Prophos} system is very selective for the DCR of nitrones and methacrolein with enantioselectivity up to 96% ee. A few structural features of



**Scheme 17** Proposed catalytic cycle

the cation of the catalyst,  $[(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}\{(R)\text{-Prophos}\}(\text{methacrolein})]^{2+}$ , account for the encountered selectivity. First, in spite of  $(R)$ -Prophos being a  $C_1$  symmetric ligand, the  $S$  at metal epimer of the catalyst is formed with complete diastereoselectivity. Moreover, although, in some instances, epimerization at the metal in chiral half-sandwich transition metal complexes is a low-demanding energy process [41], this isomerization is not observed for this complex. Second, all the diffractometric and spectroscopic data support a  $\lambda$  conformation for the Ir–P–C–C–P five-membered metallacycle formed on coordination of the  $(R)$ -Prophos ligand. The bulky  $\text{C}_5\text{Me}_5$  ring constrains this conformation forcing the methyl substituent to occupy the less hindered pseudoequatorial position. This conformation, together with the  $S$  configuration at the metal, determines the chiral bias of the catalyst pocket in which catalysis takes place. Third, methacrolein coordinates to the metallic fragment in the  $E$  geometry and adopts an  $s$ -*trans* conformation [42–44]. This conformation, which is also preferred for uncomplexed methacrolein [45], is reinforced by coordination and retained in catalytic conditions. Fourth,  $\text{CH}/\pi$ -attractive interactions between the CHO aldehyde proton and the *pro-S* phenyl ring of the  $(R)$ -Prophos ligand (Fig. 3) establish the methacrolein rotamer around the M–O bond both in the solid state and in the solution. Overall, the geometry of the methacrolein is set and, in the chiral environment in which it is located, its *Re*-face becomes much more accessible to the nitrone than its *Si*-face, which is shielded by the phenyl rings of the  $(R)$ -Prophos ligand. This situation seems also to apply for the transition-state assembly of the DCR and accounts for the outcome of the catalytic reaction.






Fig. 3 Schematic view of the CH/ $\pi$  interactions in the cation of complex 2

### 3.2 DCR of Nitrones with Other $\alpha,\beta$ -Unsaturated Aldehydes

Complexes of the formula  $(S_{Ir},R_C)-[(\eta^5-C_5Me_5)Ir\{(R)\text{-Prophos}\}(enal)](SbF_6)_2$  (**3–8**) can be diastereoselectively prepared by addition of the appropriate enal to the water complexes  $(S_{Ir},R_C)-[(\eta^5-C_5Me_5)Ir\{(R)\text{-Prophos}\}(H_2O)](SbF_6)_2$  in the presence of 4Å molecular sieves (Scheme 18).

One interesting feature of the  $^1H$  NMR spectra of these complexes is the strong shielding of the aldehyde proton resonance after coordination. At  $-25^\circ C$  in  $CD_2Cl_2$  solution, this proton resonates in the range 6.93–7.50 ppm, about 2.3 ppm shifted to higher field with respect to the corresponding free enals. It has been shown that a comparable shift, measured in the related  $(S_{Ir},R_C)-[(\eta^5-C_5Me_5)Ir\{(R)\text{-Prophos}\}(methacrolein)](SbF_6)_2$  complex, is concomitant to the existence of CH/ $\pi$  interactions between the CHO proton and one of the phenyl rings of the Profos ligand [34]. Therefore, it is likely that these interactions are also operating in complexes **3–8**. In addition, NOESY experiments show enhancement patterns compatible only with an *s-trans* conformation for the coordinated enal and a  $\lambda$  conformation for the Ir–P–C–C–P five-membered metallacycle. In summary, the NMR data indicate that the conformation of both metallic fragment and enal are significantly hampered when the enal is coordinated to the metal into the chiral pocket defined by the  $(C_5Me_5)Ir\{(R)\text{-Prophos}\}$  moiety. Thus, they are excellent candidates to be tested as catalyst for DCR: the enals present a restricted conformation and, furthermore, they

**Scheme 18** Preparation of the enal complexes**Scheme 19** 1,3-Dipolar cycloaddition reactions**Table 3** Enantioselective DCR of enals with nitrones **I** and **V**

Entry	Catalyst	Nitronium	T (°C)	t (h)	Yield (%)	ee (%)
1	<b>3</b>	<b>I</b>	-25	16	100	90
2	<b>4</b>	<b>I</b>	-25	25	100	84.5
3	<b>5</b>	<b>I</b>	-25	72	96.5	92
4	<b>6</b>	<b>I</b>	-10	72	100	84.5
5	<b>6</b>	<b>I</b>	-25	72	55	94
6	<b>3</b>	<b>V</b>	-25	25	69	66
7	<b>4</b>	<b>V</b>	-25	25	100	70

are activated by coordination to the metal. In fact, most of them efficiently catalyze the cycloaddition reaction of the nitrones *N*-benzylidene-*N*-phenylamine *N*-oxide (**I**) and 3,4-dihydroisoquinoline *N*-oxide (**V**) to the corresponding enal (Scheme 19). Table 3 lists a selection of the results. Catalysts were prepared *in situ* by treating the aqua precursors  $(S_{Ir}, R_C)-[(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}\{(R)\text{-Prophos}\}(\text{H}_2\text{O})](\text{SbF}_6)_2$  with the corresponding enal in the presence of 4 Å MS. The reactions were performed in  $\text{CH}_2\text{Cl}_2$ , in the presence of 4 Å molecular sieves, with 10 mol% of catalyst loading and a 1/70/10 catalyst/enal/nitronium molar ratio. The 1-cyclohexen-1-carboxaldehyde (**7**) and *trans*-cinnamaldehyde (**8**) derivatives were not active. The cyclic

nitronone **V** was added slowly to avoid undesired nitronone coordination (see above). Perfect diastereoselectivity for the *endo* isomer was observed in all cases, and the 3,4 regioisomer was the sole adduct (nitronone **I**) or the major one (more than 90%, nitronone **V**). The ee values achieved ranged from 66 to 94%, enantioselectivity increases as temperature decreases (compare entries 4 and 5).

The absolute configuration of the major adduct obtained from the reaction of *trans*-crotonaldehyde with nitronone **I** was determined as the 3*S*,4*R*,5*S* isomer by derivatization with (*R*)-(-)- $\alpha$ -methylbenzylamine [45]. Accordingly, the *Re*-face of the coordinated *trans*-crotonaldehyde has to be much more accessible to the nitronone than its *Si*-face during the chirality induction step.

## 4 DCR of Nitrones with Methacrylonitrile Catalyzed by Iridium Complexes


The problem of the competitive coordination between the nitronone and the alkene can be circumvented by using alkenes with a good coordinating functionality such as a cyano group. Thus, the DCR between nitrones **IV** and **V** (see Scheme 13) and methacrylonitrile was carried out using the aqua-complex ( $S_{Ir},R_C$ )-[( $\eta^5$ -C<sub>5</sub>Me<sub>5</sub>)Ir{(*R*)-Prophos}(H<sub>2</sub>O)](SbF<sub>6</sub>)<sub>2</sub> (**1**) as the catalyst precursor [46, 47]. The reactions were performed in CH<sub>2</sub>Cl<sub>2</sub>, with 10 mol% of catalyst loading and a 1/70/20 catalyst/methacrylonitrile/nitronone molar ratio. In both cases, quantitative conversion to the 3,5-*endo*-cycloadduct are achieved but the system is not enantioselective (Table 4).

Interesting information about the catalytic systems was obtained by studying the stereochemistry of the reaction of the catalyst precursor **1** with methacrylonitrile. The water molecule of complex **1** is readily displaced by methacrylonitrile rendering complex [( $\eta^5$ -C<sub>5</sub>Me<sub>5</sub>)Ir{(*R*)-Prophos}(methacrylonitrile)](SbF<sub>6</sub>)<sub>2</sub> (**9**) as a mixture of the two possible epimers at metal, namely,  $R_{Ir},R_C$  and  $S_{Ir},R_C$ , in 34% diastereomeric excess in the former (Scheme 20). In acetone, at 50°C, the  $R_{Ir},R_C$  isomer slowly epimerizes to the thermodynamically preferred  $S_{Ir},R_C$  epimer. From the solution, pure samples of the latter can be isolated that have been employed as *stoichiometric* catalysts for the DCR between methacrylonitrile and nitrones **IV** and **V**.


Reaction of pure ( $S_{Ir},R_C$ )-**9** with nitrones **IV** or **V** followed by the addition of an excess of *n*-Bu<sub>4</sub>NBr (Scheme 21) gives the corresponding 3,5-*endo*-isoxazolidines in quantitative yield. Table 5 collects the ee values obtained for the two nitrones. Comparison of the results of Table 5 with those of Table 4 indicates that, whereas a mixture of epimers complex **9** (67%  $R_{Ir},R_C$ /33%  $S_{Ir},R_C$ , molar ratio) reacts with nitrones 2,3,4,5-tetrahydropyridine *N*-oxide (**IV**) or 3,4-dihydroisoquinoline

**Table 4** Catalytic DCR of methacrylonitrile

Entry	Nitronone	Yield	3,5- <i>endo</i> (%)	ee (%)
1	<b>IV</b>	91	100	1
2	<b>V</b>	100	100	0



**Scheme 20** Preparation of the methacrylonitrile complex **9**



**Scheme 21** Stoichiometric DCR between complex  $(S_{Ir,R_C})-9$  and nitrones **IV** and **V**

**Table 5** Stoichiometric reactions


Entry	Nitronium	Temp.	ee (%)
1	<b>IV</b>	RT	84
2	<b>V</b>	RT	74
3	<b>IV</b>	0°C	93.5

*N*-oxide (**V**) rendering regio- and diastereoselectively racemates of the corresponding 3,5-cycloadducts (Table 4), pure  $(S_{Ir,R_C})-9$  stoichiometrically catalyzes these reactions with 84 and 74% ee, respectively, (Table 5) at room temperature. At lower temperatures, greater ee are obtained (entry 3, 0°C, 93.5% ee).

The impressive increase in the ee values achieved in stoichiometric reactions compared to those in catalytic runs strongly indicates that the configuration at the metal correlates with the stereochemistry of the catalytic outcome: only if  $(R_{Ir,R_C})-9$  and  $(S_{Ir,R_C})-9$  diverge in enantioselection, it is possible to achieve zero ee even working with  $R_{Ir,R_C}/S_{Ir,R_C}$  mixtures and up to 93.5% ee using pure  $(S_{Ir,R_C})-9$  as catalyst.

## 4.1 Recycling Experiments

Pure  $(S_{Ir,R_C})-9$  is a highly enantioselective catalyst for the DC reaction between methacrylonitrile and nitrones **IV** and **V**, but only stoichiometric amounts of the



**Scheme 22** Recycling experiments

product can be prepared in high enantioselectivity. To increase the ratio adduct/catalyst without loss of ee, recycling experiments in repetitive batch mode have been carried out. Scheme 22 shows the three steps of the procedure developed. In the first step, 5 equivalents of nitron were added to diastereopure  $(S_{Ir}, R_C)$ -9. After the required reaction time, excess of nitron was extracted in  $\text{Et}_2\text{O}/\text{CH}_2\text{Cl}_2$  to avoid the simultaneous presence of nitron, alkene, and catalyst in the reaction medium. Adduct was dissociated from the iridium complex by adding 20 equivalents of alkene in the second step. Simultaneously, a  $(R_{Ir}, R_C, S_{Ir}, R_C)$ -9 mixture was formed. The adduct and the excess of nitrile were extracted in  $\text{Et}_2\text{O}/\text{CH}_2\text{Cl}_2$ , and from the extract, the adduct was recovered. In the third step, the  $(R_{Ir}, R_C, S_{Ir}, R_C)$ -9 mixture was allowed to epimerize to  $(S_{Ir}, R_C)$ -9, which restarts a further catalytic run. Following these procedure, in a second catalytic run,  $(S_{Ir}, R_C)$ -9 renders the cycloadduct in 93.5% yield and 92% ee. As it can be seen, both yield and ee are essentially maintained.<sup>1</sup>

<sup>1</sup>With the homologous rhodium complex  $(S_{Rh}, R_C)$ - $[(\eta^5\text{-C}_5\text{Me}_5)\text{Rh}\{(R)\text{-Prophos}\}(\text{methacrylonitrile})](\text{SbF}_6)_2$  as catalyst, up to ten consecutive catalytic runs have been performed for the reaction between methacrylonitrile and the nitron 3,4-dihydroisoquinoline *N*-oxide (**V**). Yield decreases from 97% (first run) to 84% (tenth run) and enantioselectivity remains essentially constant at 90% ee.



## 5 Conclusions

Diastereomerically pure iridium complexes of the formula  $[(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}\{(R)\text{-Prophos}\}(\text{activated alkene})](\text{SbF}_6)_2$  (activated alkene = enal, methacrylonitrile) are active, and selective catalysts for the DCR between one point binding activated alkenes and nitrones. Enals coordinate to the metal in a completely diastereoselective way with a restricted geometry. From the point of view of the selectivity, a key point in enal coordination is the establishment of CH/ $\pi$ -attractive interactions between the CHO aldehyde proton and one (*R*)-Prophos phenyl group. This interaction fixes the methacrolein rotamer around the M–O bonds and renders the system enantioselective.

The above-mentioned complexes are the sole iridium derivatives applied to DCR, and the cycloaddition of nitrones to enals or methacrylonitrile, the unique process studied. We think that iridium-based catalysts are underrepresented in 1,3-dipolar cycloaddition chemistry. For example, no iridium (I) systems have been developed to this end. It can be anticipated that the (bidentate ligand)Ir(I) fragment could be active (and stereoselective if chiral bidentate ligands are used) in DCR such as those involving azomethine ylides.

In summary, the development of 1,3-dipolar cycloadditions will continue during the next years and the potential of iridium-based catalysts in this field is far from being exhausted.

**Acknowledgments** The authors thank financial support from Ministerio de Ciencia e Innovación (MICINN, Spain) and FEDER (Grant CTQ2009/10303BQU) and MULTICAT, Consolider Ingenio-2010 (CSD2009-00050).

## References

1. Padwa A (1991) In: Trost BM, Fleming I, Semmelhack MF (eds) Intermolecular 1,3-dipolar cycloadditions, comprehensive organic chemistry, vol 4. Pergamon, Oxford, pp 1069–1109
2. Huisgen R (1984) In: Padwa A (ed) 1,3-dipolar cycloaddition chemistry, vol 1. Wiley, New York
3. Huisgen R (1968) *J Org Chem* 33:2291–2297
4. Firestone RA (1968) *J Org Chem* 33:2285–2290
5. Ess DH, Houk KN (2008) *J Am Chem Soc* 130:10198–10187
6. Schoenebeck F, Ess DN, Jones GO, Houk KN (2009) *J Am Chem Soc* 131:8121–8133
7. Xu L, Doubleday CE, Houk KN (2009) *Angew Chem Int Ed Engl* 48:2746–2748
8. Engels B, Christl M (2009) *Angew Chem Int Ed Engl* 48:7968–7970
9. Sustmann R (1971) *Tetrahedron Lett* 12:2717–2720
10. Sustmann R, Trill H (1972) *Angew Chem Int Ed Engl* 11:838–840
11. Houk KN, Sims J, Watts CR, Luskus LJ (1973) *J Am Chem Soc* 95:7301–7315
12. Houk KN (1984) In: Padwa A (ed) 1,3-dipolar cycloaddition chemistry, vol 2. Wiley, New York
13. Frederickson M (1997) *Tetrahedron* 53:403–425
14. Gothelf KV, Jørgensen KA (1998) *Chem Rev* 98:863–909
15. Gothelf KV, Jørgensen KA (2000) *Chem Commun* 1449–1458

16. Husinec S, Savic V (2005) *Tetrahedron Asymmetry* 16:2947–2061
17. Pandey G, Banerjee P, Gader SR (2006) *Chem Rev* 106:4484–4517
18. Pellissier H (2007) *Tetrahedron* 63:3235–3285
19. Stanley LM, Sibi MP (2008) *Chem Rev* 108:2887–2902
20. Badoiu A, Brinkmann Y, Viton F, Kündig EP (2008) *Pure Appl Chem* 80:1013–1018
21. Brandi A, Cardona F, Cicchi S, Cordero FM, Goti A (2009) *Chem Eur J* 15:7808–7821
22. Kissane M, Maguire AR (2010) *Chem Soc Rev* 39:845–883
23. Viton F, Bernardinelli G, Kündig EP (2002) *J Am Chem Soc* 124:4968–4969
24. Badoiu A, Bernardinelli G, Mareda J, Kündig EP, Viton F (2008) *Chem Asian J* 3:1298–1311
25. Mita T, Ohtsuki N, Ikeno T, Yamada T (2002) *Org Lett* 4:2457–2460
26. Ohtsuki N, Kezuka S, Kogami Y, Mita T, Ashizawa T, Ikeno T, Yamada T (2003) *Synthesis* 1462–1466
27. Kezuka S, Ohtsuki N, Mita T, Kogami Y, Ashizawa T, Ikeno T, Yamada T (2003) *Bull Chem Soc Jpn* 76:2197–2207
28. Shirahase M, Kanemasa S, Oderaotoshi Y (2004) *Org Lett* 6:675–678
29. Shirahase M, Kanemasa S, Hasegawa M (2004) *Tetrahedron Lett* 45:4061–4063
30. Kano T, Hashimoto T, Maruoka K (2005) *J Am Chem Soc* 127:11926–11927
31. Hashimoto T, Omote M, Kano T, Maruoka K (2007) *Org Lett* 9:4805–4808
32. Hashimoto T, Omote M, Hato Y, Kano T, Maruoka K (2008) *Chem Asian J* 3:407–412
33. Carmona D, Lamata MP, Viguri F, Rodríguez R, Oro LA, Balana AI, Lahoz FJ, Tejero T, Merino P, Franco S, Montesa I (2004) *J Am Chem Soc* 126:2716–2717
34. Carmona D, Lamata MP, Viguri F, Rodríguez R, Oro LA, Lahoz FJ, Balana AI, Tejero T, Merino P (2005) *J Am Chem Soc* 127:13386–13398
35. Carmona D, Lamata MP, Viguri F, Rodríguez R, Fischer T, Lahoz F, Dobrinovitch IT, Oro LA (2007) *Adv Synth Catal* 349:1751–1758
36. Wang Y, Wolf J, Zavalij P, Doyle MP (2008) *Angew Chem Int Ed Engl* 47:1439–1442
37. Carmona D, Lamata MP, Viguri F, Ferrer J, García N, Lahoz F, Martín ML, Oro LA (2006) *Eur J Inorg Chem* 3155–3166
38. White C, Thompson SJ, Maitlis PM (1977) *J Chem Soc Dalton Trans* 1654–1661
39. Tanaka J, Kanemasa S (2001) *Tetrahedron* 57:899–905
40. Ali SA, Khan JH, Wazeer MIM (1988) *Tetrahedron* 44:5911–5920
41. Brunner H (2001) *Eur J Inorg Chem* 905–912
42. Carmona D, Cativiela C, Elipe S, Lahoz FJ, Lamata MP, López-Ram de VÍu MP, Oro LA, Vega C, Viguri F (1997) *Chem Commun* 2351–2352
43. Kündig EP, Saudan CM, Bernardinelli G (1999) *Angew Chem Int Ed Engl* 38:1220–1223
44. Davenport AJ, Davies DL, Fawcett J, Garrat SA, Russell DR (2000) *J Chem Soc Dalton Trans* 4432–4441
45. Ishihara K, Qingzhi G, Yamamoto H (1993) *J Am Chem Soc* 115:10412–10413
46. Carmona D, Lamata MP, Viguri F, Rodríguez R, Lahoz F, Oro LA (2007) *Chem Eur J* 13:9746–9756
47. Carmona D, Lamata MP, Viguri F, Rodríguez R, Lahoz F, Fabra MJ, Oro LA (2009) *Tetrahedron Asymmetry* 20:1197–1205



# Index

## A

- 2-Acetylfulan, 88
- Alanine, 83
- Alcohols, 77
  - C-alkylation, 100
  - oxidation, 91
- Aldehydes, reduction, transfer
  - hydrogenation, 78
- Alkanes, activation, 7
  - cyclic, metathesis, 147
  - dehydrogenation, 1, 141
  - metathesis, 139, 145
- Alkenes, 77
  - diastereoselective reduction, 50
  - 1,10-disubstituted, 31, 56
  - hydroformylation, 108
  - tetrasubstituted, 31, 59
  - transfer hydrogenation, 82
  - trisubstituted, 31, 39
- 3-Alkenoyl-oxazolidinones, 213
- Alkyl groups, dehydrogenation, 147
- C/N-Alkylation, 77
- Alkyne–carbonyl reductive couplings, 111
- Alkynes, allylmetal surrogates, transfer
  - hydrogenation, 123
  - vinylmetal surrogates, Ir-catalyzed
    - hydrogenation, 110
- Allenes, allylmetal surrogates, hydrogenation, 115, 119
- Allyl metal complexes, 171
- Allylation, 107
- Allylic acetates, 124
- Allylic alcohols, allylic substitution, 201
  - redox rearrangement to chiral aldehydes, 51
- Allylic amines, trisubstituted, 114
- Allylic carbonates, 203
- Allylic esters, 203
- Allylic substitutions, 169
  - metal-catalyzed, 195
  - iridium-catalyzed, 174, 202
- Allylmetal surrogates, 124
- Amines, alkylation with alcohols, 96
- Amino oxazolines, 86
- Ammonia, allylic substitution, 200
  - monoallylation, 201
- Anderssons quadrant model, 38
- Aniline, methyl cinnamyl carbonate, 196
- Anilines, 191
- Anion effect, 33
- Arenes, borylation, 148
  - silylation, 152
- N-Aryl acetophenone imines, 24
- Aryl difluorosilanes, 152
- N-Aryl ketimines, acetophenone based, 71
- Aryl ketones, 86
- Aryl zinc reagents, allylic carbonates, 190
- Arylacetonitriles, 102
- 1-Aryl-2-methylalkynes, 123
- Aryl–aryl coupling, 158
- Arylboronate esters, 148
- Asymmetric catalysis, 11, 31, 169
- Asymmetric hydrogenation, 11, 31
- Aza-iridacyclopentene, 114

## B

- Barbituric acid
- BDPP, 13
- Benzophenone, 79
- Benzylamine, 198
- N-Benzylaniline, 83

- Benzyloxycarbamate, 199  
Bicyclooctadiene, 203  
Biferrocene diphosphine, 16  
BINAP, 13  
Binaphane, 14  
Binaphthol, 12  
Bis(diphenylphosphino)hexahydrofuro  
[3,2-b]furan, 16  
Bis(*tert*-butylmethylphosphino)ethane, 17  
Bisoxazolines, 86  
Boronic esters, 49  
Borylation, 139  
Bromoacetophenone, 88  
*t*-Butylformimidate synthesis, 161
- C**  
C–C bond formation, 101, 108, 154  
    Ir-catalyzed hydrogenation, 110  
    Ir-catalyzed transfer hydrogenation, 119  
C–C multiple bonds, 154  
C–H bond activation, 1, 7, 139, 162  
Carbamate nucleophiles, allylic substitution,  
199  
Carbamates, 191  
Carbenes, *N*-heterocyclic, 79  
Carbon dioxide, reduction, 84  
Carbonyl allylations, 116  
*Carduus crispus*, 25  
Catalysis, asymmetric, 11, 31  
Cesium carbonate, 54  
Chromenes, asymmetric hydrogenation, 64  
Cinnamaldehyde, 91  
Conjugate reduction, 52  
Crabtree catalyst, 33  
Crispine A, 25  
Cyanoesters, 102  
Cycloadditions, 1,3-dipolar, 209  
Cyclohexanone, 80, 83  
Cyclometalation, 184  
Cyclooctadiene, 141  
Cyclopentadienyl, 7, 141
- D**  
DCR, nitrones with alkenes, 212  
    enals, 212  
    Ir catalysis, 216  
    methacrolein, 217  
    methacrylonitrile, Ir catalysis, 225  
Dehydrogenation, 139  
    catalytic, 141  
Demethyl methoxycalamenene, 45  
Diallylamines, 192
- DIAPHOXs, 180  
Diastereoselectivity, 202  
1,8-Diazabicyclo[5.4.0]undec-7-ene (DBU),  
185  
Dibenzo[*a,e*]cyclooctatetraene, 205  
Dienes, allylmetal surrogates, transfer  
hydrogenation, 121  
Dihydrocinnamaldehyde, 93  
Diisopropyl-1,10-phenanthroline, 153  
Diisopropylamine, 99  
1,3-Dimethylbarbituric acid, 102  
DIOP, 13  
Diphenylphosphinates, 62  
Diphosphine ferrocene, 14  
Diphosphines, 12  
    bidentate, 13  
Diphosphinites, 12  
    ligands, bidentate, 19  
Diphosphites, 12  
    ligands, 19  
Diphosphonites, 12  
Di-*sec*-butyl-tetrafluorodisilane, 153  
Dopa, 32  
Dynamic kinetic resolution, 94
- E**  
Enals, 209  
Enamines, asymmetric hydrogenation, 65  
    asymmetric reduction, 66  
Enantioselectivity, 107  
Enol esters/ethers, 61  
Enol phosphinates, asymmetric  
hydrogenation, 63  
Enolates, 188  
Ethyl nitroacetate, 203  
Ethylbenzene, 83
- F**  
Farnesol, 47  
Fluorenyloxycarbamate, 199  
Fluorocinnamic acid, 48  
Frontier molecular orbital theory, 211  
Furanoside diphosphinite ligands, 19  
Furanoside phosphite–phosphinite, 20  
Furans, asymmetric hydrogenation, 64
- H**  
Heteroarenes, silylation, 154  
Heterocumulenes, 160  
Heterocycles, 169  
Hydroarylation, 139, 154, 156  
Hydrogen transfer, 78

- Hydrogenation, 1, 6, 107  
  asymmetric, 11, 31  
Hydrovinylation, 154, 157  
4-Hydroxycoumarins, 102
- I**  
Imines, 11, 31, 73, 77  
  asymmetric hydrogenation, 69, 73  
  transfer hydrogenation, 83  
  vinylation, hydrogenative, 115  
*N*-Iminopyridinium ylides, 70  
Indoles, *C*-allylation, 190  
  asymmetric hydrogenation, 65  
  borylation, 151  
Ir(PHOX), 33  
Iridacyclopropene, 114  
Iridium, hydrogenation catalysis, 4  
  N/P complexes, 31  
Iridium-BINAP, 13  
Iridium ferrocenyl diphosphine, 6  
Iridium hydride intermediates, 37  
Iridium PHOX hydride complexes, 35  
Iridium-phosphoramidite catalysts, 205  
Iridium(III) vinyl bis-acetylacetonate, 155  
Isoquinoline alkaloid, 25  
Isotope exchange, 1
- K**  
Kagan's chelating diop, 3  
Ketones, 77  
  asymmetric transfer hydrogenation, 85  
Ketonitriles, 102
- L**  
Lipase, 96
- M**  
Malonate esters, 102  
Malonate nucleophiles, 188  
Metal carbenoids, 162  
Metalacyclic catalyst, 184  
Metalacyclic iridium-phosphoramidite,  
  187, 196  
Methacrylonitrile, 209  
4-Methoxy-3-nitrobenzoic acid, 122  
Methyl *N*-acetamidoacrylate, 19  
Methyl *t*-butyl ether (MTBE), 160  
Methylcinnamic ester, 38  
Methylcinnamyl carbonate, 196  
Methylpyrimidines, 102  
Metolachlor, 1  
Monodentate phosphite, 23  
Monophosphorothioite, monodentate, 26
- Morpholine, 97  
Mutisianthol, 45
- N**  
Nitrenoids, 162  
Nitroalkanes, 102, 189  
*p*-Nitroanisole, 84  
Nitroarenes, reduction, 84  
Nitroindole, 66  
Nitrones, 209
- O**  
Olefins, alkyl-substituted, 46  
  fluorinated, 48  
  trisubstituted, 59  
Organoboron reagents, 111  
Organofluorides, 48  
Organotin reagents, 111  
Organotitanium reagents, 111  
Organozinc reagents, 111  
Osborn catalysts, 3  
Oxa-iridacyclic intermediates, 113  
2-Oxazolidinone, 199  
Oxidation, 77  
Oxindoles 102
- P**  
Pentamethyldisiloxane, 152  
1-Phenethyl alcohol, 91  
Phenylacetoneitrile, 102  
Phosphine–phosphoramidite ligands, 19, 22  
Phosphines, monodentate, 27  
Phosphinooxazolines (PHOX), 33, 177  
Phosphite–oxazoline ligands, 60  
Phosphite–phosphinite ligands, 19  
Phosphoramidate, 48  
Phosphoramidites, 12, 169  
  ligand, 23, 181, 185  
  modification, 184  
Phosphorothioite ligands, 23  
Phosphorus ligands, 11  
  bidentate, 13  
  monodentate, 23  
PHOX ligands, 33  
Pinacolone, 79  
Piperdines, 69  
Potassium triethylsiloxide (TESOK), 194  
2-Propanol, 83  
Propiophenone, 85  
Pseudopteroxazole, 45  
Pumiliotoxin C, 5  
Pybox ligands, 178  
Pyridines, asymmetric hydrogenation, 68

**Q**

- Quinolines, asymmetric hydrogenation, 68
  - Ir-catalyzed hydrogenation, 19
  - transfer hydrogenation, 84
- Quinoxalines, 23

**R**

- Recycling, 226
- Reduction, 77
- Roxaticin, 127

**S**

- Schrock catalysts, 3
- Segphos, 90
- Shilov chemistry, 7
- Silylation, 139
- SIPHOX, 54
- Spiro-phosphoramidite, 25
- Stoichiometric chemistry, 140
- Streptavidin, 88
- Sulfamic acid, 204
- Sulfonamides, 191
  - alkylation, 98

**T**

- Terpinene, 35
- Tetrahydroisoquinoline, 97
- Tetrakis[3,5-bis(trifluoromethyl)phenyl]borate, 6, 17, 34
- Tetrakis(pentafluorophenyl)borate, 34

- Tetrakis(perfluoro-*tert*-butoxy)aluminate, 34
- Tetralone, 86
- Thiochromene, 64
- Tishchenko reaction, 93
- Tocopherol, 47
- Toluenesulfonamide, 98
- Transfer hydrogenation, 77, 107
  - asymmetric, 77
- Triazabicyclo[4.4.0]dec-5-ene (TBD), 185
- Tricyclohexylphosphine, 5
- Trifluoroacetamide, 192
- Trifluoromethyl alkenes, 49
- N*-(Trifluoro-1-phenylethylidene)aniline, 17
- Tri-ortho-tolylphosphine, 23
- Triphenylphosphite, 174

**V**

- Vaska's complex, 2, 140
- Vinyl boronic esters, 49
- Vinyl ether alcohols/esters, asymmetric hydrogenation, 61, 62
- Vinylation, 107, 110, 123
- Vinylboronates, asymmetric reduction, 49
- Vinylfluorides, 48
- Vitamin E, 47

**W**

- Weinreb amide, 53
- Wilkinson catalyst, 2